

1979

## Age and Growth of White Crappie, *Pomoxis annularis* Refinesque, from a Flood-Created Pond in Mississippi County, Arkansas

Stephen A. Sewell  
*Arkansas State University*

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Terrestrial and Aquatic Ecology Commons](#)

---

### Recommended Citation

Sewell, Stephen A. (1979) "Age and Growth of White Crappie, *Pomoxis annularis* Refinesque, from a Flood-Created Pond in Mississippi County, Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 33 , Article 40.  
Available at: <http://scholarworks.uark.edu/jaas/vol33/iss1/40>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact [ccmiddle@uark.edu](mailto:ccmiddle@uark.edu), [drowens@uark.edu](mailto:drowens@uark.edu), [scholar@uark.edu](mailto:scholar@uark.edu).

## Arkansas Academy of Science

River, SE of Oxford (Sec. 30, R8W, T18N) (2); Bull Pen Creek, NW of Wiseman (Sec. 10, R8W, T18N) (1). SHARP Co.: Unnamed tributary of Big Creek (Sec. 33, R5W, T16N) (3).

On 15 October 1978, a single 118 mm specimen of *Ichthyomyzon castaneus* was collected from the Strawberry River at St. Hwy. 115 (Sec. 17, T16N, R3W), Lawrence County. The specimen was still in the process of transforming into an adult; however, several clues to its identity were noted including a well-developed intestine, developing bicuspid teeth in the inner circle of the oral disc and relatively large buccal funnel. Dr. George A. Moore, Oklahoma State University, kindly verified the identification.

Commercial fishermen have provided documentation of the presence of *Scaphirhynchus platyrhynchus* and *Polyodon spathula* near the confluence of the Strawberry and Black rivers where these two large river forms probably utilize the resources of the Strawberry River when conditions are favorable.

The collection of a single specimen of *Pimephales promelas*, undoubtedly a bait release, was taken from the headwaters of the river approximately 8 mi. S. W. of Salem, Fulton County, on 4 April 1975. This represents the only record of this species in the system.

An excellent collecting site in Lawrence County just upstream from the St. Hwy. 115 bridge (Sec. 17, T15N, R3W) 1 mi. N. of Jesus has yielded records of the additional seven species denoted in this paper. Six collections have been made to date from this locality. A total of eight specimens of *Hybopsis x-punctata* have been collected at this location over small gravel in the main current. The richness of the diversity of the Hwy. 115 collecting site is demonstrated by the collection of a total of 51 species, including 17 species of etheostomatine fishes.

Five percid fishes are added to the stream list. Nine specimens of *Cystallaria* (= *Ammocrypta*) *asprella* were taken in three collections at St. Hwy. 115. A total of four *Ammocrypta clara* was taken over sand habitats at the Hwy. 115 location and N. of the St. Hwy. 58 bridge near Poughkeepsie. Fifteen specimens of *Etheostoma histrio* have been collected from the St. Hwy. 115 collecting locality. Forty-seven specimens of the two cryptic species, *Percina ouachitae* and *P. uranidea*, have been taken to date from the St. Hwy. 115 locality on four occasions. Specimens of the Strawberry River *P. uranidea* were used in the original description of the snail darter, *Percina tanasi* Etnier, in Tennessee (Etnier, 1976, *Percina (Imostoma) tanasi*, a new percid fish from the Little Tennessee River, Tennessee, Proc. Biol. Soc. Washington, 88 (44):469-488). Thirteen specimens of *Percina shumardi* have also been taken from the Hwy. 115 location, including a large, 60.2 mm SL (71.1 mm TL) male specimen.

With the addition of these 12 species the Strawberry River is now documented to have 107 species. This richness in diversity is exemplary of the upland streams in the Ozark physiographic province and indeed, compares with the richest streams in North America.

HENRY W. ROBISON, Department of Biological Sciences, Southern Arkansas University, Magnolia, Arkansas 71753.

#### AGE AND GROWTH OF WHITE CRAPPIE, *Pomoxis annularis* RAFINESQUE, FROM A FLOOD-CREATED POND IN MISSISSIPPI COUNTY, ARKANSAS

White Crappie, *Pomoxis annularis* Rafinesque, originally were found in lakes, ponds, bayous, and slow-moving streams and rivers from eastern South Dakota to western New York and south in the Mississippi River and Gulf of Mexico drainage to Alabama and Texas. They have been introduced into other suitable waters (Carlander, 1977). In Arkansas, the white crappie occurs in all major rivers (Buchanan, 1973) and reservoirs (Ball, 1972), comprising a significant portion of the annual sport harvest in both (Morais, 1975). Many growth studies have been done involving white crappie directly or as a sympatric species, as evidenced by the bulk of data compiled on the species by Carlander (1977). This paper describes the age and growth of white crappie in Butterfly Hole, a flood-created impoundment of the Mississippi River.

Butterfly Hole is located approximately 2.2 km north of Tomato, section 7, T14N, R13E, Mississippi County, Arkansas, in an area almost entirely devoted to farming. Butterfly Hole was formed during the Fall-Winter flooding of 1974, by flood-stage waters of the Mississippi River. The hold covers approximately 1.5 surface acres with bottom depths ranging from 0.3 to 10.7 meters. Due to its depth, Butterfly Hole has never been dry since its creation. Many such water bodies are created with each flood period, but most are shallow and dry up during the summer months when temperatures average 91°F (Ferguson and Gray, 1971).

Fifty-four white crappie were collected to determine the growth characteristics of the species in Butterfly Hole, following a complete rotenone kill by the Arkansas Game and Fish Commission.

There is little evidence of sex differences in the growth of white crappie (Carlander, 1977). Therefore, no sexually dimorphic growth patterns were assumed to exist, and no such differentiations were attempted.

Scales for study were selected by the use of the "key scale" method suggested by Lagler (1956). Key scales were designated as those from an area approximately 10 back from the head and 5 down from the lateral line along the right side of the fish. Approximately 20 scales were taken from each fish. Total length in mm and total weight in grams were recorded for each fish upon capture.

Scale annuli counts were made by use of an American Optical dissecting microscope in conjunction with an American Optical Model #651, 7.5 volt light source. Distances were measured on each scale using a USDA metric planners rule from focus to each annulus, and from focus to scale margin along the anterior median of each scale.

The age determinations were made by counting the number of annuli. Since collection was made in late summer, after annulus formation (Hall et al., 1954), the age referred to herein represents the number of the last complete annulus.

The length (66-356 mm) - weight (5.5-718 g) relationship was  $\text{Log } W = 0.924 \text{ Log } L + 0.134$ . The regression coefficient of 0.924 was significantly different from 3.0 ( $t_{54} = 5.33$ ), indicating that the weight of the crappie did not increase as the cube of length. Figure 1 further illustrates the rapidly decreasing rate of weight gain of all white crappie within the hole, following impoundment (1974).

The coefficient of condition, K, was calculated for each of the crappie from the expression,  $K = \frac{W}{L^3} \times 100$ . The coefficient for the individual fish ranged from 1.45 to 1.95 with an average of 1.64. The average coefficient of the Butterfly Hole crappie was higher than that reported by Whitacre (1952) in Crab Orchard Lake, Illinois (0.67), and higher than that reported by Witt (1952) from Lake Norfolk, Missouri (1.32) and Lake Taneycomo, Missouri, (1.33). Average condition coefficients for the year classes are given in Table 1.

The total length (L) - scale radius (S) relationship for the Butterfly Hole crappie was  $L = 50.98 + 48.53S$  with a correlation coefficient (r) of 0.96. The average calculated lengths at the time of annuli formation are given in Table 2. Comparison of lengths of age-groups I, II, and III revealed no difference at the 0.01 level between the year classes 1975 through 1977. Therefore, the postimpoundment growth was the same for all these year classes.

## General Notes




Figure 1: True weight gain ( $W_2 - W_1$ ) white crappie collected from Butterfly Hole by year class (Ricker, 1975).

Table 1. Average coefficient of condition ( $K_{TL}$ ) for year classes 1972 through 1977.

Year Class	Average $K_{TL}$	Number of Fish
1972	1.59	1
1973	1.55	3
1974	1.51	6
1975	1.75	31
1976	1.58	11
1978	1.93	2

Table 2. The average calculated total lengths (mm) of white crappie from Butterfly Hole.

Year Class	1	2	3	4	5
1972	124	187	250	313	342
1973	104	165	223	265	296
1974	117	172	226	263	-
1975	107	156	200	-	-
1976	106	149	-	-	-
1977	95	-	-	-	-
Average	109	166	215	280	324
Number of fish	2	11	31	6	3

## LITERATURE CITED

- BALL, R. L. 1972. The feeding ecology of black and white crappie in Beaver Reservoir. Unpubl. MS Thesis, U. of Arkansas, Fayetteville, Ark. 181 p.
- BUCHANAN, T. M., 1973. Handbook of freshwater fisheries biology, Vol. II. Iowa State Univ. Press, Ames, Iowa. 431 p.
- FERGUSON, D. V. and I. L. GRAY. 1971. Soil survey of Mississippi County, Arkansas, U. S. Gov't Printing Office, Washington, D. C. 76 p.
- HALL, G. E., R. M. JENKINS and J. C. FINNELL. 1954. The influence of environmental factors on the growth of crappie in Oklahoma waters. Okla. Fish. Res. Lab. Rpt. 40:1-56.
- STEPHEN A. SEWELL, Dept. of Biology, Arkansas State University, State University 72467. (Present address: Soil Conservation Service, P.O. Box 276, Melbourne, Arkansas 72556).
- MORAIS, D. I. 1975. Estimates of sportfisherman use and harvest, Bull Shoals Lake, 1971-1974. Unpubl. Report, USFWS. 23 p.
- RICKER, W. E. 1975. Computation and interpretation of biological statistics for fish populations. Fish. Res. Bd. Canada Bull. #191: 203-233.
- WHITACRE, M. A. 1952. The fishes of Crab Orchard Lake, Illinois. Midwest Wildl. Conf. 14:1-41.
- WITT, A. J. 1952. Age and growth of the white crappie in Missouri. Doct. Dissert. U. of Missouri, Columbia, MO. 213 p.

ADDITIONS, DELETIONS, AND CORRECTIONS FOR THE ATLAS AND ANNOTATED LIST OF THE VASCULAR PLANTS OF ARKANSAS

Since the publication of the Atlas and Annotated List of the Vascular Plants of Arkansas (Smith, 1978, Student Union Bookstore, U. of Ark. at Fayetteville, Fayetteville, Ark., 592 pp.), a large number of new county records and new reports and several new state records and deletions have come to our attention. Over 800 new county records, several new reports, and several deletions are covered in a "Quick Copy" list available from the senior author. Approximately half of the new county records in the "Quick Copy" list have been found by Gwen Barber in her study of the flora of Franklin County. The new state records are presented in the following list. Voucher specimens are on deposit in the herbarium at Fayetteville.