

University of Arkansas, Fayetteville

ScholarWorks@UARK

Animal Science Undergraduate Honors Theses

Animal Science

5-2017

Genetic Polymorphisms of the Glucocorticoid Receptor and Interleukin-8 Receptor Genes are Related to Production Traits and Hair Coat Score in Crossbred Cattle

Avery Deaton

Follow this and additional works at: <https://scholarworks.uark.edu/anscuht>



Part of the [Molecular Genetics Commons](#), and the [Other Animal Sciences Commons](#)

Citation

Deaton, A. (2017). Genetic Polymorphisms of the Glucocorticoid Receptor and Interleukin-8 Receptor Genes are Related to Production Traits and Hair Coat Score in Crossbred Cattle. *Animal Science Undergraduate Honors Theses* Retrieved from <https://scholarworks.uark.edu/anscuht/13>

This Thesis is brought to you for free and open access by the Animal Science at ScholarWorks@UARK. It has been accepted for inclusion in Animal Science Undergraduate Honors Theses by an authorized administrator of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Genetic Polymorphisms of the Glucocorticoid Receptor and Interleukin-8 Receptor Genes are
Related to Production Traits and Hair Coat Scores in Crossbred Cattle

Avery Deaton

University of Arkansas

Abstract

The objective of this thesis was to identify polymorphisms in the glucocorticoid receptor (GR) and interleukin-8 receptor (CXCR2) genes and to associate genotypes between the above mentioned polymorphisms and production traits in crossbred cattle. The hypothesis was that polymorphisms will exist for GR and CXCR2 genes and will be linked to production traits. Glucocorticoid receptors have been positively associated with higher milk yields, lactose content, feed intake, and feed conversion rates. Interleukin-8 genes are part of the innate immune response and help with many aspects of female reproduction health, such as protecting the embryo from the maternal immune system during pregnancy. Despite these things, very little is known about how GR and CXCR2 gene polymorphisms affect phenotypes in cattle. Blood samples were collected from ninety-four crossbred cattle over a period of three years (2012, 2013, 2014) and the DNA was extracted, amplified, and sent to GeneSeek in Lincoln, Nebraska, to be analyzed and genotyped for single nucleotide polymorphisms (SNP). Phenotypic data was collected from the ninety-four crossbred cattle and analyzed alongside the genotypic results, including: cow pre-breeding BCS and weight, Julian calving date, calf birth weight, cow weaning BCS and weight, calf weaning weight, calf adjusted 205-day weight, cow efficiency, and HCS. Significant relationships were determined using t-tests. It is expected that SNPs will be found for the GR and CXCR2 genes and that these polymorphisms will be significantly related to the production traits in cattle. Scientists and breeders could manipulate these genes to produce cattle that are more efficient and possess more desirable production traits.

TABLE OF CONTENTS

Introduction and Literature Review	4
Glucocorticoids	4
Interleukin-8	5
Materials and Methods	6
Results and Discussion	7
T105G Polymorphisms	7
C777G Polymorphisms	7
Cow Age and Year	7
Hair Coat Score	8
Conclusion	8
References	9
Tables	
Table 1. Description of Hair Coat Scores.....	10
Table 2. Distribution of SNP in the bovine genes.....	11
Table 3. Main effects of GR (T105G) and CXCR2 (C777G) SNP on bovine productivity traits.....	12
Table 4. Main effects of cow age group and year on bovine productivity traits.....	13
Figures	
Figure 1. Genotypic effects of CXCR2 on average hair coat score over three consecutive years (2012-2014).....	14

Introduction and Literature Review

Glucocorticoids

Steroid hormones can be classified as anabolic, results in increased lean body mass, and catabolic, which facilitate metabolism. Glucocorticoids, specifically cortisol and corticosterone, are the primary catabolic steroid hormones which assists with homeostasis in the body by mobilization of glucose, and increased glucose synthesis through gluconeogenesis in the liver. In response to stress, the hypothalamic-pituitary-adrenal (HPA) axis is activated and glucocorticoids are synthesized and released by the adrenal cortex which triggers the fight or flight response. Specifically, when an animal becomes stressed, the hypothalamus releases corticotropin releasing hormone which triggers the anterior pituitary to secrete adrenocorticotropin hormone (ACTH) into the bloodstream. Circulating ACTH acts on the adrenal cortex which then releases glucocorticoids which circulate throughout the body. Intracellular glucocorticoid receptors (GR) are located in the cytoplasm, and nucleus which are members of the nuclear receptor subfamily of ligand-dependent transcription factors. The GR remains inactive until glucocorticoids bind to it. The GR then transports the glucocorticoid to the nucleus through the use of nuclear pores. Once bound to the DNA, the GR acts as a transcription factor and can either induce or repress certain target genes (Oakley & Cidlowski, 2013).

In an experiment with Brown Swiss cows, a polymorphism of the glucocorticoid receptor DNA-binding factor 1 (GRLF1) was identified and positively associated with milk yields and lactose percentages. Two SNP had previously been found that were associated with feed intake and feed conversion rates in cattle (Cecchinato et al., 2014). In a study done on meat quality

traits in male Nellore cattle, GR polymorphisms were found to be associated with various traits, including: glucocorticoid sensitivity, bone mineral density, body mass index, abdominal obesity, cholesterol, and lower concentrations of plasma cortisol (Poleti et al., 2014). This supported the theory that the GR is related to energy production in cattle (Cecchinato et al., 2014).

Interleukin 8

Chemokines are a family of small (8-10 kDa) chemotactic cytokines that help coordinate the movement of cells. They are grouped into four families based on their amino acid sequences: α , β , γ , and δ (Tizard, 2013). The β , or CXC, chemokines are further classified into two subgroups, ELR- and ELR+. These subgroups are based on the presence of the amino acid sequence glutamic acid-leucine-arginine, or ELR (Umasuthan et al., 2014). Interleukin-8 (IL-8) is a β chemokine that attracts and activates neutrophils for inflammatory and immune responses. It is produced by macrophages, and is often denoted as CXCL8 due to its structure and function as a ligand (Tizard, 2013). It is ELR+ and typically found in or associated with liver, acute lung injury, and atherosclerotic lesions. The most common receptor for IL-8 is CXCR2, which binds all ELR+ CXC chemokines. Recruitment of myeloid cells is facilitated by CXCR2 for inflammation sites in the liver, lungs, and atherosclerotic lesions (Olson & Ley, 2002). Different genotypes for CXCR2 have been linked to impaired neutrophil migration and increased occurrences of mastitis (Tizard, 2013). The CXCR family of genes including CXCR1 and CXCR2 have affinities for IL-8 that may result in altered animal form and(or) function.

Very little is known about the impacts of the polymorphisms of the GR and CXCR family of genes. If identified, scientists and breeders could begin to manipulate those genes for a desired phenotype. Cattle could be produced that are more efficient and possess more desirable

production traits. The objective of this study was to identify polymorphisms in the GR and CXCR2 genes, and associate a specific phenotype between these polymorphisms and production traits in crossbred cattle.

Materials and Methods

This study used samples from ninety-four crossbred Angus cows grazing mixed grass pastures over a period of three consecutive years (2012, 2013, 2014). All cows were on a fall calving schedule with a 100% calving rate during those three years. Blood samples were taken from the jugular vein and immediately put in ice. The samples were centrifuged and then buffy coat was extracted. Buffy coats were stored at -20° C until DNA analysis.

Genomic DNA was extracted from the buffy coats and plated into 96-well plates. These plates were then shipped to GeneSeek in Lincoln, Nebraska to be sequenced and SNP analysis. Genotyping for GR and CXCR2 SNP was performed using the Sequenom technique.

Hair coat scores (HCS) were recorded for all ninety-four cows during each of the three years (2012, 2013, 2014). Trained personnel determined HCS for each cow monthly and used a scale ranging from 1 to 5 (Table 1).

Dependent variables collected from the 94 crossbred cows included: cow pre-breeding BCS and weight, Julian calving date, calf birth weight, cow weaning BCS and weight, calf weaning weight, calf adjusted 205-day weight, cow efficiency, and HCS. The experimental unit was the cow. Significance of the genotypes and their relation to the mentioned phenotypic traits was determined using t-tests where significance was determined at $P < 0.05$.

Results and Discussion

The SNP T105 and C777G were identified for the GR and CXCR2 genes, respectively. Seventy-eight cows were homozygous dominant for the GR gene. Sixteen cows were heterozygous for the polymorphism and zero cows were homozygous recessive. The minor allele frequency was 8.5% in the population (Table 2). Of the ninety-four cows, seventy-nine were homozygous dominant for the CXCR2 gene. Twelve cows were heterozygous and two were homozygous for the recessive allele resulting in an 8.6% minor allele frequency (Table 2).

T105G Polymorphism

Cow pre-breeding weight was affected ($P=0.02$) by genotype. Cow weight at weaning and calf weight at weaning tended ($P=0.08$, $P=0.11$) to be affected by genotype. All other dependent variables were not affected by the genotype (Table 3).

C777G Polymorphism

Cow pre-breeding body condition score and pre-breeding weight tended ($P=0.06$, $P=0.08$) to be affected by CXCR2 genotype. Calf birth weight was affected ($P=0.0003$) by genotype. Calf weaning weight also was affected ($P=0.05$) by genotype. Cow body condition score at weaning and weight at weaning were affected ($P<0.05$) by genotype. Calf adjusted 205 day weight tended ($P=0.10$) to be affected by genotype (Table 3).

Cow Age and Year

Cow pre-breeding weight, calf birth weight, cow weaning BCS, cow weaning weight, calf weaning weight, and cow efficiency were all affected ($P<0.05$) by cow age group (Table 4).

Cow pre-breeding BCS, cow pre-breeding weight, Julian calving date, cow weaning weight, calf adjusted 205 day weight, and cow efficiency were affected ($P < 0.05$) by year (Table 4).

Hair Coat Score

Cow HCS was affected by year and month ($P < 0.0001$). Over the course of twelve months, HCS tended to be affected ($P = 0.07$) by the genotype C777G. By looking at a period of four months, HCS was affected ($P = 0.008$) by the genotype C777G (Figure 1).

Conclusion

The polymorphism T105G for the glucocorticoid receptor gene was found to be associated with production traits in crossbred cows. This SNP was related to cow pre-breeding weight and tended to affect cow weight at weaning and calf weaning weight. The interleukin-8 receptor gene, CXCR2, polymorphism C777G was shown to be associated with calf birth weight, calf weaning weight, cow BCS at weaning, and cow weight at weaning. When looking at a period over four months, it also affected HCS in cows. This study also proved that cow age and year has effects on production traits and HCS in cattle. Future research should be done in order to determine if these mutations are related to other traits in cattle and if they occur in great enough frequency to be useful for further genetic modifications.

References

- Cecchinato, A., Ribeca, C., Chessa, S., Cipolat-Gotet, C., Maretto, F., Casellas, J. and Bittante, G. (2014). Candidate gene association analysis for milk yield, composition, urea nitrogen and somatic cell scores in brown swiss cows. *Animal: an International Journal of Animal Bioscience*, 8.7, 1062-1070. doi:10.1017/S1751731114001098
- Gray, K. A., Smith, T., Maltecca, C., Overton, P., Parish, J. A., and Cassady, J. P. (2011). Differences in hair coat shedding, and effects on calf weaning weight and BCS among Angus dams. *Livestock Science*, 140, 68-71. <http://dx.doi.org.library.uark.edu/10.1016/j.livsci.2011.02.009>
- Oakley, R. H., & Cidlowski, J. A. (2013). The biology of the glucocorticoid receptor: new signaling mechanisms in health and disease. *J Allergy Clin Immunol*, 132(5), 1033–1044. doi: 10.1016/j.jaci.2013.09.007
- Poleti, M.D., DeRijk, R.H., Rosa, A.F., Moncau, C.T., Oliveira, P.S., Coutinho, L.L., Eler, J.P., Balieiro, J.C.C. (2014). Genetic variants in glucocorticoid and mineralocorticoid receptors are associated with concentrations of plasma cortisol, muscle glycogen content, and meat quality traits in male Nellore cattle. *Domestic Animal Endocrinology*, 51, 105-113. doi: 10.1016/j.domaniend
- Tizard, I. R. (1977). *Veterinary immunology (9th ed.)*. St. Louis, MO: Elsevier Saunders.
- Umasuthan, N., Wan, Q., Revathy, K. S., Whang, I., Noh, J. K., Kim, S., Park, M., Lee, J. (2014). Molecular aspects, genomic arrangement and immune responsive mRNA expression profiles of two CXC chemokine receptor homologs (CXCR1 and CXCR2) from rock bream, *Oplegnathus fasciatus*. *Fish & Shellfish Immunology*, 40, 304-318. doi: 10.1016/j.fsi.2014.06.006

Table 1. Description of hair coat scores (HCS)

HCS	Description
5	Full winter coat, 0% shed
4	Initial shedding
3	Half way shed
2	Almost shed
1	Slick summer coat, 100% shed

(Gray et al., 2011)

Table 2. Distribution of SNP in the bovine genes

Polymorphism ^a	Genotype distribution ^b			MAF ^c
	Homo	hetero	homo	
T105G	78	16	0	8.5
C777G	79	12	2	8.6

^aSingle nucleotide polymorphism occurred at the number indicated. First letter indicates the primary allele and the letter following the digits is the minor allele

^bNumber of cows that were homozygous for the primary allele (Homo), heterozygous (hetero), and homozygous for the minor allele (homo)

^cMinor allele frequency expressed as percent

Table 3. Main effects of GR (T105G) and CXCR2 (C777G) SNP on bovine productivity traits

Productivity Traits ^b	T105G				C777G			
	TT	TG	SEM	<i>P</i> -value	CC	CG	SEM	<i>P</i> -value
No. ^c	234	48	-	-	237	42	-	-
Pre-Breeding								
BCS	5.5	5.4	0.12	0.26	5.6	5.1	0.15	0.06
Weight, kg	527	500	8.31	0.02	528	493	12.5	0.08
Calving								
Julian	268	271	2.32	0.42	267	270	2.94	0.78
Birth Weight, kg	34	33	0.88	0.51	35	29	1.01	0.0003
Weaning								
BCS	5.0	4.9	0.08	0.61	5.0	4.7	0.10	0.04
Cow Weight, kg	497	476	8.94	0.08	500	456	13.5	0.05
Calf Weight, kg	198	189	4.45	0.11	199	179	6.35	0.05
Adj. 205 Wt., kg	206	206	4.20	0.94	208	192	6.04	0.10
Cow Efficiency	43	45	1.26	0.26	43	43	1.26	0.63

^aSingle nucleotide polymorphism occurred at the number indicated. First letter indicates the primary allele and the letter following the digits is the minor allele; SEM represents the mean standard error of least square means

^bBCS = Body Condition Score; Cow efficiency calculated by dividing the Adj. 205 Weight by cow weight at weaning

^cNo. = total number of records over three consecutive years (2012, 2013, 2014)

Table 4. Main effects of cow age group and year on bovine productivity traits

Productivity Traits ^b	Cow Age Group ^a					Year				
	≤ 3	4 - 10	≥ 11	SEM	P-value	2012	2013	2014	SEM	P-value
No. ^c	53	207	22	-	-	94	94	94	-	-
Pre-Breeding										
BCS	5.4 ^x	5.6 ^x	5.0 ^x	0.18	0.07	5.1 ^y	5.4 ^{xy}	5.6 ^x	0.14	0.002
Weight, kg	476 ^y	546 ^x	510 ^{xy}	14.5	0.0001	501 ^y	489 ^y	542 ^x	10.8	0.0001
Calving										
Julian Date	271 ^x	270 ^x	267 ^x	3.58	0.71	264 ^y	268 ^{xy}	275 ^x	3.11	0.02
Birth Weight, kg	30 ^y	34 ^x	31 ^{xy}	1.18	0.005	31 ^x	32 ^x	32 ^x	0.95	0.41
Weaning										
BCS	5.0 ^z	5.1 ^x	4.5 ^y	0.12	0.007	4.9 ^x	4.9 ^x	4.7 ^x	0.10	0.30
Cow Weight., kg	453 ^y	505 ^x	477 ^{xy}	16.4	0.002	462 ^{xy}	488 ^x	484 ^x	37.0	0.008
Calf Weight, kg	166 ^{xy}	216 ^x	186 ^x	8.10	0.0001	193 ^x	178 ^x	196 ^x	6.88	0.06
Adj. 205 Wt., kg	207 ^x	206 ^x	187 ^x	7.60	0.34	199 ^x	186 ^{xy}	214 ^x	6.12	0.0003
Cow Efficiency	48 ^x	41 ^y	40 ^{yz}	1.55	0.0004	45 ^x	39 ^{xy}	45 ^x	1.41	0.0008

^aCow age groups = ≤ 3 years of age, young; 4 – 10 years of age, adult; ≥ 11 years of age, mature adult

^bBCS = Body Condition Score; Cow efficiency calculated by dividing the Adj. 205 Weight by cow weight at weaning

^cNo. = total number of animals over three consecutive years (2012, 2013, 2014)




Figure 1. Genotypic effects of CXCR2 on average hair coat score over three consecutive years (2012 – 2014). Based on a scale of 1-5; 5- 0% shed, 1- 100% shed.