2015

Crepidostomum cornutum (Digenea: Allocreadiidae) from Midget Crayfish, *Orconectes (Procericambarus) nana* (Decapoda: Cambaridae), from Northwestern Arkansas

C. T. McAllister

Eastern Oklahoma State College, cmcallister@se.edu

W. F. Font

Southeastern Louisiana University

H. W. Robison

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Zoology Commons](#)

Recommended Citation

McAllister, C. T.; Font, W. F.; and Robison, H. W. (2015) "Crepidostomum cornutum (Digenea: Allocreadiidae) from Midget Crayfish, *Orconectes (Procericambarus) nana* (Decapoda: Cambaridae), from Northwestern Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 69 , Article 27.

DOI: <https://doi.org/10.54119/jaas.2015.6921>

Available at: <https://scholarworks.uark.edu/jaas/vol69/iss1/27>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

***Crepidostomum cornutum* (Digenea: Allocreadiidae) from Midget Crayfish, *Orconectes (Procericambarus) nana* (Decapoda: Cambaridae), from Northwestern Arkansas**

C.T. McAllister^{1*}, W.F. Font² and H.W. Robison³

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

Department of Biological Sciences, Southeastern Louisiana University, Hammond, LA 70402

³9717 Wild Mountain Drive, Sherwood, AR 72120

*Correspondence: cmcallister@se.edu

Running Title: Trematode from Midget Crayfish

The midget crayfish, *Orconectes (Procericambarus) nana* Williams is a diminutive species (maximum length = 3.0 cm) that is found in the Neosho River Basin of northwest Arkansas and northeast Oklahoma (Reimer 1963, Hayes 1973, Morehouse and Tobler 2013). In addition, *O. nana* has been reported in the Illinois River (Bergey et al. 2005), and into the White River drainage (Prairie Creek) of Arkansas (C. Taylor, *pers. comm.*). In Arkansas, it has been collected from sites in Benton and Washington counties. The midget crayfish is found in clear-flowing, permanent small creeks and larger streams with substrate consisting of limestone gravel and cobble (Williams 1952, 1954). It was once thought to consist of two subspecies (Williams 1952, Hayes 1973) but a recent mtDNA study by Dillman et al. (2010) supported full species status originally suggested by Hobbs (1972) and Fitzpatrick (1987) for both *O. nana* and the Neosho midget crayfish, *O. macrus* Williams. This species' habitat is under constant threat from agriculture, road construction and urbanization, causing sedimentation and water pollution, in addition to construction of dams. Thus, this species has a State Heritage rank of S2 (imperiled) by NatureServe (2014) and has been assessed as vulnerable by the American Fisheries Society (Taylor et al. 2007).

Although information is available on the ecology of *O. nana* (see citations above), nothing, to our knowledge, has been published on any parasite of this crayfish. Here, we report a trematode parasite from *O. nana* from Arkansas.

Nine adult *O. nana* were collected on 19 November 2011 with a dipnet by overturning rocks at Flint Creek (Illinois River drainage) off St. Hwy. 59S, 0.5 km S of Gentry, Benton County (36.242699°N, 94.487472°W). In addition, on the same date, 7 adult *O. macrus* were taken with a dipnet from a tributary of Spavinaw Creek off St. Hwy 102, SE of Maysville, Benton County (36.364594°N, 94.551124°W).

Specimens were placed in aerated creek water and transported to the laboratory within 24 hr. They were killed by immersion in a concentrated Chloretone solution and their antennal glands, hepatopancreas and heart were placed in Petri dishes containing 0.6% w/v saline and examined for encysted parasites. Metacercariae were carefully teased from antennal gland cysts using insulin needles and fixed in hot ethanol, stained with acetocarmine and mounted in Canada balsam. Parasites were deposited in the United States National Parasite Collection (USNPC), Beltsville, Maryland, and vouchers of crayfish were deposited in the Henderson State University (HSU) Collection, Arkadelphia.

None of the *O. macrus* were found to be infected, however, 2 of 9 (22%) *O. nana* were each found to be harboring one metacercaria (USNPC 105277–105278) in their antennal glands that fit the description of *Crepidostomum cornutum* (Osborn, 1903) Stafford, 1904, given by Cheng (1957). Morphological features of these 2 metacercariae are as follows: genital pore slightly posterior to cecal bifurcation; anterior extent of vitellaria at genital pore; posterior extent of vitellaria near posterior end; prostate in anterior one-half of cirrus sac; posterior extent of cirrus sac at anterior margin of anterior testis; prostate in anterior two-thirds of cirrus sac. Measurements are as follows (in μm , length $\times$ width): body, 1,872–2,541 $\times$ 474–498; oral sucker, 269–284 $\times$ 316–395; pharynx, 98–117 $\times$ 82; ventral sucker, 222–254 $\times$ 230–242; ovary, 144–148 $\times$ 117–137; anterior testis, 234–250 $\times$ 203–238; posterior testis, 226–242 $\times$ 211–234.

Crepidostomum cornutum has been reported previously in various crayfish (*Cambarus* and *Orconectes* spp.) from Illinois, Kansas, Louisiana, Michigan, Minnesota, Mississippi, New York, Oklahoma, Virginia, and Wisconsin, and Ontario and Quebec, Canada (Faust 1918, Hopkins 1934, Henderson 1938). The life cycle involves cercaria in

Figure 1. *Crepidostomum cornutum* metacercariae. A. Specimen showing ovary (O) and testes (T). B. Close-up of anterior end of specimen showing lappets (L) characteristic of the genus.

first intermediate freshwater bivalves (*Musculium* and *Sphaerium* spp.), metacercaria encyst in second intermediate host crayfishes, and adults are found in the pyloric ceca and intestinal tract of fish definitive hosts (Ameel 1937, Cheng 1957). There are many fishes reported as hosts including those in the genera *Ambloplites*, *Amia*, *Anguilla*, *Carassius* (experimental), *Gasterosteus*, *Ictalurus*, *Lepomis*, *Micropterus*, *Morone*, *Notemigonus*, *Perca*, *Pomoxis*, *Pylodictus*, *Salmo*, and *Salvelinus* (Hoffman 1999).

Previous reports of *C. cornutum* from Arkansas include specimens from game fishes, including Channel Catfish (*Ictalurus punctatus*), Warmouth (*Lepomis gulosus*), Bluegill (*Lepomis macrochirus*), Spotted Bass (*Micropterus punctulatus*), Largemouth Bass (*Micropterus salmoides*) and White Crappie (*Pomoxis annularis*) (Becker et al. 1966, Becker and Houghton 1969, Becker and Cloutman 1975, Cloutman 1975, Kilambi and Becker 1977). In addition, McAllister et al. (2014) recently reported *Crepidostomum cooperi* Hopkins, 1931 from Banded

Sculpin (*Cottus caroliniae*) from Flint Creek.

In summary, we document, to our knowledge, the first parasite ever reported from *O. nana*. Turner (1999) reported *Alloglossidium cardicolum* (Corkum and Turner) Smythe and Font from White River crayfish, *Procambarus acutus* from Arkansas. In addition, Turner (2006) reported *Allocorrigia filiformis* Turner and Corkum from red swamp crayfish, *Procambarus clarkii* from Arkansas, and Turner (2009) found *Alloglossidium dolandi* (Turner and McKeever) Smythe and Font in *P. acutus* from Arkansas. The only other previous report on Arkansas crayfish parasites was by McAllister et al. (2011) who reported metacercaria of *Alloglossidium corti* (Lamont) Van Cleave and Mueller from red-spotted stream crayfish, *Orconectes acares* and western painted crayfish, *Orconectes palmeri longimanus*. As Arkansas supports approximately 58 species of crayfishes, additional surveys should increase our knowledge of their parasites, of which we know very little about.

Acknowledgments

The Arkansas Game and Fish Commission issued Scientific Collecting Permits to CTM and HWR. Dr. Renn Tumblison (HSU) and Pat Pilitt (USNPC) provided expert curatorial assistance.

Literature Cited

- Ameel DJ.** 1937. The life history of *Crepidostomum cornutum* (Osborn). Journal of Parasitology 23:218-220.
- Becker DA and DG Cloutman.** 1975. Parasites of selected game fishes of Lake Fort Smith, Arkansas. Proceedings of the Arkansas Academy of Science 29:12-18.
- Becker DA, RG Heard and PD Holmes.** 1966. A pre-impoundment survey of helminth and copepod parasites of *Micropterus* spp. of Beaver Reservoir in northwest Arkansas. Transactions of the American Fisheries Society 95:23-34.
- Becker DA and WC Houghton.** 1969. A survey of the helminth parasites of selected game fishes of Lake Fort Smith, Arkansas. Proceedings of the Arkansas Academy of Science 23:110-117.
- Bergey EA, SN Jones and DB Fenolio.** 2005. Surveys and studies of Oklahoma crayfish and the grotto salamander. Final Report to the Oklahoma Biological Survey, University of Oklahoma, Tulsa, Oklahoma, August 2005. 26 p.

Trematode from Midget Crayfish

- Cheng TC.** 1957. A study of the metacercaria of *Crepidostomum cornutum* (Osborn, 1903), (Trematoda: Allocreadiidae). Proceedings of the Helminthological Society of Washington 24:107-109.
- Cloutman DG.** 1975. Parasite community structure of Largemouth Bass, Warmouth, and Bluegill in Lake Fort Smith, Arkansas. Transactions of the American Fisheries Society 104:277-283.
- Dillman CB, BK Wagner and RM Wood.** 2010. Phylogenetic estimation of species limits in dwarf crayfishes from the Ozarks: *Orconectes macrus* and *Orconectes nana* (Decapoda: Cambaridae). Southeastern Naturalist 9 (special issue 3):185-198.
- Faust EC.** 1918. Studies on American Stephanophialinae. Transactions of the American Microscopical Society 37:183-198.
- Fitzpatrick JF Jr.** 1987. The subgenera of the crawfish genus *Orconectes* (Decapoda: Cambaridae). Proceedings of the Biological Society of Washington 100:44-74.
- Hayes WA III.** 1973. Occurrence of the crawfish *Orconectes nana macrus* Williams (Decapoda, Astacidae) in Oklahoma. Proceedings of the Oklahoma Academy of Science 53:111.
- Henderson HE.** 1938. The cercaria of *Crepidostomum cornutum* (Osborn). Transactions of the American Microscopical Society 57:167-172.
- Hobbs HH Jr.** 1972. Crayfishes (Astacidae) of North and Middle America. Biota of Freshwater Ecosystems Identification Manual No. 9, US Government Printing Office, Washington, DC. 173 p.
- Hoffman GL.** 1999. Parasites of North American freshwater fishes. Second Ed. Ithaca (NY): Comstock Publishing Associates. 539 p.
- Hopkins SH.** 1934. The papillose Allocreadiidae. Illinois Biological Monographs 13:45-124.
- Kilambi RV and DA Becker.** 1977. Population dynamics and species diversity of ichthyoparasitofauna of the Buffalo National River. Arkansas Water Resources Center, University of Arkansas 48:1-73.
- McAllister CT, MB Connior, WF Font and HW Robison.** 2014. Helminth parasites of the Banded Sculpin, *Cottus carolinae* (Scorpaeniformes: Cottidae), from Northern Arkansas, U.S.A. Comparative Parasitology 81:203-209.
- McAllister CT, HW Robison and WF Font.** 2011. Metacercaria of *Alloglossidium corti* (Digenea: Macroderoididae) from 3 species of crayfish (Decapoda: Cambaridae) in Arkansas and Oklahoma, U.S.A. Comparative Parasitology 78:382-386.
- Morehouse RL and M Tobler.** 2013. Crayfishes (Decapoda: Cambaridae) of Oklahoma: Identification, distributions, and natural history. Zootaxa 3717:101-157.
- NatureServe.** 2014. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>. (Accessed: 9 February 2015).
- Reimer RD.** 1963. The crawfish of Arkansas. [MS thesis]. Fayetteville (AR): University of Arkansas. 170 p.
- Taylor CA, GA Schuster, JE Cooper, RJ DiStefano, AG Eversole, P Hamr, HH Hobbs Jr, et al.** 2007. A reassessment of the conservation status of crayfishes of the United States and Canada after 10+ years of increased awareness. Fisheries 32: 272-389.
- Turner HM.** 1999. Distribution and prevalence of *Alloglossoides caridicola* (Trematoda: Macroderoididae), a parasite of the crayfish *Procambarus acutus* within the state of Louisiana, U.S.A., and into adjoining states. Journal of the Helminthological Society of Washington 66:86-88.
- Turner HM.** 2006. Distribution and prevalence of *Allocorrigia filiformis* (Trematoda: Dicrocoeliidae), a parasite of the crayfish *Procambarus clarkii*, within the state of Louisiana and the lower Mississippi River Valley, U.S.A. Comparative Parasitology 73:274-278.
- Turner HM.** 2009. Additional distribution and prevalence records for *Alloglossidium dolandi* (Digenea: Macroderoididae) and a comparison with the distribution of *Alloglossidium caridicolum*, parasites of procambarid crayfish, within the coastal plains of the southeastern United States. Comparative Parasitology 76:283-286.
- Williams AB.** 1952. Six new crayfishes of the genus *Orconectes* (Decapoda: Astacidae) from Arkansas, Missouri and Oklahoma. Transactions of the Kansas Academy of Science 55: 330-351.
- Williams AB.** 1954. Speciation and distribution of the crayfishes of the Ozark Plateaus and Ouachita Provinces. University of Kansas Science Bulletin 36:803-918.