2014

The Introduced Dirt-Colored Seed Bug, *Megalonotus sabulicola* (Hemiptera: Rhyparochromidae): New for Arkansas

S. W. Chordas III
Ohio State University

C. T. McAllister
Eastern Oklahoma State College, cmcallister@se.edu

H. W. Robison

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Entomology Commons](#)

Recommended Citation

Chordas, S. W. III; McAllister, C. T.; and Robison, H. W. (2014) "The Introduced Dirt-Colored Seed Bug, *Megalonotus sabulicola* (Hemiptera: Rhyparochromidae): New for Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 68 , Article 22.

DOI: <https://doi.org/10.54119/jaas.2014.6817>

Available at: <https://scholarworks.uark.edu/jaas/vol68/iss1/22>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

The Introduced Dirt-Colored Seed Bug, *Megalonotus sabulicola* (Hemiptera: Rhyparochromidae): New for Arkansas

S.W. Chordas III¹, C.T. McAllister^{2*}, and H.W. Robison³

¹Center for Life Sciences Education, The Ohio State University, 260 Jennings Hall, 1735 Neil Avenue, Columbus, OH 43210

²Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

³9717 Wild Mountain Drive, Sherwood, AR 72120

*Correspondence: cmcallister@se.edu

Over the past decade, we have published several new records of Hemiptera from Arkansas in this journal (Chordas et al. 2005, 2011, Chordas and Kovarik 2008a, 2008b, Chordas and Kremers 2009). We now document the introduced dirt-colored seed bug, *Megalonotus sabulicola* (Thomson, 1870) for the first time from Arkansas.

On 20 April 2013, CTM and HWR collected various hemipterans at a watershed 8.0 km N of Huntsville off St. Hwy. 23W at Withrow Springs State Park (36.156195°N, 93.733369°W), Madison County (Fig. 1). Specimens were collected with a standard insect sweep net or an aquatic dip net and placed in individual vials of 70% ethanol. Bugs were forwarded to the senior author for identification, photography, and deposition of voucher specimens into the C.A. Triplehorn Insect Collection (The Ohio State University, Columbus, Ohio). Ashlock and Slater (1988), Maw et al. (2000), Scudder and Footitt (2006), and Wheeler (1992) were used as distribution references. Wheeler (1989, 1992) were used as taxonomic and identification references.

Figure 1. Location of study site (dot) in Madison County, Arkansas.

A single noteworthy specimen of *M. sabulicola* (Fig. 2) was collected in emergent vegetation. Our specimen matches the identifying characters and figure in Wheeler (1989). The bug possesses a mottled brown-testaceous hemelytra with a piceous head, pronotum and scutellum (Fig. 2A). It also has a prominent spine on the lateral, ventral aspect of the swollen profemur (Fig. 2B) and an antennal segment II almost all yellow with black at the distal edge (Fig. 2C). The tibia are yellowish with both the meso- and metafemurs are yellow proximally and piceous distally (Figs. 2A-B).

Figure 2. A. Dorsal habitus of *Megalonotus sabulicola*. Bar = 1 mm. B. Ventrolateral profemur spines (arrows). C. Antennae.

Megalonotus sabulicola was introduced from Europe in the early 1900's and has been reported to feed on seeds of knapweed, *Centaurea* spp. (Wheeler 1989). Documented mainly from both Atlantic and Pacific coastal regions of North America (see distribution herein), *M. sabulicola* was unexpected for Arkansas.

Tabulation of 21 records for *M. sabulicola* (Fig. 4) are as follows: USA (18 records): Arkansas [New Record], California, Connecticut, Delaware, Idaho, Massachusetts, Maryland, Michigan, New Jersey, New

York, North Dakota, Oregon, Pennsylvania, Rhode Island, Utah, Virginia, West Virginia, Washington. Canada (3 records): British Columbia, Ontario, Quebec.

Our Arkansas record represents a significant range extension of >1,000 km from the closest reported location (western Virginia) for this species. Previous records were primarily concentrated along the eastern and western portions of the U.S. and Canada with a few additional records in the Great Lakes region and a report from North Dakota (Fig. 3).

In summary, more data and collections are needed to determine the persistence of this bug in Arkansas. How and when it arrived in the state would be pure speculation at this point. For detailed biological information on *M. sabulicola*, see Wheeler (1989).

Figure 3. Distribution of *Megalonotus sabulicola* north of Mexico. Light shading (previous records); dark shading (new state record).

Acknowledgments

We thank the Arkansas Game and Fish Commission for Scientific Collecting Permits issued to CTM and HWR.

Literature Cited

Ashlock PD and A Slater. 1988. Family Lygaeidae Schilling, 1829. In Henry TJ and RC Froeschner, editors. Catalog of the Heteroptera, or true bugs, of Canada and the continental United States. New York (NY): E.J. Brill. p 167-245.

Chordas SW III, EG Chapman, PL Hudson, MA Chriscinske and RL Stewart. 2002. New Midwestern state records of aquatic Hemiptera (Corixidae; Notonectidae). Entomological News 113:310-314

Chordas SW III and PW Kovarik. 2008a. Two Coreidae (Hemiptera), *Chelinidea vittiger* and *Anasa armigera*, new for Arkansas, U.S.A. Journal of the Arkansas Academy of Science 62:145-146.

Chordas SW III and PW Kovarik. 2008b. Two Lygaeoidea (Hemiptera), *Ischnodemus slossonae* and *Cryphula trimaculata*, new for Arkansas, U.S.A. Journal of the Arkansas Academy of Science 62:147.

Chordas SW III and J Kremers. 2009. Backyard bug” collecting results in 6 new state records for Arkansas, U.S.A. Journal of the Arkansas Academy of Science 63:177-179.

Chordas SW III, HW Robison, EG Chapman, BG Crump and PW Kovarik. 2005. Fifty-four state records of true bugs (Hemiptera: Heteroptera) from Arkansas. Journal of the Arkansas Academy of Science 59:43-50.

Chordas SW III, R Tumilson, HW Robison and J Kremers. 2011. Twenty three true bug state records for Arkansas, with two for Ohio, U.S.A. Journal of the Arkansas Academy of Science 65:153-159.

Jansson A. 2002. New records of Corixidae (Heteroptera) from northeastern USA and eastern Canada, with one new synonymy. Entomologica Fennica 13:85-88.

Maw HEL, RG Foottit, KGA Hamilton and GGE Scudder. 2000. Checklist of the Hemiptera of Canada and Alaska. Ottawa, Ontario (Canada): NRC Research Press. 220 p.

Scudder GGE and RG Foottit. 2006. Alien true bugs (Hemiptera: Heteroptera) in Canada: composition and adaptations. The Canadian Entomologist 138:24-51.

Wheeler Jr AG. 1989. *Megalonotus sabulicola* (Heteroptera: Lygaeidae), an immigrant seed predator of *Centaurea* spp. (Asteraceae): Distribution and habitats in eastern North America. Proceedings of the Entomological Society of Washington 91:538-544.

Wheeler Jr AG. 1992. Holarctic insects adventive in Michigan: New and additional records (Homoptera, Heteroptera, Coleoptera, Neuroptera). Great Lakes Entomologist 25:99-106.