2013

Ectoparasites of Sciurid Rodents in Arkansas, Including New State Records for *Neohaematopinus* spp. (Phthiraptera: Anoplura: Polyplacidae)

C. T. McAllister

Eastern Oklahoma State College, cmcallister@se.edu

M. B. Connior

South Arkansas Community College

L. A. Durden

Georgia Southern University

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Animal Diseases Commons](#), and the [Zoology Commons](#)

Recommended Citation

McAllister, C. T.; Connior, M. B.; and Durden, L. A. (2013) "Ectoparasites of Sciurid Rodents in Arkansas, Including New State Records for *Neohaematopinus* spp. (Phthiraptera: Anoplura: Polyplacidae)," *Journal of the Arkansas Academy of Science*: Vol. 67 , Article 34.

DOI: <https://doi.org/10.54119/jaas.2013.6729>

Available at: <https://scholarworks.uark.edu/jaas/vol67/iss1/34>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Ectoparasites of Sciurid Rodents in Arkansas, Including New State Records for *Neohaematopinus* spp. (Phthiraptera: Anoplura: Polyplacidae)

C.T. McAllister^{1*}, M.B. Connior², and L.A. Durden³

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

²Health and Natural Sciences, South Arkansas Community College, 300 S. West Ave., El Dorado, AR 71730

³Department of Biology, Georgia Southern University, Statesboro, GA 30458

*Correspondence: cmcallister@se.edu

Running Title: Ectoparasites of Sciurid Rodents in Arkansas

Compared to its surrounding states, little is known about the ectoparasites (bugs, fleas, flies, lice, mites, ticks) infesting the wild mammals of Arkansas (AR) (Bequaert 1946, Sealander and Young 1954, Tugwell and Lancaster 1962, Schiefer and Lancaster 1970, Lancaster 1973, Whitaker and Wilson 1974, Ellis 1975, Price et al. 1982, Steward et al. 1986, Durden and Musser 1994). However, some studies specifically on ectoparasites of mammals have been published. For example, Richardson et al. (1994) reported records of various ectoparasites from raccoons (*Procyon lotor*) in the state, and Elrod et al. (1996) reported ectoparasites from the endemic Ozark pocket gopher (*Geomys bursarius ozarkensis*). Here, we report information on a flea, 3 ticks, a chigger, and 3 sucking lice species from sciurid rodents, including the first report of the sucking lice, *Neohaematopinus sciurinus* (Mjöberg) and *Neohaematopinus sciuropteri* (Osborn) from the state.

Various sciurid rodents, including 2 eastern gray squirrels (*Sciurus carolinensis*) and their nests, 2 fox squirrels (*Sciurus niger*), and a single eastern chipmunk, (*Tamias striatus*) from Marion Co., and 10 southern flying squirrels (*Glaucomys volans*) from Union Co. were collected using live box traps from Dec. 2012 to Feb. 2013. After being euthanized, according to accepted guidelines (Sikes et al. 2011), specimens were processed following standard methods (Gardner 1996) and their ectoparasites identified using appropriate guides (Wharton and Fuller 1952, Keirans and Clifford 1978, Benton 1983, Kim et al. 1986, Keirans and Durden 1998). Voucher specimens of hosts were deposited in the mammal collection at Henderson State University (HSU). Ectoparasites were deposited in the General Ectoparasite Collection in the Department of Biology at Georgia Southern University (accession nos. L3549–3553, L3559–3560, L3570–3571, L3593–3594).

Both *S. carolinensis* and *S. niger* were infested with sucking lice, including 1 *S. carolinensis* with 2

female *Neohaematopinus sciuri* Jancke, 2 *S. niger* with *N. sciurinus* (1 squirrel with 1 male louse, the other with 2 female lice), and 1 *G. volans* with a female *N. sciuropteri*. In addition, a single *S. niger* was infested with a female squirrel tick, *Ixodes marxi* Banks and 1 larval American dog tick, *Dermacentor variabilis* (Say). Two *S. niger* were infested with lone star ticks, *Amblyomma americanum* (Linnaeus); 1 squirrel had 5 nymphs and the other, 26 nymphs; the former host was multiply parasitized by 2 male *D. variabilis* and 2 larval chiggers, *Eutrombicula alfreddugesi* (Oudemans). A single *G. volans* was infested with a larval *D. variabilis*. Two *S. carolinensis*, 1 *S. niger* and 2 of 10 (20%) *G. volans* harbored fleas, *Orchopeas howardi* (Baker); the single *T. striatus* was negative.

Siphonaptera: Ceratophyllidae

Orchopeas howardi was previously reported in AR from *S. carolinensis*, *S. niger*, and *G. volans* (Schiefer and Lancaster 1970), and commonly infests tree squirrels and some other rodents in many other states (Lewis 2000). It was therefore not surprising to find this flea on these hosts and associated nests.

Acari: Ixodidae

Ixodes marxi is not commonly reported and there are few records of this hard tick in AR (Lancaster 1973). This tick is a parasite of squirrels and is known to occur both east and west of the Mississippi River from at least 18 states (plus the District of Columbia) and Canada (Cooney and Hays 1972, Lancaster 1973, Keirans and Clifford 1978, Durden and Keirans 1996). It has been previously reported from the state on *S. carolinensis* (Tugwell and Lancaster 1962).

Immature stages (both larvae and nymphs) of *D. variabilis*, parasitize a wide range of small mammals and a few medium-sized mammals mainly in the eastern USA (Bequaert 1946, Tugwell and Lancaster 1972, Cooney and Hays 1972, Nicholson et al. 2009).

(there is also an isolated population of *D. variabilis* along the west coast of North America). However, Pung et al. (2000) recorded only 2 tick specimens (one *Amblyomma maculatum* and 1 *Ixodes scapularis*) from 70 *G. volans* in Georgia which suggests that ticks are not common on flying squirrels. They speculated that this is because *Glaucomys* spp. spend more time in arboreal habitats than on the ground where most ticks quest for hosts. *Dermacentor variabilis* is the principal vector of *Rickettsia rickettsii*, the causative agent of Rocky Mountain spotted fever, in the eastern USA (Chapman et al. 2006).

Amblyomma americanum is a widespread tick in AR (Lancaster 1973) and the southeastern USA where all active stages parasitize a variety of mammalian hosts including humans and immature stages also parasitize birds (Keirans and Durden 1998, Kinsey et al. 2000). This tick has become more abundant in some regions in recent years, and a number of zoonotic pathogens are known to be transmitted by it including *Ehrlichia chaffeensis* and *Ehrlichia ewingii*, causative agents of human ehrlichiosis, and *Borrelia lonestari* and *Rickettsia amblyommii*, putative agents of Southern Tick Associated Rash Illness (Childs and Paddock 2003, Billeter et al. 2007).

Acari: Trombiculidae

Only the larval stages of chiggers are ectoparasitic; nymphs and adults are predators of small arthropods or their eggs (Wharton and Fuller 1952). *Eutrombicula alfreddugesi* is a widespread pest chigger species in North America that parasitizes a wide variety of reptiles, birds and mammals, including humans (Wrenn and Loomis 1984). This species has previously been recorded from AR (Wicht and Rowland 1987).

Phthiraptera: Polyplacidae

Neohaematopinus sciuri is a Holarctic species that parasitizes *S. carolinensis* in North America (and introduced populations of this mammal in some other parts of the world) and the Eurasian red squirrel (*Sciurus vulgaris*) in Europe (Kim et al. 1986, Durden and Musser 1994). It has previously been reported from *S. carolinensis* in St. Francis Co., AR (Kim et al. 1986). However, *N. sciurinus*, to our knowledge, has not been previously reported from AR mammals and we document herein a new state record for this sucking louse. This louse parasitizes *S. niger* in North America and at least 6 other species of *Sciurus* in Central America (Durden and Musser 1994). *Neohaematopinus sciuropteri* is widespread across North America on both *G. volans* and the northern flying squirrel,

Glaucomys sabrinus (Durden and Musser 1994). However, there do not appear to be any previous records of this louse from Arkansas (Kim et al. 1986). *Neohaematopinus sciuropteri* is a vector of the zoonotic agent of sporadic epidemic typhus (caused by certain strains of *Rickettsia prowazekii*) to flying squirrels which act as reservoir hosts for this agent (Bozeman et al. 1981).

In conclusion, some sciurid ectoparasites that had not previously been reported from AR are probably widespread in the state. We suggest additional ectoparasite surveys on AR mammals, which appear to be uncommonly reported from hosts in the state. For example, prior to our study, only 1 species of parasitic mite has been reported from native mammals in AR (Richardson et al. 1994).

Acknowledgments

We thank the Arkansas Game and Fish Commission for a Scientific Collecting Permit issued to MBC, and Dr. R. Tumilson (HSU) for expert curatorial assistance.

Literature Cited

- Benton AH.** 1983. An illustrated key to the fleas of the eastern United States. Marginal Media (Fredonia, NY). 34 p.
- Bequaert JC.** 1946. The ticks, or Ixodoidea, of the northeastern United States and eastern Canada. Entomological America 25:73-232.
- Billeter SA, HL Blanton, SE Little, MG Levy and EB Breitschwerdt.** 2007. Detection of "*Rickettsia amblyommii*" in association with a tick bite rash. Vector-Borne and Zoonotic Diseases 7:607-610.
- Bozeman FM, DE Sonenshine, MS Williams, DP Chadwick, DM Lauer and BL Elisberg.** 1981. Experimental infection of ectoparasitic arthropods with *Rickettsia prowazekii* (GvF-16 strain) and transmission to flying squirrels. American Journal of Tropical Medicine and Hygiene 30:253-263.
- Chapman AS, SM Murphy, LJ Demma, RC Holman, AT Curns, JH McQuiston, JW Krebs and DL Swerdlow.** 2006. Rocky Mountain spotted fever in the United States, 1997-2002. Vector-Borne and Zoonotic Diseases 6:170-178.
- Childs JE and CD Paddock.** 2003. The ascendancy of *Amblyomma americanum* as a vector of pathogens affecting humans in the United States. Annual Review of Entomology 48:307-337.

Ectoparasites of Sciurid Rodents in Arkansas

- Cooney JC** and **KL Hays**. 1972. The ticks of Alabama. Auburn University Agricultural Experimental Station Bulletin 426:1-40.
- Durden LA** and **JE Keirans**. 1996. Nymphs of the genus *Ixodes* (Acari: Ixodidae) in the United States: taxonomy, identification key, distribution, hosts and medical/veterinary importance. Thomas Say Foundation Monographs, Entomological Society of America (Lanham, MD). 95 p.
- Durden LA** and **GG Musser**. 1994. The sucking lice (Insecta, Anoplura) of the world: a taxonomic checklist with records of mammalian hosts and geographical distributions. Bulletin of the American Museum of Natural History 218:1-90.
- Ellis HR**. 1975. Common ticks in Arkansas. Journal of the Arkansas Medical Society 81:380-382.
- Elrod DA**, **GA Heidt**, **DMA Elrod**, **M Birdsong** and **EG Zimmerman**. 1996. A second species of pocket gopher in Arkansas. Southwestern Naturalist 41:395-398.
- Gardner SL**. 1996. Field parasitology techniques for use with mammals. In: DE Wilson, editor. Measuring and monitoring biological diversity. Standard methods for mammals. Smithsonian Institution Press (Washington, DC). p. 291-298.
- Keirans JE** and **CM Clifford**. 1978. The genus *Ixodes* in the United States: a scanning electron microscope study and key to the adults. Journal of Medical Entomology, Supplement 2:1-149.
- Keirans JE** and **LA Durden**. 1998. An illustrated key to nymphs of the tick genus *Amblyomma* (Acari: Ixodidae) of the United States. Journal of Medical Entomology 35:489-495.
- Kim, KC**, **HD Pratt** and **CJ Stojanovich**. 1986. The sucking lice of North America: an illustrated manual for identification. The Pennsylvania State University Press (State University). 241 p.
- Kinsey AA**, **LA Durden** and **JH Oliver Jr**. 2000. Tick infestations of birds in coastal Georgia and Alabama. Journal of Parasitology 86:251-254.
- Lancaster JL Jr**. 1973. A guide to the ticks of Arkansas. University of Arkansas (Fayetteville) Agricultural Experiment Station Bulletin 779:1-39.
- Lewis RE**. 2000. A taxonomic review of the North American genus *Orchopeas* Jordan, 1933 (Siphonaptera: Ceratophyllidae: Ceratophyllinae). Journal of Vector Ecology 25:164-189.
- Nicholson WL**, **DE Sonenshine**, **RS Lane** and **G Uilenberg**. 2009. Ticks (Ixodoidea). In: GR Mullen & LA Durden, editors. Medical and Veterinary Entomology, 2nd edition. Elsevier (Amsterdam). p. 493-542.
- Price AD**, **VR McDaniel** and **R Tumilson**. 1986. An infestation of the bat bug *Cimex pilosellus* on an Arkansas population of big brown bats (*Eptesicus fuscus*). Proceedings of the Arkansas Academy of Science 36:98.
- Pung OJ**, **LA Durden**, **MJ Patrick**, **T Conyers** and **LR Mitchell**. 2000. Ectoparasites and gastrointestinal helminths of southern flying squirrels in southeast Georgia. Journal of Parasitology 86:1051-1055.
- Richardson DJ**, **LA Durden** and **DE Snyder**. 1994. Ectoparasites of the raccoon (*Procyon lotor*) from north-central Arkansas. Journal of the Kansas Entomological Society 67:208-212.
- Sealander JA** and **H Young**. 1954. Preliminary observations on the cave bats of Arkansas. Proceedings of the Arkansas Academy of Science 7:21-31.
- Schiefer BA** and **JL Lancaster Jr**. 1970. Some Siphonaptera from Arkansas. Journal of the Kansas Entomological Society 43:177-181.
- Sikes RS**, **WL Gannon**, and the **Animal Care and Use Committee of the American Society of Mammalogists**. 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. Journal of Mammalogy 92:235-253.
- Steward TW**, **VR McDaniel** and **DR England**. 1986. Additional records of distribution and hosts for the bat bug, *Cimex pilosellus* in Arkansas. Proceedings of the Arkansas Academy of Science 40:95-96.
- Tugwell P** and **JL Lancaster Jr**. 1962. Results of a tick-host study in northwest Arkansas. Journal of the Kansas Entomological Society 35:202-211.
- Wharton GW** and **HS Fuller**. 1952. A manual of the chiggers. Memoirs of the Entomological Society of Washington 4:1-185.
- Whitaker JO, Jr** and **N Wilson**. 1974. Host and distribution lists of mites (Acari), parasitic and phoretic, in the hair of wild mammals of North America, north of Mexico. American Midland Naturalist 91:1-67.
- Wicht MC** and **AC Rowland**. 1987. Fauna and distribution of living chiggers (Acarina: Trombiculidae) in Arkansas. Proceedings of the Arkansas Academy of Science 41:115-116.
- Wrenn WJ** and **RB Loomis**. 1984. Host selectivity in the genus *Eutrombicula* (Acari: Trombiculidae). In: DA Griffiths & CE Bowman, editors. Proceedings of the VI International Congress of Acarology, Vol. 1. Ellis Horwood Ltd., (Chichester, England). p. 160-165.