2013

New Host Records for the Fish Leech *Cystobranchnus klemmi* (Hirudinida: Piscicolidae) on Cyprinid Fishes from Arkansas and Oklahoma

Dennis J. Richardson
Quinnipiac University, dennis.richardson@quinnipiac.edu

R. Tumlison
Henderson State University

J. W. Allen Jr.
Quinnipiac University

W. E. Moser
National Museum of Natural History

C. T. McAllister
Follow this and additional works at: <https://scholarworks.uark.edu/jaas>
Eastern Oklahoma State College

Part of the [Zoology Commons](#)

See next page for additional authors

Recommended Citation

Richardson, Dennis J.; Tumlison, R.; Allen, J. W. Jr.; Moser, W. E.; McAllister, C. T.; Trauth, S. E.; and Robison, H. W. (2013) "New Host Records for the Fish Leech *Cystobranchnus klemmi* (Hirudinida: Piscicolidae) on Cyprinid Fishes from Arkansas and Oklahoma," *Journal of the Arkansas Academy of Science*: Vol. 67 , Article 39.

DOI: <https://doi.org/10.54119/jaas.2013.6734>

Available at: <https://scholarworks.uark.edu/jaas/vol67/iss1/39>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

New Host Records for the Fish Leech *Cystobranchus klemmi* (Hirudinida: Piscicolidae) on Cyprinid Fishes from Arkansas and Oklahoma

Authors

Dennis J. Richardson, R. Tumilson, J. W. Allen Jr., W. E. Moser, C. T. McAllister, S. E. Trauth, and H. W. Robison

New Host Records for the Fish Leech *Cystobranchus klemmi* (Hirudinida: Piscicolidae) on Cyprinid Fishes from Arkansas and Oklahoma

D.J. Richardson^{1*}, R. Tumilson², J.W. Allen, Jr¹, W.E. Moser³, C.T. McAllister⁴,
S.E. Trauth⁵, and H.W. Robison⁶

¹Department of Biological Sciences, Quinnipiac University, 275 Mt. Carmel Avenue, Hamden, Connecticut 06518

²Department of Biology, Henderson State University, Arkadelphia, Arkansas 71999

³Smithsonian Institution, National Museum of Natural History, Department of Invertebrate Zoology, Museum Support Center-MRC 534, 4210 Silver Hill Road, Suitland, Maryland 20746

⁴Science and Mathematics Division, Eastern Oklahoma State College, Idabel, Oklahoma 74745

⁵Department of Biological Sciences, Arkansas State University, P.O. Box 599, State University, Arkansas 72467

⁶Department of Biology, Southern Arkansas University, Magnolia, Arkansas 71754

*Correspondence: Dennis.Richardson@quinnipiac.edu

Running Title: New Host Records for the Fish Leech *Cystobranchus klemmi*

The Piscicolid fish leech *Cystobranchus klemmi* was originally described by Williams and Bureson (2005) as *Gonimosobdella klemmi* from the stoneroller, *Campostoma anomalum pullum*, and largescale stoneroller, *Campostoma oligolepis*, from the Little Red River in Searcy County, Arkansas. Williams and Bureson (2005) also reported *C. klemmi* from stonerollers from Bollinger, Cape Girardeau, Perry, Wayne and Washington counties in Missouri, and the redbfin shiner, *Lythrurus umbratilis*, and red shiner, *Cyprinella lutrensis*, from Union County, Illinois. Williams and Bureson (2006) synonymized the genus *Gonimosobdella* with *Cystobranchus*.

Examination of fish in the collection of Henderson State University, Arkadelphia, Arkansas revealed the presence of *C. klemmi* from central stonerollers, *C. anomalum*, from Franklin and Madison counties in Arkansas, and Cherokee County, Oklahoma, highland stonerollers, *Campostoma spadiceum*, from Garland, Hot Spring, Howard, Montgomery, Pike, and Sevier counties in Arkansas, a southern redbelly dace, *Chrosomus erythrogaster*, from Carroll County, Arkansas, and creek chubs, *Semotilus atromaculatus*, from Madison, Montgomery, and Independence counties in Arkansas (Figs. 1-2). Host vouchers are deposited in the vertebrate collection of the Biology Department at Henderson State University, Arkadelphia, Arkansas. All leeches were deposited in the Peabody Museum of Natural History, Yale University, in New Haven, Connecticut. Collection details along with accession numbers are given in Table 1.

This constitutes the first report of *C. klemmi* since its original description by Williams and Bureson

(2005) and the first report of this species from Oklahoma. Williams and Bureson (2005) described *C. klemmi*, in part, based on specimens collected from *C. a. pullum* from the Little Red River in Searcy County, Arkansas and from Apple Creek, Perry County, and Juden Creek and the St. Francis River in

Figure 1. *Cystobranchus klemmi* on a highland stoneroller, *Campostoma spadiceum*, from the Saline River in Howard County, Arkansas. Scale bar = 1cm.

Figure 2. *Cystobranchus klemmi* taken from a creek chub, *Semotilus atromaculatus* from a spring flowing out of Big Spring Mill in Independence County, Arkansas. Scale bar = 1mm.

Table 1. Summary of *Cystobranthus klemmi* collected in this study.

Host	Geographic Location	Number of Fish infested/ Number of fish in sample	Number of Leeches from each host	Collection Date	Host Accession #	Parasite Accession #
<i>Campostoma anomalum</i> (Central stoneroller)	Franklin Co. AR, Mulberry River	2/2	(5,1)	1 February 2003	HSU2855	YPM58384
	Madison Co. AR, White River	1/1	(8,3*)	24 March 1999	HSU2620	YPM58385
	Cherokee CO. OK, Rocky Ford State Park	1/11	(1,3*)	27 April 1991	HSU75	YPM58386
<i>Campostoma spadiceum</i> (Highland stoneroller)	Pike Co. AR, Caddo River	1/3	(1)	2 April 1994	HSU333	YPM58374
	Montgomery Co. AR, Lick Creek	1/6	(1)	2 April 1994	HSU376	YPM58375
	Pike Co. AR, Caddo River	1/3	(1)	20 February 1999	HSU2273	YPM58376
	Hot Spring Co. AR, DeRoche Creek	1/15	(1,2*)	14 January 1999	HSU2517	YPM58377
	Garland Co. AR	1/1	(4,1*)	9 March 1997	HSU1532	YPM58436
	Montgomery Co. AR, Long Creek	1/1	(1)	25 March 1997	HSU2080	YPM58437
	Hot Spring Co. AR, Blakely Creek	1/2	(2,1*)	9 March 1997	HSU1454	YPM58435
	Howard Co. AR, Saline river	1/2	(12)	21 February 1999	HSU2587	YPM58378
	Hot Spring Co. AR, Big Hill Creek	1/1	(1)	28 January 1999	HSU2442	YPM58379
	Sevier Co. AR, Woolsey Creek	1/2	(3)	16 February 1997	HSU1957	YPM58380
	Hot Spring Co. AR, Sanders Creek	1/3	(1,1*)	2 February 1999	HSU2556	YPM58381
	Hot Spring Co. AR, Bayou Dee	1/3	(1)	16 February 1997	HSU1468	YPM58382
	Hot Spring Co. AR, Marcus Creek	2/4	(2,4)	5 February 1994	HSU994	YPM58383
	Carroll Co. AR, Dry Creek	1/1	(1)	22 March 1999	HSU2609	YPM58394
	Madison Co. AR, War Eagle Creek	1/1	(2)	23 March 1999	HSU2614	YPM58387
- For HSU370: also see <i>Campostoma spadiceum</i> HSU376 from the same date and location	Montgomery Co. AR, Lick creek	3/5	(2,1,1,2*)	2 April 1994	HSU370	YPM58434
	Independence Co. AR, Big Spring Mill	1/1	(1)	23 February 2013	HSU3489	YPM58393

(All Leeches found on one specimen unless noted as (fish1, fish2); (*) indicates loose specimens found in jar unattached to fish. For more specific information regarding geographic collection localities and site of attachment on host refer to museum catalog numbers.)

Wayne County, Missouri. *Campostoma spadiceum* was resurrected and redescribed by Cashner et al. (2010) who indicated that in many instances specimens previously reported as *C. a. pullum* were actually *C. spadiceum*. Host vouchers were not mentioned by Williams and Bureson (2005) so it is not clear whether some of the host specimens that they examined were *C.*

a. pullum or *C. spadiceum*. In either case, this study definitively documents both *C. anomalum* and *C. spadiceum* as hosts for *C. klemmi*.

Cystobranthus klemmi was always attached to host fins. Chi square analysis revealed a significant preference in caudal attachment over pectoral ($\chi^2=10.00$; 1df) or pelvic ($\chi^2=8.80$; 1df). Pelvic or

New Host Records for the Fish Leech *Cystobranthus klemmi*

pectoral attachment sites were significantly preferred over dorsal or anal attachment sites ($\chi^2=9.84$, 1df). Attachment sites of 56 individuals are summarized in Fig. 3. The findings of *C. klemmi* on *C. erythrogaster* and *S. atromaculatus* represent new host records.

All occurrences of *C. klemmi* documented in this study were from cyprinid fishes occurring in streams and other lotic environments. Examination of a substantial number of fishes from lakes and reservoirs failed to reveal the presence of *C. klemmi*. Thus, *C. klemmi* may be characterized as a leech of cyprinid fishes of small streams and other lotic habitats. The known distribution of *C. klemmi* is given in Table 1 and Fig. 4.

Figure 3. Attachment sites of 56 specimens *Cystobranthus klemmi* on cyprinid fishes. Drawing after McAllister (1990).

Figure 4. Known distribution of *Cystobranthus klemmi* (+) indicated the type locality given by Williams and Bureson (2005). X indicates other localities reported by Williams and Bureson (2005). * indicates localities reported herein.

Literature Cited

- Cashner RC, WJ Matthews, E Marsh-Matthews, PJ Unmack and FM Cashner. 2010. Recognition and redescription of a distinctive stoneroller from the southern interior highlands. *Copeia* 2010 (no. 2):300-311.
- McAllister DE. 1990. A list of fishes of Canada. Syllogeus No. 64. National Museum of Natural Science, Ottawa, Canada, 310 p.
- Williams JI and EM Bureson. 2005. *Gonimosobdella klemmi* n. gen., n. sp. (Hirudinida: Piscicolidae) from cyprinid fishes in Arkansas, Illinois, and Missouri, U.S.A. *Comparative Parasitology* 72:166-172.
- Williams JI and EM Bureson. 2006. Phylogeny of the fish leeches (Oligochaeta, Hirudinida, Piscicolidae) based on nuclear and mitochondrial genes and morphology. *Zoologica Scripta* 35:627-639.