

2-28-2017

## Delivery of therapeutic agents by a collagen binding protein

Tulasi Ponnappakkam

Sagaya Theresa Leena Philominathan

Joshua Sakon  
*University of Arkansas, Fayetteville*

Ranjitha Katikaneni

Takaki Koide

*See next page for additional authors*

Follow this and additional works at: <https://scholarworks.uark.edu/pat>

---

### Citation

Ponnappakkam, T., Philominathan, S. T., Sakon, J., Katikaneni, R., Koide, T., Matsushita, O., Gensure, R. C., & Nishi, N. (2017). Delivery of therapeutic agents by a collagen binding protein. *Patents Granted*. Retrieved from <https://scholarworks.uark.edu/pat/318>

This Patent is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Patents Granted by an authorized administrator of ScholarWorks@UARK. For more information, please contact [scholar@uark.edu](mailto:scholar@uark.edu), [uarepos@uark.edu](mailto:uarepos@uark.edu).

---

## **Inventors**

Tulasi Ponnappakkam, Sagaya Theresa Leena Philominathan, Joshua Sakon, Ranjitha Katikaneni, Takaki Koide, Osamu Matsushita, Robert C. Gensure, and Nozomu Nishi



US009579273B2

(12) **United States Patent**  
**Ponnappakkam et al.**

(10) **Patent No.:** **US 9,579,273 B2**  
(45) **Date of Patent:** **Feb. 28, 2017**

(54) **DELIVERY OF THERAPEUTIC AGENTS BY A COLLAGEN BINDING PROTEIN**

(71) Applicants: **The Board of Trustees of the University of Arkansas**, Little Rock, AR (US); **The Kitasato Institute**, Tokyo (JP); **MONTEFIORE MEDICAL CENTER**, New York, NY (US); **NATIONAL UNIVERSITY CORPORATION KAGAWA UNIVERSITY**, Kagawa (JP)

(72) Inventors: **Tulasi Ponnappakkam**, New York, NY (US); **Sagaya Theresa Leena Philominathan**, Cheshire, CT (US); **Joshua Sakon**, Fayetteville, AR (US); **Ranjitha Katikaneni**, New York, NY (US); **Takaki Koide**, Tokyo (JP); **Osamu Matsushita**, Kanagawa (JP); **Robert C. Gensure**, New York, NY (US); **Nozomu Nishi**, Kagawa (JP)

(73) Assignees: **The Kitasato Institute** (JP); **National University Corporation Kagawa University** (JP); **Montefiore Medical Center**; **The Board of Trustees of the University of Arkansas**

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/365,226**

(22) PCT Filed: **Dec. 14, 2012**

(86) PCT No.: **PCT/US2012/069831**

§ 371 (c)(1),

(2) Date: **Jun. 13, 2014**

(87) PCT Pub. No.: **WO2013/090770**

PCT Pub. Date: **Jun. 20, 2013**

(65) **Prior Publication Data**

US 2014/0377215 A1 Dec. 25, 2014

**Related U.S. Application Data**

(60) Provisional application No. 61/570,620, filed on Dec. 14, 2011, provisional application No. 61/596,869, filed on Feb. 9, 2012.

(51) **Int. Cl.**

**A61K 8/65** (2006.01)

**A61K 8/99** (2006.01)

(Continued)

(52) **U.S. Cl.**

CPC ..... **A61K 8/66** (2013.01); **A61K 8/4913** (2013.01); **A61K 8/64** (2013.01); **A61K 8/65** (2013.01); **A61K 8/99** (2013.01); **A61K 38/179** (2013.01); **A61K 38/18** (2013.01); **A61K 38/1825** (2013.01); **A61K 38/1841** (2013.01); **A61K 38/1858** (2013.01); **A61K 38/1866**

(2013.01); **A61K 38/193** (2013.01); **A61K 38/29** (2013.01); **A61K 38/30** (2013.01); **A61K 39/44** (2013.01); **A61K 47/48246** (2013.01); **A61Q 7/00** (2013.01); **A61Q 7/02** (2013.01); **A61K 2039/505** (2013.01); **A61K 2039/6031** (2013.01); **A61K 2800/57** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56)

**References Cited**

**U.S. PATENT DOCUMENTS**

5,914,126 A 6/1999 Li et al.  
6,362,163 B1 3/2002 Gardella et al.  
(Continued)

**FOREIGN PATENT DOCUMENTS**

EP	0207751	1/1987
WO	WO 0006195	2/2000
WO	WO 0049159	8/2000
WO	WO 03052091	6/2003
WO	WO 2004071543	8/2004
WO	WO 2005082941	9/2005
WO	2006072623	7/2006
WO	WO 2010087397	8/2010
WO	WO 2010042425	11/2011
WO	WO 2012157339	11/2012

**OTHER PUBLICATIONS**

Philominathan et al, JBC, Apr. 17, 2009, 284/16:10868-10876.\*  
Locklin, R.M. et al., "Mediators of the biphasic responses of bone to intermittent and continuously administered parathyroid hormone," J Cell Biochem. (2003) 89(1):180-90.

(Continued)

*Primary Examiner* — Nita M Minnifield

(74) *Attorney, Agent, or Firm* — Andrus Intellectual Property Law, LLP

(57)

**ABSTRACT**

Methods of delivering therapeutic agents by administering compositions including a bacterial collagen-binding polypeptide segment linked to the therapeutic agent to subjects in need of treatment with the therapeutic agent are provided. In these methods, the therapeutic agent is not a PTH/PTHrP receptor agonist or antagonist, basic fibroblast growth factor (bFGF) or epidermal growth factor (EGF). The bacterial collagen-binding polypeptide segment delivers the agent to sites of partially untwisted or under-twisted collagen. Methods of treating collagenopathies using a composition including a collagen-binding polypeptide and a PTH/PTHrP receptor agonist are also provided. In addition, methods of treating hyperparathyroidism, and hair loss using compositions comprising a collagen binding polypeptide and a PTH/PTHrP receptor agonist are provided. Finally, methods of reducing hair regrowth by administering a composition including a collagen binding polypeptide and a PTH/PTHrP receptor antagonist are provided.

**12 Claims, 21 Drawing Sheets**  
**(15 of 21 Drawing Sheet(s) Filed in Color)**

(51) **Int. Cl.**

<b>A61K 38/29</b>	(2006.01)
<b>C07K 14/635</b>	(2006.01)
<b>A61Q 7/00</b>	(2006.01)
<b>A61K 8/66</b>	(2006.01)
<b>A61K 47/48</b>	(2006.01)
<b>A61K 8/49</b>	(2006.01)
<b>A61K 8/64</b>	(2006.01)
<b>A61K 38/17</b>	(2006.01)
<b>A61K 38/18</b>	(2006.01)
<b>A61K 38/19</b>	(2006.01)
<b>A61K 38/30</b>	(2006.01)
<b>A61K 39/44</b>	(2006.01)
<b>A61Q 7/02</b>	(2006.01)
<b>A61K 39/00</b>	(2006.01)

## (56)

**References Cited**

## U.S. PATENT DOCUMENTS

8,450,273 B2	5/2013	Sakon et al.	
9,062,300 B2	6/2015	Gensure et al.	
2002/0102709 A1	8/2002	Ishikawa et al.	
2002/0164719 A1	11/2002	Hall et al.	
2003/0187232 A1	10/2003	Hubbell et al.	
2004/0053368 A1	3/2004	Ishikawa et al.	
2004/0220094 A1	11/2004	Skinner et al.	
2005/0124537 A1	6/2005	Kostenuik et al.	
2005/0180986 A1	8/2005	Rich et al.	
2006/0014687 A1	1/2006	Crine et al.	
2006/0257376 A1	11/2006	Scadden et al.	
2008/0108562 A1	5/2008	Riviere et al.	
2009/0305352 A1 *	12/2009	Dai et al.	435/69.7
2010/0129341 A1 *	5/2010	Sakon	A61K 38/29 424/94.6
2013/0337017 A1 *	12/2013	Gensure	A61K 38/29 424/400
2014/0377215 A1 *	12/2014	Ponnappakkam	A61K 38/29 424/85.1
2015/0038423 A1 *	2/2015	Ponnappakkam	A61K 38/29 514/17.2
2015/0284701 A1 *	10/2015	Gensure	A61K 38/29 514/11.8

## OTHER PUBLICATIONS

Locus BAA06251 (GI 710023), Collagenase precursor from *Clostridium histolyticum*, Jan. 30, 2003. This amino acid sequence is disclosed in this application as SEQ ID No. 6. The sequence of residues 901-1021 of BAA06251 corresponds to the collagen binding domain included in the fusion protein of SEQ ID No. 1.

Locus EAW68494 (GI 119588900), Parathyroid hormone isoform from *Homo sapiens*, Dec. 18, 2006. Residues 64-147 of EAW68494 correspond to the PTH of SEQ ID No. 7.

Lotinun, S. et al., "Differential effects of intermittent and continuous administration of parathyroid hormone on bone histomorphometry and gene expression," *Endocrine*. (2002) 17(1):29-36.

Lotinun, S. et al., "Triazolopyrimidine (trapidil), a platelet-derived growth factor antagonist, inhibits parathyroid bone disease in an animal model for chronic hyperparathyroidism," *Endocrinology*. (2003) 144(5):2000-7.

Lumachi, F. et al., "Lumbar spine bone mineral density changes in patients with primary hyperparathyroidism according to age and gender," *Ann N Y Acad Sci*. (2007) 1117:362-6. Epub Jul. 26, 2007.

Ma, Y.L. et al., "Catabolic effects of continuous human PTH (1-38) in vivo is associated with sustained stimulation of RANKL and inhibition of osteoprotegerin and gene-associated bone formation," *Endocrinology* (2001) 142(9):4047-54.

Machado Do Reis, L. et al., "Accentuated osteoclastic response to parathyroid hormone undermines bone mass acquisition in osteonectin-null mice," *Bone* (2008) 43(2):264-73, Epub Apr. 13, 2008.

Malluche, H.H. et al., "Endogenous calcitonin does not protect against hyperparathyroid bone disease in renal failure," *Miner. Electrolyte Metab.* (1986) 12(2):113-8.

Malluche, H.H. et al., "Osteomalacia and hyperparathyroid bone disease in patients with nephrotic syndrome," *J Clin Invest.* (1979) 63(3):494-500.

Malluche, H.H. et al., "Influence of the parathyroid glands on bone metabolism," *Eur J Clin Invest.* (2006) 36(Suppl 2):23-33.

Malluche, H.H. et al., "Effects of long-term infusion of physiologic doses of 1-34 PTH on bone" *Am J Physiol.* (19S2) 242(2):F197-201.

Masi, L. et al., "Molecular, biochemical and cellular biology of PTH anabolic action," *J Endocrinol Invest.* (2005) 28(8 Suppl):37-40.

Mathias, R. et al., "Renal bone disease in pediatric and young adult patients on hemodialysis in a children's hospital," *J Am Soc Nephrol.* (1993) 3(12):1938-46.

Matsushita, O. et al., "A study of the collagen-binding domain of a 116-kDa *Clostridium histolyticum* collagenase," *J Biological Chem* (1998) 273(6):3643-3648.

Matsushita, O. et al., "Gene duplication and multiplicity of *C. Histolyticum* collagenases," *J Bacteriol.* (1999) 181:923-933.

Matsushita, O. et al., "Substrate recognition by the collagen-binding domain of *Clostridium histolyticum* class I collagenase," *J of Biological Chem* (2001) 276(12):8761-8770.

Matsushita, O., "Studies on the Clostridial Collagenases," *Nippon Saikingaku Zasshi* (1999) 54(4):753-761.

McCauley, L.K. et al., "PTH/PTHrP receptor is temporally regulated during osteoblast differentiation and is associated with collagen synthesis," *J Cell Biochem* (1996) 61:638-647.

McCauley, L.K. et al., "Proto-oncogene c-fos is transcriptionally regulated by parathyroid hormone (PTH) and PTH-related protein in a cyclic adenosine monophosphate-dependent manner in osteoblastic cells," *Endocrinology* (1997) 138(12):5427-33.

McCauley, L.K. et al., "Parathyroid hormone stimulates fra-2 expression in osteoblastic cells in vitro and in vivo," *Endocrinology* (2001) 142(5):1975-81.

Minisola, S. et al., "Trabecular bone mineral density in primary hyperparathyroidism: relationship to clinical presentation and biomarkers of skeletal turnover," *Bone Miner.* (1993) 20(2):113-23.

Minisola, S. et al., "Uneven deficits in vertebral bone density in postmenopausal patients with primary hyperparathyroidism as evaluated by posterior-anterior and lateral dual-energy absorptiometry," *Osteoporos Int.* (2002) 13(8):618-23.

Mitlak, B.H. et al., "Asymptomatic primary hyperparathyroidism," *J Bone Miner Res.* (1991) 6(Suppl 2):S103-10; discussion S121-4.

Miyachi, Y. et al., "Long-term safety and efficacy of high-concentration (20 mug/g) tacalcitol ointment in psoriasis vulgaris," *Eur J Dermatol* (2002) 12(5):463-468.

Morley, P. et al., "Anabolic effects of parathyroid hormone on bone," *Trends Endocrinol. Metab.* (1997) 8(6):225-31.

Morley, P. et al., "Parathyroid hormone: an anabolic treatment for osteoporosis," *Curr Pharm Des.* (2001) 7(8):671-87.

Murray, E.J. et al., "E64d, a membrane-permeable cysteine protease inhibitor, attenuates the effects of parathyroid hormone on osteoblasts in vitro," *Metabolism* (1997) 46(9):1090-4.

Nasu, M. et al., "Stimulatory effects of parathyroid hormone and 1,25-dihydroxyvitamin D3 on insulin-like growth factor-binding protein-5 mRNA expression in osteoblastic UMR-106 cells: the difference between transient and continuous treatments," *FEBS Lett.* (1997) 409(1):63-6.

Neer, R.M. et al., "Effect of parathyroid hormone (1-34) on fractures and bone mineral density in postmenopausal women with osteoporosis," *N. Engl. J. Med.* (2001) 344(19):1434-1441.

Nemeth, E.F., "Pharmacological regulation of parathyroid hormone secretion," *Corr Pharm. Des.* (2002) 8(23):2077-87.

Nilsson, P., "Bone disease in renal failure. Clinical and histomorphometric studies," *Scand J Urol Nephrol Suppl.* (1984) 84:1-68.

Nishi, N. et al., "Collagen-binding growth factors; Production and characterization of functional fusion proteins having a collagen-binding domain," *PNAS* (1998) 95(12):7018-7023.

(56)

## References Cited

## OTHER PUBLICATIONS

- O'Brien, C.A. et al., "IL-6 is not required for parathyroid hormone stimulation of RANKL expression, osteoclast formation, and bone loss in mice," *Am J Physiol Endocrinol Metab.* (2005) 289(5):E784-93. Epub Jun. 14, 2005.
- Okazaki, R., "Parathyroid hormone—its mechanisms of action and issues on clinical application," *Clin Calcium.* (2005) 15(5):845-51.
- Olgaard, K. et al., "Can hyperparathyroid bone disease be arrested or reversed?," *Clin J Am Soc Nephrol.* (2006) 1(3):367-73. Epub Mar. 29, 2006.
- Onyia, J.E. et al., "Molecular profile of catabolic versus anabolic treatment regimens of parathyroid hormone (PTH) in rat bone: an analysis by DNA microarray," *J Cell Biochem.* (2005) 95(2):403-18.
- Owens, R.J. et al., "Mapping the collagen-binding site of human fibronectin by expression in *Escherichia coli*," *The EMBO Journal* (1986) 5(11):2825-2830.
- Pailiard, M. et al., "Determinants of parathormone secretion in primary hyperparathyroidism," *Horm Res.* (1989) 32(1-3):89-92.
- Parfitt, A.M., "The actions of parathyroid hormone on bone: relation to bone remodeling and turnover, calcium homeostasis, and metabolic bone disease. Part IV of IV parts: The state of the bones in uremic hyperparathyroidism—the mechanisms of skeletal resistance to PTH in renal failure and pseudohypoparathyroidism and the role of PTH in osteoporosis, osteopetrosis, and osteofluorosis," *Metabolism.* (1976) 25(10):1157-88.
- Parfitt, A.M. et al., "Hypercalcemia due to constitutive activity of the parathyroid hormone (PTH)/PTHrP-related peptide receptor: comparison with primary hyperparathyroidism" *J Clin Endocrinol Metab.* (1996) 81(10):3584-8.
- Peters, E.M.J. et al., "A new strategy for modulating chemotherapy-induced alopecia, using PTH/PTHrP receptor agonist and antagonist," *J Invest Dermatol* (2001) 117(2):173-178.
- Pettway et al., "Anabolic actions of PTH (1-34): Use of a novel tissue engineering model to investigate temporal effects on bone," *Bone* (2005) 36(6):959-970.
- Phelps, E. et al., "Parathyroid hormone induces receptor activity modifying protein-3 (RAMP3) expression primarily via 3',5'-cyclic adenosine monophosphate signaling in osteoblasts," *Calcif Tissue Int.* (2005) 77(2):96-103 Epub Aug. 11, 2005.
- Philominathan et al., "Unidirectional binding of Clostridial Collagenase to Triple Helical Substrates," *Journal of Biological Chemistry* (2009) 284(16):10868-10876.
- Pirih, F.Q. et al., "Parathyroid hormone induces the NR4A family of nuclear orphan receptors in vivo," *Biochem Biophys Res Commun.* (2005) 332(2) 494-503.
- Podbesek, R. et al., "Effects of two treatment regimens with synthetic human parathyroid hormone fragment on bone formation and the tissue balance of trabecular bone in greyhounds," *Endocrinology* (1983) 112(3):1000-6.
- Poole, K.E. et al., "Parathyroid hormone—a bone anabolic and catabolic agent," *Curr Opin Pharmacol.* (2005) 5(6):612-7. Epub Sep. 21, 2005.
- Watson, P.H. et al., "Enhanced osteoblast development after continuous infusion of hPTH(1-84) in the rat," *Bone* (1999) 24(2):89-94.
- Weir, E.C. et al., "Synthetic parathyroid hormone-like protein (1-74) is anabolic for bone in vivo," *Calcif Tissue Int.* (1992) 51(1):30-4.
- Whitfield, J.F., "Taming Psoriatic Keratinocytes-PTHs' uses go up another notch," *J. Cell. Biochem.* (2004) 93:251-256.
- Wilson, J.J. et al., "A bacterial collagen-binding domain with novel calcium-binding motif controls domain orientation," *EMBO Journal* (2003) 22(8):1743-1752.
- Xu, M. et al., "Basal bone phenotype and increased anabolic responses to intermittent parathyroid hormone in healthy male COX-2 knockout mice," *Bone* (2010) 47(2):341-52. Epub May 13, 2010.
- Yang, C. et al., "Effects of continuous and pulsatile PTH treatments on rat bone marrow stromal cells," *Biochem Biophys Res Commun.* (2009) 380(4):791-6. Epub Feb. 3, 2009.
- Yoshihara, K. et al., "Cloning and nucleotide sequence analysis of the coIH gene from *Clostridium histolyticum* encoding a collagenase a gelatinase," *J Bacteriol* (1994) 176:6489-6496.
- Younes, N.A. et al., "Laboratory screening for hyperparathyroidism," *Clin Chim Acta.* (2005) 353(1-2):1-12.
- Zang, X.Y. et al., "Effects of parathyroid hormone and estradiol on proliferation and function of human osteoblasts from fetal long bone: An in vitro study," *Chin Med J (Engl).* (1994) 107(8):600-3.
- Zaruba, M.M. et al., "Parathyroid hormone treatment after myocardial infarction promotes cardiac repair by enhanced neovascularization and cell survival," *Cardiovasc Res* (2008) 77(4):722-731.
- Zhou, H. et al., "Anabolic action of parathyroid hormone on cortical and cancellous bone differs between axial and appendicular skeletal sites in mice," *Bone* (2003) 32(5):513-520.
- International Search Report and Written Opinion of the International Searching Authority for Application No. PCT/US08/004589 dated Oct. 28, 2008 (17 pages).
- Extended European Search Report for Application No. 08742686.2 dated Aug. 4, 2010 (8 pages).
- United States Patent Office Action for U.S. Appl. No. 12/594,547 dated Aug. 6, 2012 (12 pages).
- International Search Report and Written Opinion of the International Searching Authority for Application No. PCT/US12/69831 dated Mar. 14, 2013 (12 pages).
- International Search Report and Written Opinion of the International Searching Authority for Application No. PCT/US2013/25541 dated Jun. 17, 2013 (16 pages).
- Fukayama, S. et al., "New insights into interactions between the human PTH/PTHrP receptor and agonist/antagonist binding," *Am. J. Physiol. Endocrinol. Metab.* (1998) 274:297-301.
- Gao, Y. et al., "T cells potentiate PTH-induced cortical bone loss through CD40L signaling," *Cell Metab.* (2008) 8(2):132-45.
- Gardella, T.J., et al., "Converting Parathyroid Hormone-related Peptide (PTHrP) into a Potent PTH-2 Receptor Agonist," *J. of Biological Chemistry*, (1996) 271(31):19888-19893.
- Gensure, R.C. et al., "Parathyroid hormone and parathyroid hormone-related peptide, and their receptors," *Biochem Biophys Res Commun.* (2005) 328(1):666-78.
- Gensure, R.C. et al., "Parathyroid hormone without parathyroid glands," *Endocrinology* (2005) 146(2):544-546.
- Gevers, E.F. et al., "Bone marrow adipocytes: a neglected target tissue for growth hormone," *Endocrinology* (2002) 143(10):4065-73.
- Goltzman, D., "Studies on the mechanisms of the skeletal anabolic action of endogenous and exogenous parathyroid hormone," *Arch Biochem Biophys.* (2008) 473(2):218-24. Epub Mar. 10, 2008.
- Gopalakrishnan, R. et al., "Role of matrix Gla protein in parathyroid hormone inhibition of osteoblast mineralization," *Cells Tissues Organs* (2005) 181(3-4):166-75.
- Gosavi, A. et al., "An unusual presentation of parathyroid adenoma—a case report," *Indian J Pathol Microbiol.* (2005) 48(2):208-10.
- Gu, W.X. et al., "Mutual up-regulation of thyroid hormone and parathyroid hormone receptors in rat osteoblastic osteosarcoma 17/2.8 cells," *Endocrinology* (2001) 142(1):157-64.
- Hall, A.K. et al., "The effects of parathyroid hormone on osteoblast-like cells from embryonic chick calvaria," *Acta Endocrinol(Copenh).* (1985) 108(2):217-23.
- Han, B. et al., "Collagen-targeted BMP3 fusion proteins arrayed on collagen matrices or porous ceramics impregnated with Type I collagen enhance osteogenesis in a rat cranial defect model," *J Orthopaedic Research* (2002) 20:747-755.
- Headley, C.M., "Hungry bone syndrome following parathyroidectomy," *Anna J.*, (1998) 25(3):283-9; quiz 290-1.
- Heath, H., 3rd, "Clinical spectrum of primary hyperparathyroidism: evolution with changes in medical practice and technology," *J Bone Miner Res.* (1991) 6(Suppl 2):S63-70: discussion S83-4.
- Hoare, S.R. et al., "Specificity and stability of a new PTH1 receptor antagonist, mouse TIP(7-39)," *Peptides* (2002) 23(5):989-998.
- Hock, J.M. et al., "Human parathyroid hormone-(1-34) increases bone mass in ovariectomized and orchidectomized rats," *Endocrinology* (1988) 124(6):2899-2904.

(56)

## References Cited

## OTHER PUBLICATIONS

- Hock, J.M. et al., "Effects of continuous and intermittent administration and inhibition of resorption on the anabolic response of bone to parathyroid hormone," *J Bone Miner Res.* (1992) 7(1):65-72.
- Holick, M.F. et al., "Topical PTH (1-34) is a novel, safe and effective treatment for psoriasis: a randomized self-controlled trial and an open trial," (2003) *British J. Dermatology* 149:370-376.
- Homme, M. et al., "Differential regulation of RGS-2 by constant and oscillating PTH concentrations," *Calcif Tissue Int.* (2009). 84(4):305-12. Epub Feb. 20, 2009.
- Horwitz, M.J. et al., "Parathyroid hormone-related protein for the treatment of postmenopausal osteoporosis: defining the maximal tolerable dose," *J Clin Endocrinol Metab.* (2010) 95(3):1279-87.
- Horwitz, M.J. et al., "Continuous PTH and PTHrP infusion causes suppression of bone formation and discordant effects on 1,25(OH)<sub>2</sub> vitamin D," *J Bone Miner Res.* (2005) 20(10):1792-803. Epub Jun. 6, 2005.
- Hruska, K.A. et al., "Regulation of skeletal remodeling by parathyroid hormone," *Contrib Nephrol.* (1991) 91:38-42.
- Iida-Klein, A. et al., "Short-term continuous infusion of human parathyroid hormone 1-34 fragment is catabolic with decreased trabecular connectivity density accompanied by hypercalcemia in C57BL/6 mice," *J Endocrinol.* (2005) 186(3):549-57.
- Ishii, H. et al., "Daily intermittent decreases in serum levels of parathyroid hormone have an anabolic-like action on the bones of uremic rats with low-turnover bone and osteomalacia," *Bone* (2000) 26(2):175-82.
- Ishikawa, T. et al., "Delivery of a growth factor fusion protein having collagen-binding activity to wound tissues," *Artif. Organs* (2003) 27(2):147-154.
- Ishikawa, T. et al., "Production of a biologically active epidermal growth factor fusion protein with high collagen affinity," *J. Biochem.* (2001) 129(4):627-433.
- Ishizuya, T. et al., "Parathyroid hormone exerts disparate effects on osteoblast differentiation depending on exposure time in rat osteoblastic cells," *J Clin Invest.* (1997) 99(12):2961-70.
- Ito, M., "Parathyroid hormone and bone quality," *Clin Calcium.* (2005) 15(12):31-7.
- Ito, M., "Parathyroid and bone: Effect of parathyroid hormone on bone quality," *Clin Calcium.* (2007) 17(12):1858-64.
- Jeon, E. et al., "Engineering and application of collagen-binding fibroblast growth factor 2 for sustained release," (2013) *J. of Biomed. Materials Research: Part A*.
- Jilka, R.L., "Molecular and cellular mechanisms of the anabolic effect of intermittent PTH," *Bone* (2007) 40(6):1434-1446. Epub Apr. 6, 2007.
- Jilka, R.L. et al., "Continuous elevation of PTH increases the number of osteoblasts via both osteoclast-dependent and -independent mechanisms," *J Bone Miner Res.* (2010) 25(II):2427-37.
- Kaji, H., "Parathyroid and bone: Effects of parathyroid hormone on bone resorption and formation: differences between intermittent and continuous treatment," *Clin Calcium.* (2007) 17(12):1836-42.
- Katikaneni et al., "Treatment for chemotherapy-induced alopecia in mice using parathyroid hormone agonists and antagonists linked to collagen binding domain," *Int. J. Cancer* (2012) 131(5):E813-821.
- Kaye, M. et al., "Elective total parathyroidectomy without autotransplant in end-stage renal disease," *Kidney Int.* (1989) 35(6):1390-9.
- Khan, A. et al., "Primary hyperparathyroidism: pathophysiology and impact on bone," *Cmaj.* (2000) 163(2):184-7.
- Kido, S. et al., "Mechanism of PTH actions on bone," *Clin Calcium.* (2003) 13(1):14-8.
- Kistler, H., "Primary hyperparathyroidism: An analysis of 152 patients with special references to acute life threatening complications (acute hyperparathyroidism)," *Schweiz Med Wochenschr.* (1976) 106(Suppl 3): 1-61.
- Kitazawa, R. et al., "Effects of continuous infusion of parathyroid hormone and parathyroid hormone-related peptide on rat bone in vivo: comparative study by histomorphometry," *Bone Miner.* (1991) 12(3):157-66.
- Klempa, I., "Treatment of secondary and tertiary hyperparathyroidism—surgical viewpoints," *Chirurg.* (1999) 70(10):1089-101.
- Koh, A.J. et al., "3', 5'-Cyclic adenosine monophosphate activation in osteoblastic cells: effects on parathyroid hormone-1 receptors and osteoblastic differentiation in vitro," *Endocrinology* (1999) 140(7):3154-62.
- Komarova, S.V., "Mathematical model of paracrine interactions between osteoclasts and osteoblasts predicts anabolic action of parathyroid hormone on bone," *Endocrinology*, (2005) 146(8):3589-95. Epub Apr. 28, 2005.
- Kousteni, S. et al., "The cell biology of parathyroid hormone in osteoblasts," *Curr Osteoporos. Rep.* (2008)6(2):72-6.
- Kroll, M.H., "Parathyroid hormone temporal effects on bone formation and resorption," *Bull Math Biol.* (2000) 62(1):163-88.
- Lemaire, V. et al., "Modeling the interactions between osteoblast and osteoclast activities in bone remodeling," *J Theor Biol.* (2004) 229(3):293-309.
- Li, X. et al., "Determination of dual effects of parathyroid hormone on skeletal gene expression in vivo by microarray and network analysis," *J Biol. Chem.* (2007) 282(45):33086-97. Epub Aug. 9, 2007.
- Li, X. et al., "In vivo parathyroid hormone treatments and RNA isolation and analysis," *Methods Mol Biol.* (2008) 455:79-87.
- Liu, J. et al., "Intermittent PTH administration: a novel therapy method for periodontitis-associated alveolar bone loss." *Med Hypotheses.* (2009) 72(3):294-6. Epub Nov. 30, 2008.
- Potter, L.K. et al., "Response to continuous and pulsatile PTH dosing: a mathematical model for parathyroid hormone receptor kinetics," *Bone* (2005) 37(2):159-169.
- Potts, J.T., "Parathyroid hormone: past and present," *J Endocrinology* (2005) 187:311-325.
- Qin, L. et al., "Parathyroid hormone: a double-edged sword for bone metabolism," *Trends Endocrinol Metab.* (2004) 15(2):60-5.
- Rattanakul, C. et al., "Modelin of bone formation and resorption mediated by parathyroid hormone: response to estrogen/PTH therapy" *Biosystems* (2003) 70(1):55-72.
- Richardson, M.L. et al., "Bone mineral changes in primary hyperparathyroidism," *Skeletal Radiol.* (1986) 15(2):85-95.
- Rickard, D.J. et al., "Intermittent treatment with parathyroid hormone (PTH) as well as a non-peptide small molecule agonist of the PTH1 receptor inhibits adipocyte differentiation in human bone marrow stromal cells," *Bone* (2006) 39(6):1361-1372. Epub Aug. 10, 2006.
- Rixon, R.H. et al., "Parathyroid hormone fragments may stimulate bone growth in ovariectomized rats by activating adenyllyl cyclase." *J Bone Miner Res.* (1994) 9(8):1179-89.
- Robinson, J.A. et al., "Identification of a PTH regulated gene selectively induced in vivo during PTH-mediated bone formation," *J Cell Biochem.* (2006) 98(5):1203-20.
- Rosen, C.J., "The cellular and clinical parameters of anabolic therapy for osteoporosis," *Crit Rev Eukaryot Gene Expr.* (2003) 13(1):25-38.
- Rubin, M.R. et al., "The potential of parathyroid hormone as a therapy for osteoporosis," *Int J Fertil Womens Med.* (2002) 47(3):103-15.
- Rubin, M. et al., "The anabolic effects of parathyroid hormone therapy," *Osteoporosis International* (2002) 13(4):267-277.
- Rubin, M.R. et al., "The anabolic effects of parathyroid hormone therapy," *Clin Geriatr Med.* (2003) 19(2):415-32.
- Scafer, F., "Pulsatile parathyroid hormone secretion in health and disease," *Novartis Found Symp.* (2000) 227:225-39; discussion 239-43.
- Schluter, K.-D. et al., "A N-terminal PTHrP peptide fragment void of a PTH/PTHrP-receptor binding domain activates cardiac ETA receptors," *British Journal of Pharmacology* (2001) 132:427-432.
- Schmitt, C.P. et al., "Intermittent administration of parathyroid hormone (1-37) improves growth and bone mineral density in uremic rats," *Kidney Int.* (2000) 57(4):1484-92.
- Schmitt, C.P. et al., "Structural organization and biological relevance of oscillatory parathyroid hormone secretion," *Pediatr Nephrol.* (2005) 20(3):346-51. Epub Feb. 8, 2005.
- Schneider, A. et al., "Skeletal homeostasis in tissue-engineered bone," *J Orthop Res.* (2003) 21(5):859-64.

(56)

## References Cited

## OTHER PUBLICATIONS

- Seeman, E. et al., "Reconstructing the skeleton with intermittent parathyroid hormone," *Trends Endocrinol Metab.* (2001) 12(7):281-3.
- Shen, V. et al., "Skeletal effects of parathyroid hormone infusion in ovariectomized rats with or without estrogen repletion," *J Bone Miner Res.* (2000) 15(4):740-6.
- Shinoda, Y. et al., "Mechanisms underlying catabolic and anabolic functions of parathyroid hormone on bone by combination of culture systems of mouse cells," *J. of Cellular Biology* (2010) 109(4):755-63.
- Silver, J. et al., "Harnessing the parathyroids to create stronger bones," *Curr Opin Nephrol Hypertens.* (2004) 13(4):471-6.
- Silverberg, S.J. et al., "Skeletal disease in primary hyperparathyroidism," *J Bone Miner Res.*, (1989) 4(3):283-91.
- Skripitz, R. et al., "Parathyroid hormone—a drug for orthopedic surgery?," *Acta Orthop Scand.* (2004) 75(6):654-62.
- Skripitz, R. et al., "Stimulation of implant fixation by parathyroid hormone (1-34)—A histomorphometric comparison of PMMA cement and stainless steel," *J Orthop Res.* (2005) 23(6):1266-70. Epub Jun. 16, 2005.
- Smajilovic, S. et al., "Effect of intermittent versus continuous parathyroid hormone in the cardiovascular system of rats," *Open Cardiovasc. Med. J.* (2010) 4:110-6.
- Spurney, R.F. et al., "Anabolic effects of G protein-coupled receptor kinase inhibitor expressed in osteoblasts," *J Clin Invest.* (2002) 109(10):1361-71.
- Stewart, A.F., "PTHrP(1-36) as a skeletal anabolic agent for the treatment of osteoporosis," *Bone* (1996) 19(4):303-306.
- Stracke, S. et al., "Long-term outcome after total parathyroidectomy for the management of secondary hyperparathyroidism," *Nephron Clin Pract.* (2009) 111(2):c102-9. Epub Jan. 13, 2009.
- Strewler, G.J., "Local and systemic control of the osteoblast," *J. of Clin. Invest.* (2001) 107:271-272.
- Suttamanatwong, S. et al., "Regulation of matrix Gla protein by parathyroid hormone in MC3T3-E1 osteoblast-like cells involves protein kinase A and extracellular signal-regulated kinase pathways," *J Cell Biochem.* (2007) 102(2):496-505.
- Suttamanatwong, S. et al., "Sp proteins and Runx2 mediate regulation of matrix gla protein (MGP) expression by parathyroid hormone," *J Cell Biochem.* (2009) 107(2):284-92.
- Suzuki, A. et al., "PTH/cAMP/PKA signaling facilitates canonical Wnt signaling via inactivation of glycogen synthase kinase-3beta in osteoblastic Saos-2 cells," *J Cell Biochem.* (2008) 104(1):304-17.
- Swarthout, J.T. et al., "Parathyroid hormone-dependent signaling pathways regulating genes in bone cells," *Gene* (2002) 282(1-2):1-17.
- Swarthout, J.T. et al., "Stimulation of extracellular signal-regulated kinases and proliferation in rat osteoblastic cells by parathyroid hormone is protein kinase C-dependent" *J Biol Chem.* (2001) 276(10):7586-92, Epub Dec. 6, 2000.
- Takada, H. et al., "Response of parathyroid hormone to anaerobic exercise in adolescent female athletes," *Acta Paediatr Jpn.* (1998) 40(1):73-7.
- Takasu, H. et al., "Dual signaling and ligand selectivity of the human PTH/PTHrP receptor," *J Bone Miner Res.* (1999) 14(1):11-20.
- Talmage, R. V. et al., "Calcium homeostasis: reassessment of the actions of parathyroid hormone," *Gent Comp Endocrinol.* (2008) 156(1):1-8. Epub Nov. 12, 2007.
- Tam, C.S. et al., "Parathyroid hormone stimulates the bone apposition rate independently of its resorptive action: differential effects of intermittent and continuous administration," *Endocrinology* (1982) 10(2):506-12.
- Tawfeek, H. et al., "Disruption of PTH receptor 1 in T cells protects against PTH-induced bone loss," *PLoS* (2010) 5(8):e12290.
- Tokumoto, M. et al., "Parathyroid cell growth in patients with advanced secondary hyperparathyroidism: vitamin D receptor, calcium sensing receptor, and cell cycle regulating factors," *Ther Apher Dial.* (2005) 9(Suppl. 1.):S27-34.
- Tollin, S.R. et al., "Serial changes in bone mineral density and bone turnover after correction of secondary hyperparathyroidism in a patient with pseudohypoparathyroidism type Ib," *J Bone Miner Res.* (2000). 15(7):1412-6.
- Toyoshima, T. et al., "Collagen-binding domain of a Clostridium histolyticum collagenase exhibits a broad substrate spectrum both in vitro and in vivo," *Connective Tissue Research* (2001) 42(4):281-290.
- Uzawa, T. et al., "Comparison of the effects of intermittent and continuous administration of human parathyroid hormone(1-34) on rat bone," *Bone* (1995) 16(4):477-84.
- Vanstone, M.B. et al., "Rapid correction of bone mass after parathyroidectomy in an adolescent with primary hyperparathyroidism," *J. Clin. Endocrinol. Metab.* (2011) 96(2): E347-50. Epub Nov. 24, 2010.
- Wan, Q. et al., "Intra-articular injection of parathyroid hormone in the temporomandibular joint as a novel therapy for mandibular asymmetry," *Med Hypotheses* (2009) 74(4):685-7.
- Wang, C.A. et al., "Natural history of parathyroid carcinoma. Diagnosis, treatment, and results," *Am J Surg.* (1985) 149(4):522-7.
- Wang, Y. et al., "A theoretical model for simulating effect of parathyroid hormone on bone metabolism at cellular level," *Viol Cell Biomech.* (2009) 6(2):101-12.
- Wang, Y. et al., "Gender differences in the response of CD-1 mouse bone to parathyroid hormone: potential role of IGF-I," *J Endocrinol.* (2006) 189(2):279-87.
- Abdelhadi, M. et al., "Bone mineral recovery after parathyroidectomy in patients with primary and renal hyperparathyroidism," *J Clin Endocrinol Metab.* (1998) 83(11):3845-51.
- Abe, Y. et al., "Enhancement of graft bone healing by intermittent administration of human parathyroid hormone (1-34) in a rat spinal arthodesis model," *Bone* (2007) 41(5):775-785.
- Abshirini, H. et al., "Pathologic fractures: a neglected clinical feature of parathyroid adenoma," *Case* (2010) p. 357029. Epub Nov. 29, 2010.
- Akimoto, M. et al., "Effects of CB-VEGF-A injection in rat flap models for improved survival," (2013) *Plast. Reconstr. Surg.* 131(14):717-725.
- Aleksyniene, R. et al., "Parathyroid hormone—possible future drug for orthopedic surgery," *Medicina (Kaunas)* (2004) 40(9):842-9.
- Andrade, M.C., et al., "Bone mineral density and bone histomorphometry in children on long-term dialysis," *Pediatr Nephrol.* (2007) 22(10):1767-72. Epub Aug. 7, 2007.
- Barros, S.P., et al., "Parathyroid hormone protects against periodontitis-associated bone loss," *J Dent Res.* (2003) 82(10):791-5.
- Bauer, R., et al., "Structural comparison of ColH and ColG collagen-binding domains from Clostridium histolyticum," (2013), *Journal of Bacteriology*, 195(2), 318-327.
- Bedi, B., et al., "Inhibition of antigen presentation and T cell costimulation blocks PTH-induced bone loss," *Ann N Y Acad Sci.* (2010) 1192:215-21.
- Belinsky, G.S. et al., "Direct measurement of hormone-induced acidification in intact bone," *J Bone Miner Res.*, (2000) 15(3):550-6.
- Bellido, T., et al., "Chronic elevation of parathyroid hormone in mice reduces expression of sclerostin by osteocytes: a novel mechanism for hormonal control of osteoblastogenesis," *Endocrinology* (2005) 146(11):4577-83. Epub Aug. 4, 2005.
- Bergensstock, M.K. et al., "Parathyroid hormone stimulation of noncanonical Wnt signaling in bone" *Ann N Y Acad Sci.* (2007) 1116:354-9.
- Bergwitz, C. et al., "Rapid desensitization of parathyroid hormone dependent adenylate cyclase in perfused human osteosarcoma cells (SaOS-2)," *Biochem Biophys Acta.* (1994) 1222(3):447-56.
- Bianchi, E.N. et al., "Beta-arrestin2 regulates parathyroid hormone effects on a p38 MAPK and NFkappaB gene expression network in osteoblasts" *Bone* (2009) 45(4):716-25. Epub Jun. 25, 2009.
- Bilezikian, J.P., et al., "Asymptomatic primary hyperparathyroidism: new issues and new questions—bridging the past with the future," *J Bone Miner Res.* (2002) 17(Suppl 2):N57-67.

(56)

**References Cited****OTHER PUBLICATIONS**

- Bilezikian, J.P. et al., "Characterization and evaluation of asymptomatic primary hyperthyroidism," *J Bone Miner Res.* (1991) 6(Suppl 2):S85-9; discussion S121-4.
- Blachowicz, A. et al., "Serum I-84 and 7-84 parathyroid hormone concentrations and bone in patients with primary hyperparathyroidism," *Langenbecks Arch Surg.* (2008) 393(5):709-13, Epub Jul. 11, 2008.
- Buargub, M.A. et al., "Prevalence and pattern of renal osteodystrophy in chronic hemodialysis patients: a cross sectional study of 103 patients," *Saudi J Kidney Dis Transpl.* (2006) 17(3):401-7.
- Calvi, L.M. et al., "Activated parathyroid hormone/parathyroid hormone-related protein receptor in osteoblastic cells differentially affects cortical and trabecular bone," *J. Clin. Invest.* (2001) 107:277-286.
- Calvi, L.M. et al., "Osteoblastic cells regulate the haematopoietic stem cell niche," *Nature* (2003) 425:841-846.
- Canalis, E., "Effect of hormones and growth factors on alkaline phosphatase activity and collagen synthesis in cultured rat calvariac," *Metabolism* (1983) 32(1):14-20.
- Canalis, E. et al., "Insulin-like growth factor I mediates selective anabolic effects of parathyroid hormone in bone cultures," *J Clin Invest.* (1989) 83(1):60-5.
- Carter, P.H. et al., "Selective and Nonselective Inverse Agonists for Constitutively Active Type-1 Parathyroid Hormone Receptors: Evidence for Altered Receptor Conformations," *Endocrinology* (2001) 142(4):1534-1545.
- Chan, H.W. et al., "Prospective study on dialysis patients after total parathyroidectomy without autoimplant," *Nephrology* (2009) 15(4):441-7.
- Chen, B. et al., "Homogeneous osteogenesis and bone regeneration by demineralized bone matrix loading with collagen-targeting bone morphogenetic protein-2," *Biomaterials* (2007) 28:1027-1035.
- Chen, Q. et al., "Effects of an excess and a deficiency of endogenous parathyroid hormone on volumetric bone mineral density and bone geometry determined by peripheral quantitative computed tomography in female subjects," *J Clin Endocrinol Metab.* (2003) 88(10):4655-8.
- Cherian, P.P. et al., "Role of gap junction, hemichannels, and connexin 43 in mineralizing in response to intermittent and continuous application of parathyroid hormone," *Cell Commun Adhes.* (2008) 15(1):43-54.
- Chevalley, T. et al., "Bone and hormones, Effects of parathyroid hormone on the bone," *Presse Med.* (1999) 28(10):547-53.
- Cohen, A. et al., "Osteoporosis in adult survivors of adolescent cardiac transplantation may be related to hyperparathyroidism, mild renal insufficiency, and increased bone turnover," *J Heart Lung Transplant* (2005) 24(6):696-702.
- Compston, J.E., "Skeletal actions of intermittent parathyroid hormone: effects on bone remodelling and structure," *Bone* (2007) 40(6):1447-1452.
- Cormier, C., "Parathyroid hormone in osteoporosis," *Presse Med.* (2006) 35(3 Pt 2):495-501.
- Corsi, A. et al., "Osteomalacic and hyperparathyroid changes in fibrous dysplasia of bone: core biopsy studies and clinical correlations," *J Bone Miner Res.* (2003) 18(7): 1235-46.
- Cosman, F., "Parathyroid hormone treatment for osteoporosis," *Current Opinion in Endocrinology, Diabetes & Obesity* (2008) 15:495-501.
- Cundy, T. et al., "Hyperparathyroid bone disease in chronic renal failure," *Ulster Med J.* (1985) 54(Suppl):S34-43.
- Datta, N.S. et al., "Distinct roles for mitogen-activated protein kinase phosphatase-1 (MJP-1) and ERK-MAPK in PTH I R signaling during osteoblast proliferation and differentiation," *Cell* (2010) 22(3):457-66, Epub.
- Deal, C., "The use of intermittent human parathyroid hormone as a treatment for osteoporosis," *Curr Rheumatol Rep.* (2004) 6(1): 49-58.
- Demiralp, B. et al., "Anabolic actions of parathyroid hormone during bone growth are dependent on c-fos," *Endocrinology* (2002) 143(10):4038-47.
- Dobnig, H. et al., "The effects of programmed administration of human parathyroid hormone fragment (1-34) on bone histomorphometry and serum chemistry in rats," *Endocrinology* (1997) 138(11):4607-12.
- Drake, M.T. et al., "Parathyroid hormone increases the expression of receptors for epidermal growth factor in UMR 106-01 cells," *Endocrinology* (1994) 134(4):1733-7.
- Endo, K. et al., "1,25-dihydroxyvitamin D3 as well as its analogue OCT lower blood calcium through inhibition of bone resorption in hyperealemic rats with continuous parathyroid hormone-related peptide infusion," *J Bone Miner Res.* (2000) 15(1):175-81.
- Etoh, M. et al., "Repetition of continuous PTH treatments followed by periodic withdrawals exerts anabolic effects on rat bone," *J Bone Miner Metab.* (2010) 28(6):641-649.
- Fitzpatrick, L.A. et al., "Acute primary hyperparathyroidism," *Am J. Med.* (1987) 82(2) 75-82.
- Fleming, A. et al., "High-throughput in vivo screening for bone anabolic compounds with zebrafish," *J Biomol Screen.* (2005) 10(8):823-31. Epub Oct. 18, 2005.
- Fouda, M.A., "Primary hyperparathyroidism: King Khalid University Hospital Experience," *Ann Saudi Med.* (1999) 19(2):110-5.
- Fox, J. et al., "Effects of daily treatment with parathyroid hormone 1-84 for 16 months on density, architecture and biomechanical properties of cortical bone in adult ovariectomized rhesus monkeys," *Bone* (2007) 41(3):321-330.
- Fraher, L.J. et al., "Comparison of the biochemical responses to human parathyroid hormone-(1-31)NH2 and hPTH-(1-34) in healthy humans," *J Clin Endocrinol Metab.* (1999) 84(8):2739-43.
- Frolik, C.A. et al., "Anabolic and catabolic bone effects of human parathyroid hormone (1-34) are predicted by duration of hormone exposure," *Bone* (2003) 33(3):372-379.
- Fujita, T., "Parathyroid hormone in the treatment of osteoporosis," *BioDrugs* (2001) 15(11):721-728.

\* cited by examiner



Fig. 1




Fig. 1 (Continued)







Fig. 2

PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>


Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-  
Hyp-Gly-Pro-Hyp-Ala-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-NH<sub>2</sub>

PROXYL-(POG)<sub>5</sub>POA(POG)<sub>4</sub>


Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-  
Hyp-Ala-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-NH<sub>2</sub>

PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>


Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Ala-Pro-  
Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-NH<sub>2</sub>

PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>

Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Ala-Pro-Hyp-Gly-Pro-  
Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-NH<sub>2</sub>

(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-PROXYL

H-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Ala-  
Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Cys-NH<sub>2</sub>

<sup>11</sup>PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>

H-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Cys-Gly-Pro-  
Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-Pro-Hyp-Gly-NH<sub>2</sub>

Fig. 3A




Fig. 3B




Fig. 4




Fig. 5

Fig. 5A




Fig. 5B




Fig. 5C




Fig. 6

Fig. 6A




Fig. 6B




Fig. 6C




Fig. 6 (cont.)

Fig. 6D




Fig. 6E




Fig. 6F






Fig. 6 (cont.)

Fig. 6G




Fig. 6H




Fig. 6I




Fig. 7

Fig. 7A




Fig. 7B





Fig. 7C




Fig. 7D

Fig. 8

Fig. 8A, 8B




Fig. 8C, 8D




Fig. 8E, 8F




Fig. 8G, 8H




Fig. 9




Fig. 10

Fig. 10A




Fig. 10B





Fig. 10C




Fig. 10D

Fig. 11




Fig. 12






Fig. 13


A) Normal Control



B) CYP



C) CYP + Agonist



D) CYP + Antagonist




Fig. 14






Fig. 15



**Grey Scale Analysis at Injection Site**

Fig. 16




Fig. 17



**Chemotherapy****Prophylaxis****Therapy**

Fig. 18




Fig. 19





Fig. 20



1

## DELIVERY OF THERAPEUTIC AGENTS BY A COLLAGEN BINDING PROTEIN

### CROSS-REFERENCE TO RELATED APPLICATIONS

This patent application is a national stage filing under 35 U.S.C. 371 of International Application No. PCT/US2012/069831, filed Dec. 14, 2012, which claims the benefit of priority of United States Provisional Patent Application No. 61/570,620, filed Dec. 14, 2011 and of United States Provisional Patent Application No. 61/596,869, filed Feb. 9, 2012, all of which are incorporated herein by reference in their entireties.

### STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

This invention was made with United States government support awarded by the National Institutes of Health grant number NCRR COBRE 8P30GM103450 and INBRE GM103429. The United States may have certain rights in this invention.

### SEQUENCE LISTING

A Sequence Listing accompanies this application and is incorporated herein by reference in its entirety. The Sequence Listing was filed with the application as a text file on Dec. 14, 2012.

### INTRODUCTION

Delivery of therapeutic agents to sites within the body of a subject where a particular therapeutic agent is needed in order to be effective is a developing area. Such delivery systems will allow more efficient use of therapeutic agents while reducing toxicity caused by some therapeutic agents. Use of targeted liposomes or polypeptides, such as antibodies, to target therapeutic agents to particular sites within the body has proved successful, but additional delivery agents are needed.

Alopecia (hair loss) is a psychologically and emotionally distressing event with multiple causes. Alopecia occurs most commonly in male-pattern baldness, affecting approximately two thirds of males by age 35; a similar pattern of hair loss can be observed in females with polycystic ovarian syndrome. In both of these disorders, the hair loss is androgen mediated. Alopecia can also occur as an autoimmune disease, termed alopecia areata; a disorder which affects 1.7% of the population. It can occur as a side-effect of medical treatments, particularly in chemotherapy, with 65-85% of chemotherapy patients experiencing some degree of alopecia. Psychological consequences of hair loss have been well studied in the chemotherapy setting. Chemotherapy-induced alopecia (CIA) can result in anxiety, depression, a negative body image, lowered self-esteem and a reduced sense of well-being. In fact, 47-58% of female cancer patients consider hair loss to be the most traumatic aspect of chemotherapy, and 8% would decline treatment for fear of hair loss. In addition to these studies in chemotherapy patients, evidence exists in other forms of alopecia to support therapy to reduce psychological consequences of hair loss. Thus a new treatment to stop hair loss or speed hair regrowth would be beneficial.

While drugs with mild anti-androgenic effects (i.e. spironolactone) had been used with limited success as

2

therapy for alopecia, the first effective medication for alopecia was minoxidil (Rogaine). This antihypertensive has an observed side-effect of causing hair growth, and is now used as topical therapy for many forms of alopecia. However, responses are incomplete, with some subjects showing only slowing of hair loss rather than actual regrowth. Finasteride (Propecia) is a newer agent that blocks conversion of testosterone to dihydrotestosterone, resulting in improvements in androgenic alopecia at the expense of partial systemic androgen blockade. However, response rates with long-term (10 years) therapy are only around 50%. Overall, despite considerable research in this area, there is still no adequate therapy for hair loss.

In addition, unwanted hair growth is a cosmetic issue many people deal with on a regular basis. Unwanted hair growth on the face, legs, arms, chest or back is a growing cosmetic problem. Many people use laser therapy, waxing or other therapies to remove unwanted hair. There are currently no topical pharmaceuticals to limit hair growth.

Collagenopathies represent a large number of diseases in which collagen structure or formation is not normal. This group of diseases results in a broad spectrum of symptoms including bone defects, vascular defects, and skin defects. Many of these diseases have no or only ineffective treatments available.

For example, osteogenesis imperfecta (OI), also known as brittle bone disease, is caused by an inborn mutation of type I collagen. Approximately 25,000 to 50,000 Americans are affected and the effects of the disease range from mild, in which many individuals are unaware of the disease, to severe in which individuals cannot live a normal life due to recurrent broken bones. Most OI patients carry a mutation which causes an amino acid change in collagen changing a glycine to a bulkier amino acid which results in disruption of the triple helix structure of the collagen and under-twisting. The body may respond by hydrolyzing the collagen and this may result in a reduction in bone strength. There is currently no cure and few treatments for OI.

### SUMMARY

Provided herein are methods of delivering therapeutic agents by administering compositions including a bacterial collagen-binding polypeptide segment linked to the therapeutic agent to subjects in need of treatment with the therapeutic agent. In these methods, the therapeutic agent is not a PTH/PTHrP receptor agonist or antagonist, basic fibroblast growth factor (bFGF) or epidermal growth factor (EGF) and the bacterial collagen-binding polypeptide segment delivers the agent to sites of partially untwisted or under-twisted collagen.

In another aspect, methods of treating a subject with a collagenopathy, such as osteogenesis imperfecta, by administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor agonist to a subject in an amount effective to treat the collagenopathy are provided. The bacterial collagen-binding polypeptide segment delivers the agent to sites of partially untwisted or under-twisted collagen.

In yet another aspect, methods of treating hyperparathyroidism by administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor agonist to a subject are provided.

In still a further aspect, methods of slowing hair growth or regrowth after removal by administering a composition

comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor antagonist to a subject are provided.

In a still further aspect, methods of increasing hair growth or the speed of hair re-growth after removal or loss by administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor agonist to a subject are provided.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The patent application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the office upon request and payment of the necessary fee.

FIG. 1 is a sequence alignment showing the alignment of several M9B bacterial collagenases from the *Bacillus* and *Clostridium* families. The residues shown in blue are important for collagen binding activity, those shown in green are important for maintaining the architecture or protein folding. Both of these are also underlined for the top and bottom sequences. Residues shown in red are critical for  $\text{Ca}^{2+}$  binding and those in orange are critical for positioning the  $\text{Ca}^{2+}$  binding residues.

FIG. 2 is a set of drawings showing the chemical structures of synthesized peptides.

FIG. 3A is a graph showing the circular dichroism spectra of the collagenous peptides measured at 4° C.

FIG. 3B is a graph showing the thermal denaturation profile of the various collagenous peptides. The temperature was increased at the rate of 0.3° C./min.

FIG. 4A is a graph showing the scattering profile with the intensity  $I(Q)$  plotted against the scattering vector  $Q$ .

FIG. 4B is a graph showing the pair-distance distribution function  $P(r)$  in the real space obtained using GNOM for [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub>:CBD complex (Red), [PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub>:CBD complex (Blue), [PROXYL-(POG)<sub>5</sub>POA(POG)<sub>4</sub>]<sub>3</sub>:CBD complex (Green), [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub>:CBD complex (Orange) and [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub>:CBD complex (Cyan).

FIG. 5 is a set of plots showing HSQC NMR data obtained using the collagen binding domain (CBD)—collagenous peptide interactions. FIG. 5A shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonance of V973, G975 and S979 are present during this titration. FIG. 5B shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub>:CBD complex (red) at 1:1 ratio. Amide resonances of V973, G975 and S979 disappeared because of their proximity to the spin-labeled group. FIG. 5C is a cartoon showing the structure of CBD and the CBD residues that are line broadened upon titration with [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub>.

FIG. 6 is a set of plots showing HSQC NMR data obtained using the CBD—collagenous peptide interactions. FIG. 6A shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonances of Q972, V973, G975 and S979 are present during this titration. FIG. 6B shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>4</sub>]<sub>3</sub>:CBD complex (red) at ratio 1:1. Amide resonances of Q972, V973, G975 and S979 are line broadened due to the PROXYL moiety. FIG. 6C is a cartoon of the structure of CBD showing the CBD residues that are

uniquely line broadened upon titration with [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>4</sub>]<sub>3</sub>. FIG. 6D shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonances of L946, Q972, V973, G975 and S979 are present during this titration. FIG. 6E shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub>:CBD complex (red) at 1:1 ratio. Amide resonances of L946, Q972, V973, G975 and S979 disappeared because of the spin-label. FIG. 6F shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub>:CBD (cyan) at ratio 1:1. In the absence of spin label, amide resonances of L946, Q972, V973, G975 and S979 are not line broadened. FIG. 6G shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonances of L946, G953, Q972, V973, D974, G975, N976, V978, S979 are present during this titration. FIG. 6H shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub>:CBD complex (red) at ratio 1:1. Amide resonances of L946, G953, Q972, V973, D974, G975, N976, V978, S979 are line broadened due to the PROXYL moiety. FIG. 6I is a cartoon of the structure of CBD showing the CBD residues that are line broadened by the spin label of [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub>.

FIG. 7 is a set of graphs showing the intensity drop of (A) Q972, (B) G975, (C) S979 and (D) L924 on CBD as a function of increasing concentrations of mini-collagen i.e. [(POG)<sub>10</sub>]<sub>3</sub> (black), [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub> (red), [PROXYL-(POG)<sub>5</sub>POA(POG)<sub>4</sub>]<sub>3</sub> (blue), [PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub> (green), and [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub> (cyan).

FIG. 8 is a set of plots showing HSQC NMR data obtained using the CBD—collagenous peptide interactions. FIG. 8A shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonances of S906, S997 and G998 are present during this titration. FIG. 8B shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-PROXYL]<sub>3</sub>:CBD complex (red) at ratio 1:1. Amide resonances of S906, S997 and G998 are line broadened due to the PROXYL moiety. FIG. 8C shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-carbamidomethyl]<sub>3</sub>:CBD (cyan) at 1:1 ratio. In the absence of spin label, amide resonances of S906, S997 and G998 are not line broadened. FIG. 8D is a cartoon of the structure of CBD showing the CBD residues that are line broadened due to the spin label of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-PROXYL]<sub>3</sub>. Amide resonances of S906, S997 and G998 (red) disappeared upon titration with [(POG)<sub>4</sub>POA(POG)<sub>5</sub>-PROXYL]<sub>3</sub>. FIG. 8E shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>10</sub>]<sub>3</sub>:CBD complex (green) at 1:1 ratio. Amide resonances of S906, Q972, V973, G975, S979, S997 and G998 are present during this titration. FIG. 8F shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub>:CBD complex (red) at 1:1 ratio. Amide resonances of S906, Q972, V973, G975, S979, S997 and G998 disappeared because of the spin-label. FIG. 8G shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (black) and <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub>:CBD (cyan) at ratio of 1:1. Resonances of S906, Q972, V973, G975, S979, S997 and G998 are intact in the

5

absence of the spin label. FIG. 8H is a cartoon of the structure of CBD showing the residues that are line broadened upon titration with [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub>. Only amide resonances of S906, R929, S997, and G998 (red) disappeared at 0.2:1 ratio. When the peptide ratio was raised to 0.3:1, additional resonances of V973, G975, S979 (blue) disappeared.

FIG. 9 is a set of structure drawings derived from SAXS scattering profiles using ab initio simulated annealing calculations for (A) [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub>:CBD complex, (B) [PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub>:CBD complex, (C) [PROXYL-(POG)<sub>5</sub>POA(POG)<sub>4</sub>]<sub>3</sub>:CBD complex and (D) [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub>:CBD complex, (E) [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-PROXYL]<sub>3</sub>:CBD complex, (F) [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-carbamidomethyl]<sub>3</sub>:CBD. The Gly→Ala mutation sites are highlighted. FIGS. 9G and 9H show two probable binding modes of [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub>:CBD complex.

FIG. 10 is a set of plots showing HSQC NMR data obtained using the CBD—collagenous peptide interactions. FIG. 10A is an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [POGPO-<sup>15</sup>N-G-(POG)<sub>8</sub>]<sub>3</sub> (black) with <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [POGPO-<sup>15</sup>N-G-(POG)<sub>8</sub>]<sub>3</sub>:CBD complex (red) at 1:1 ratio. FIG. 10B shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [POGPO-<sup>15</sup>N-G-(POG)<sub>2</sub>POA-(POG)<sub>5</sub>]<sub>3</sub> (black) with <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [POGPO-<sup>15</sup>N-G-(POG)<sub>2</sub>POA-(POG)<sub>5</sub>]<sub>3</sub>:CBD complex (red) at 1:1 ratio. FIG. 10C shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>8</sub>-PO-<sup>15</sup>N-G-POG]<sub>3</sub> (black) with <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>8</sub>-PO-<sup>15</sup>N-G-POG]<sub>3</sub>:CBD complex (red) at 1:1 ratio. FIG. 10D shows an overlay of <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>4</sub>-POA-PO-<sup>15</sup>N-G-POG]<sub>3</sub> (black) with <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of [(POG)<sub>4</sub>-POA-PO-<sup>15</sup>N-G-POG]<sub>3</sub>:CBD complex (red) at 1:1 ratio.

FIG. 11 shows the tissue distribution of S<sup>35</sup>-PTH-CBD 1 hour and 12 hours after subcutaneous injection. Note the skin outline.

FIG. 12 is a set of photographs documenting the hair growth on the back of mice at day 36 after depilation, treatment groups as indicated (Antagonist=PTH(7-33)-CBD, Agonist=PTH-CBD).

FIG. 13 is a set of photographs showing the histology at Day 36 after the indicated treatment. Skin samples were taken from the dorsal region and processed for Hematoxylin and Eosin (H&E) staining. Representative sections are shown from each treatment group as indicated. (Antagonist=PTH(7-33)-CBD, Agonist=PTH-CBD).

FIG. 14 is a graph showing the hair follicle counts per high powered field. Anagen VI hair follicles were counted by two independent observers in a blinded fashion. Results are expressed as mean±standard deviation. \*\*=p<0.01 vs. no chemo ANOVA followed by Dunnett's test. (Antagonist=PTH(7-33)-CBD, Agonist=PTH-CBD).

FIG. 15 is a set of photographs showing the hair growth on the back of the mice after each of the indicated treatments and a graph showing the results of a grey scale analysis of the hair at the injection site over time after the injection.

FIG. 16 is a set of photographs showing the hair on the back of mice after the indicated treatment without prior depilation.

FIG. 17 is a set of photographs and a graph showing the grey scale analysis of hair growth on the backs of mice comparing the indicated treatments with the PTH-CBD being administered prior to the chemotherapy as opposed to after chemotherapy began.

6

FIG. 18 is a photograph of three mice 13 days after waxing to remove hair and treatment with PTH-CBD, PTH antagonist-CBD or vehicle alone.

FIG. 19 is a set of photographs of mice showing hair regrowth in a model of alopecia areata after treatment with a control or with PTH-CBD.

FIG. 20 is a graph showing the endogenous parathyroid hormone levels in ovariectomized aged rats injected with a single dose of human PTH-CBD 6 months prior to sacrifice.

## DETAILED DESCRIPTION

Methods of delivering a therapeutic agent by administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a therapeutic agent to a subject in need of treatment with the therapeutic agent are provided herein. In this embodiment, the therapeutic agent is not a PTH/PTHrP receptor agonist or antagonist and is not a bFGF or EGF polypeptide. The bacterial collagen-binding polypeptide segment delivers the therapeutic agent to sites of partially untwisted or under-twisted collagen.

In addition, methods of treating collagenopathies, such as osteogenesis imperfecta (OI), by administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor agonist to a subject in need of treatment for a collagenopathy are provided. Collagenopathies include but are not limited to osteogenesis imperfecta, Stickler's syndrome, Ehlers-Danlos syndrome, Alport's syndrome, Caffey's disease, and localized collagen or cartilage damage. Many of these diseases are caused by genetic defects that result in the collagen in certain tissues being under twisted or partially untwisted.

For example, individuals with OI carry a mutation which causes an amino acid change in collagen changing a glycine to a bulkier amino acid which results in disruption of the triple helix structure of the collagen and under-twisting of the collagen. In the Examples, we demonstrate that the bacterial collagen-binding polypeptides described herein target and bind to these areas of under-twisted collagen. Thus, use of the collagen-binding polypeptides described herein to deliver a therapeutic agent capable of treating OI to the sites of under-twisted collagen may allow more effective treatment.

The collagen-binding polypeptide segment and the therapeutic agent may be chemically cross-linked to each other or may be polypeptide portions of a fusion protein. The terms "fusion protein" and "fusion polypeptide" may be used to refer to a single polypeptide comprising two functional segments, e.g., a collagen-binding polypeptide segment and a polypeptide based therapeutic agent, such as PTH/PTHrP receptor agonist polypeptide segment. The fusion proteins may be any size, and the single polypeptide of the fusion protein may exist in a multimeric form in its functional state, e.g., by cysteine disulfide connection of two monomers of the single polypeptide. A polypeptide segment may be a synthetic polypeptide or a naturally occurring polypeptide. Such polypeptides may be a portion of a polypeptide or may comprise one or more mutations. The two polypeptide segments of the fusion proteins can be linked directly or indirectly. For instance, the two segments may be linked directly through, e.g., a peptide bond or chemical cross-linking, or indirectly, through, e.g., a linker segment or linker polypeptide. The peptide linker may be any length and may include traditional or non-traditional amino acids. For example, the peptide linker may be 1-100 amino acids long, suitably it is 5, 10, 15, 20, 25 or more amino acids long such that the collagen binding portion of the fusion polypeptide



can mediate collagen binding and the therapeutic agent can have its therapeutic effect. Peptide linkers may include but are not limited to a PKD (polycystic kidney disease) domain from a collagenase or other protein such as in SEQ ID NO: 2, a GST or His-tag, or a Ser or Gly linker.

The collagen-binding polypeptide segment is a polypeptide that binds collagen and may be part of a larger fusion protein, bioactive agent, or pharmaceutical agent. Determination of whether a composition, polypeptide segment, fusion protein, or pharmaceutical or bioactive agent binds collagen can be made as described in U.S. Patent Publication No. 2010/0129341, which is incorporated herein by reference in its entirety. Briefly, it is incubated with collagen in binding buffer, and the mixture is then filtered through a filter that would otherwise allow it to pass through but that blocks the collagen and therefore holds back materials that bind to the collagen. The filtrate is then assayed for the presence of the composition, polypeptide segment, fusion protein, or pharmaceutical or bioactive agent. Suitably, at least 80%, 85%, 90%, 95%, 98% or more suitably at least 99% of the collagen-binding composition, polypeptide segment, fusion protein, or pharmaceutical or bioactive agent is retained by the filter in this assay, as compared to when the filtration is performed without collagen.

The collagen-binding polypeptide segment may be a bacterial collagen-binding polypeptide segment. It may be a *Clostridium* collagen-binding polypeptide segment. The collagen-binding polypeptide segment may be a segment of a collagenase, or a bacterial collagenase, or a *Clostridium* collagenase. Suitably the polypeptide segment is only a portion of the collagenase and the collagen-binding polypeptide segment does not have collagenase activity. The collagen-binding polypeptide may be a bacterial M9B (including those derived from *Bacillus* spp. and *Clostridium* spp.) or M9A (including those derived from *Vibrio* spp.) collagen-binding protein or a collagen-binding peptide derived from such a protein. By "derived from" we mean that the peptide is a fragment of the full-length protein, a peptide that has amino acid changes relative to the wild-type protein or a combination thereof. The key is that the peptide retains the ability to bind collagen. For example, a peptide may be derived from a protein by selecting a region of the protein capable of binding to collagen. Compositions including a bacterial collagenase as a collagen binding peptide are described in US Patent Publication No. 2010/0129341, which is hereby incorporated herein by reference in its entirety.

FIG. 1 shows a sequence alignment of the collagen-binding region of several M9B bacterial collagen-binding proteins included as SEQ ID NOs: 13-34. As can be seen from the sequence alignment, these proteins have a relatively small amount of sequence identity (about 30%), but they all bind to collagen in a similar fashion and are believed to have similar conformation as discussed in the Examples. Thus any of the peptides shown in FIG. 1 or collagen-binding fragments thereof can be used in the compositions and methods described herein. In FIG. 1, the amino acid residues critical for the conformation of the peptide and for the collagen-binding activity are underlined and shown in green and blue respectively. The key amino acid residues for collagen-binding are a tyrosine or phenylalanine at position 970 of ColG, position 977 of the ColH sequence of SEQ ID NO:1 (position 937 in FIG. 1) or a similar position of one of the sequences shown in FIG. 1; a tyrosine at position 994 of ColG, position 1000 of the ColH sequence of SEQ ID NO:1 (position 962 in FIG. 1) or a similar position of one of the sequences shown in FIG. 1; a tyrosine, phenylalanine or

histidine at position 996 of ColG, position 1002 of the ColH sequence of SEQ ID NO:1 (position 964 in FIG. 1) or a similar position of one of the sequences shown in FIG. 1. Thus a peptide with relatively low sequence identity, sharing the structure and function of the ColG protein may also be used as a collagen binding domain (CBD) herein.

In one embodiment, the collagenase is ColH, SEQ ID NO: 6. The collagen-binding polypeptide segment may be or may include residues 901-1021 of SEQ ID NO:6 (residues 34-158 of SEQ ID NO:1), or a fragment of residues 34-158 of SEQ ID NO:1 at least 8, 10, 12, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110 or 120 amino acid residues in length. The collagen-binding polypeptide segment is at least 50%, 60%, 70%, 80%, or at least 85%, at least 90%, at least 95%, at least 96%, at least 98%, or at least 99% identical to residues 34-158 of SEQ ID NO: 1. The collagen-binding polypeptide segment may be or may include residues 807-1021 of SEQ ID NO:6 (residues 37-251 of SEQ ID NO:2), or a fragment of residues 807-1021 of SEQ ID NO:6 at least 8, 10, 12, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110 or 120 amino acid residues in length. Residues 807-901 comprise the polycystic kidney disease (PKD) domain of the collagen-binding protein. Those of skill in the art will appreciate that other linkers could be used to link the collagen-binding peptide to a therapeutic agent as outlined above. The collagen-binding polypeptide segment may be or may comprise a fragment of residues 901-1021 of SEQ ID NO:6, e.g., a fragment of at least 8, at least 10, at least 20, at least 30 at least 40, or at least 50 consecutive amino acid residues of residues 901-1021 of SEQ ID NO:6. Suitably the collagen-binding polypeptide consists of residues 894-1008, 894-1021, 901-1021, or 901-1008 of SEQ ID NO: 6 or a homolog thereof as shown by the sequence alignment in FIG. 9.

Among other proteins the collagen-binding segment can be derived from are ColG (Matsushita et al., (1999) J. Bacteriol. 181:923-933), a class I collagenase from *Clostridium histolyticum*. ColH is a class II collagenase (Yoshihara et al., (1994) J. Bacteriol. 176: 6489-6496). The collagen-binding polypeptide segment may also be a polypeptide segment from any one of the protein sequences provided in FIG. 1 which aligns collagen-binding peptides from members of *Clostridium* and *Bacillus*. Those of skill in the art will appreciate that other members of this collagen-binding protein family may be useful in the methods described herein.

The therapeutic agents linked to the collagen-binding polypeptide may be any suitable pharmaceutical or other active agent, including but not limited to, osteogenic promoters, antimicrobials, anti-inflammatory agents, polypeptides such as recombinant proteins, cytokines or antibodies, small molecule chemicals or any combination thereof. Suitably the therapeutic agents are capable of promoting bone growth, decreasing inflammation, promoting collagen stability. Suitably, the therapeutic agent is one whose therapeutic effect is in the region of collagen or damaged collagen. The therapeutic agent may include, but is not limited to, bone morphogenic protein (BMP), G-CSF, FGF, BMP-2, BMP-3, FGF-2, FGF-4, anti-sclerostin antibody, growth hormone, IGF-1, VEGF, TGF- $\beta$ , KGF, FGF-10, TGF- $\alpha$ , TGF- $\beta$ 1, TGF- $\beta$  receptor, CT, GH, GM-CSF, EGF, PDGF, celineprolol, activins and connective tissue growth factors. In alternative embodiments, the active agent may be a PTH/PTHrP receptor agonist or antagonist.

Bone loss due to a collagenopathy such as osteogenesis imperfecta, Stickler's syndrome or others which put an individual at higher risk for a bone fracture due to a collagen

defect could be treated by administration of a bone anabolic peptide. The CBD may target the bone anabolic agents to sites where the collagen is malformed and thus may prevent fracture.

Vascular fragility due to defects such as Ehlers-Danlos syndrome type IV, Alport's syndrome or other diseases where blood vessel rupture is more likely due to a defect in collagen formation may be administered peptides that stimulate vascular growth or repair. The CBD will target the peptide to the areas having collagen damage and these areas are likely to have damaged vessels. The therapeutic agents will stimulate growth and repair at the site of damage and prevent vessel rupture.

Skin fragility due to disorders such as Ehlers-Danlos syndrome, Caffey's disease or other diseases where weakening of the skin due to a collagen defect leads to hyperelasticity, easy bruising or poor wound healing. Dermal and epidermal growth factors may serve as therapeutic agents which when linked to CBD and delivered to areas of damaged collagen will stimulate growth and repair of the skin, preventin striae and improving healing.

Collagen defects may also lead to cartilage malformation or insufficiency. Cartilage growth factors could be delivered locally to sites of damaged cartilage to aid in repair and restore function.

The PTH/PTHrP receptor agonist polypeptide segment may be a synthetic polypeptide or a naturally occurring polypeptide. Such polypeptides may be a portion of a polypeptide or may comprise one or more mutations. The mutations may make the PTH/PTHrP receptor agonist a better or worse agonist as compared to the wild-type PTH/PTHrP. Agonist activity with the PTH/PTHrP receptor can be assayed as described in Example 3 below by a cAMP stimulation assay. An agonist will stimulate cAMP synthesis in the assay described. Suitably, an agonist can activate receptor activity at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or even 110% or 120% as much as wild-type PTH(1-34).


The PTH/PTHrP receptor agonist polypeptide segment is a PTH or PTHrP polypeptide segment. One human isoform of PTH is SEQ ID NO:7. One human isoform of PTHrP is SEQ ID NO:8. While the human isoforms are provided, those of skill in the art will appreciate that other non-human-derived isoforms may be used as well. Such non-human-derived isoforms may be able to interact with human PTH/PTHrP receptor and vice versa. The PTH/PTHrP receptor agonist polypeptide segment may be or may include residues 1-33 of SEQ ID NO:1 (residues 1-33 of PTH (SEQ ID NO:7)). The PTH/PTHrP receptor agonist polypeptide segment may be or may include residues 1-34 of PTH (SEQ ID NO:7). In other embodiments, it is a fragment of residues 1-34 of PTH (SEQ ID NO:7). In other embodiments, the PTH/PTHrP receptor agonist polypeptide segment may be or may include residues 1-84 of PTH (SEQ ID NO:7). In other embodiments, the PTH/PTHrP receptor agonist polypeptide segment may be or may include residues 1-14 of PTH (SEQ ID NO:7). In still other embodiments, the PTH/PTHrP receptor agonist is a PTH or PTHrP polypeptide segment for any other species.

The PTH/PTHrP receptor antagonist can include in one embodiment PTH(7-34), i.e., residues 7-34 of PTH (SEQ ID NO:7). In another embodiment, it is or includes residues 7-33 of PTH (SEQ ID NO:7). In other embodiments, it is a fragment of residues 7-34 of SEQ ID NO: 8. In another embodiment, the PTH/PTHrP receptor antagonist includes PTH(7-14), i.e., residues 7-14 of PTH (SEQ ID NO:7). In another embodiment, the PTH/PTHrP receptor antagonists

include ((-1)-33) of PTH/PTHrP. In another embodiment, the PTH/PTHrP receptor antagonists include residues 1-14 of PTH with an N-terminal extension. Adding an N-terminal extension to PTH or active N-terminal fragments of PTH converts the PTH peptides to antagonists. The N-terminal extension can be 1, 2, 3, 4, 5, or more amino acids in length. The identity of the amino acids in the N-terminal extension is typically not important. In one embodiment, the PTH/PTHrP receptor antagonist includes residues 1-33 of PTH with a Gly-Ser extension at the N-terminus (SEQ ID NO:11). In another embodiment, the PTH/PTHrP receptor antagonist includes PTHrP(7-34), i.e., residues 7-34 of SEQ ID NO:8, or a fragment of residues 7-34 of SEQ ID NO:8. In another embodiment, the PTH/PTHrP receptor antagonist includes mouse TIP(7-39) (See Hoare S R, Usdin T B. 2002. Specificity and stability of a new PTH1 receptor antagonist, mouse TIP(7-39). Peptides 23:989-98.). Other PTH/PTHrP receptor antagonists that may be used in the fusion proteins are also disclosed in Hoare et al. The PTH/PTHrP receptor antagonist may be a fragment of at least 8, 10, 12 or more amino acids from residues 1-34 of SEQ ID NO:7. In other embodiments the PTH/PTHrP receptor antagonist may be PTH/PTHrP receptor antagonist polypeptide from another species.

In one embodiment, the therapeutic agent or PTH/PTHrP receptor agonist or antagonist polypeptide segment is N terminal to the collagen-binding polypeptide segment in the fusion protein. That is, the two polypeptide segments each have an N-terminal and a C-terminal, and the N-terminal of the collagen-binding polypeptide segment is linked directly or indirectly, e.g., through a linker polypeptide segment (such as PKD, a Glycine or Serine linker) to the C-terminal of the therapeutic agent or PTH/PTHrP agonist or antagonist polypeptide segment.

The fusion proteins described above comprising (a) a collagen-binding polypeptide segment linked to (b) a therapeutic agent or a PTH/PTHrP receptor agonist or antagonist polypeptide segment can be replaced by pharmaceutical agents comprising (a) a collagen-binding polypeptide segment linked to (b) a therapeutic agent or PTH/PTHrP receptor agonist or a non-peptidyl PTH/PTHrP receptor agonist. An example of a non-peptidyl PTH/PTHrP receptor agonist is compound AH3960 (Rickard et al., (2007) Bone 39:1361-1372).



AH3960

AH3960 contains two amino groups. Amino groups in small chemical molecules such as AH3960 can be used to cross-link the therapeutic agent to amino groups on the collagen-binding polypeptide segment through a cross-linker such as DSG (disuccinimidyl glutarate) or through the combination of SANH (succinimidyl-4-hydrazinonicotinate acetone hydrazone) and SFB (succinimidyl-4-formyl benzoate). Therapeutic agents can be cross-linked through their amino group to a carboxyl group of the collagen-binding

polypeptide segment by EDC (1-ethyl-3-[3-dimethylamino-propyl]carbodiimide hydrochloride) or vice versa. These cross-linking products are available from Pierce (pierce.net.com, Thermo Fisher Scientific Inc., Rockford, Ill.). Protocols and reaction conditions are also available in the product literature from Pierce (pierce.net.com).

In another embodiment of the pharmaceutical agents comprising (a) a collagen-binding polypeptide segment; linked to (b) a polypeptide therapeutic agent or a PTH/PTHrP receptor agonist or antagonist polypeptide segment, segment (a) and segment (b) are separate polypeptides, and the two polypeptides are linked by chemical cross-linking. The two polypeptides can be cross-linked through amino groups by reagents including DSG (disuccinimidyl glutarate) or glutaraldehyde. They can also be cross-linked through amino groups by derivatizing one polypeptide with SANH (succinimidyl-4-hydrazinonicotinate acetone hydrazone) and the other with SFB (succinimidyl-4-formyl benzoate), and then mixing the two derivatized polypeptides to cross-link. The two polypeptides can be cross-linked between an amino group of one polypeptide and a carboxyl of the other by reaction with EDC (1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride). The polypeptides can also be cross-linked (e.g., covalently coupled) by any other suitable method known to a person of ordinary skill in the art. These cross-linking reagents are available from Pierce (pierce.net.com, Thermo Fisher Scientific Inc., Rockford, Ill.). Protocols and reaction conditions are also available in the product literature from Pierce (pierce.net.com). These and other applicable cross-linking methods are described in U.S. published patent applications 2006/0258569 and 2007/0224119.

Also provided herein are methods of treating hyperparathyroidism by administering PTH-CBD to a subject in need of treatment for hyperparathyroidism. In one embodiment the PTH administered to the subject may be a PTH from a different species. As shown in the Examples a single administration of CBD-PTH to ovariectomized aged rats was able to reduce the amount of endogenous PTH produced by the animal. Thus, administration of PTH-CBD to individuals suffering from hyperparathyroidism may experience a decrease in symptoms associated with hyperparathyroidism and have decreased levels of PTH after administration of PTH-CBD.

The effects of PTH agonists and antagonists on hair growth have been studied for over almost 15 years. PTH has a common receptor with PTH-related peptide (PTHrP), which is normally produced by dermal fibroblasts. PTHrP affects keratinocyte proliferation/differentiation and modulates the hair cycle. Most of the testing on hair growth effects has been performed with PTH antagonists, as indications from initial testing were that these were the most effective agents. Both injected and topical formulations have been tested in animal models of chemotherapy-induced alopecia and in the SKH-1 hairless mouse. Part of the effect of PTH antagonists on hair growth is to transition the hair follicles into a dystrophic catagen stage, which protects them from chemotherapeutic damage. However, clinical trials of topical PTH antagonists for chemotherapy-induced alopecia by IGI Pharmaceuticals were discontinued in phase 2 because of limited efficacy. Thus new compositions for treating alopecia are needed.

The problems of delivery and retention of PTH to the skin can be overcome by using collagen-targeted PTH analogs. To accomplish this, we synthesized several fusion proteins of different PTH agonists and antagonists linked to a collagen binding domain derived from the ColH1 collagenase of

*Clostridium histolyticum*. In the studies described in the Examples, we found that the agonist compound PTH-CBD promotes transition of hair follicles to the anagen phase and has potent effects on hair growth. The antagonist compound PTH(7-33)-CBD had little effect on hair growth in chemotherapy models and had a deleterious effect on hair regrowth after depilation. Compounds such as PTH-CBD, which promote anagen phase transition of hair follicles, have been sought after due to their potential to treat a large variety of disorders of hair loss. PTH-CBD appears to have a similar mechanism of action to cyclosporine, which also promotes transition of hair follicles to anagen phase, although the mechanism is less likely to be the result of direct effects on WNT signaling. While clinical use of cyclosporine for this purpose is limited by systemic toxicity, PTH-CBD has not shown toxic effects, even with systemic administration.

Thus in another aspect, methods of increasing hair growth are provided herein. The methods include administering a CBD linked to a PTH/PTHrP receptor agonist to a subject in need of treatment to induce hair growth or stop hair loss. The method is applicable to individuals with alopecia, including chemotherapy induced alopecia, but also alopecia areata, alopecia caused by male pattern baldness, polycystic ovarian syndrome or other hair loss. The compositions may be administered locally or topically to treat hair loss.

In another aspect, methods of slowing hair growth or regrowth after a hair removal procedure by administering a CBD linked to a PTH/PTHrP receptor antagonist to a subject are provided. In one embodiment, the PTH antagonist composition is applied locally, topically. The PTH antagonist may be applied after a hair removal procedure to prevent or slow hair regrowth. As described in the Examples, we have demonstrated that hair regrowth is slowed after waxing in animals treated with CBD-PTH antagonist as compared to control animals treated with PTH-CBD or vehicle alone. The compositions may be administered locally or topically to block hair growth.

The compositions described herein may be administered by any means known to those skilled in the art, including, but not limited to, oral, topical, intranasal, intraperitoneal, parenteral, intravenous, intramuscular, intradermal or subcutaneous. Thus the compositions may be formulated as an ingestible, injectable, topical or suppository formulation. The composition may be formulated for administration by injection to result in systemic administration or local administration. The compositions may also be delivered with in a liposomal or time-release vehicle. The compositions may also be delivered in a site-directed delivery vehicle, such as but not limited to, a targeted liposome or an absorbable collagen sponge carrier or other implant.

The inventors have found that when administering compositions including a CBD subcutaneously it binds locally at the site of injection if the composition is dissolved in neutral pH buffer. But if the composition is dissolved in a low pH buffer, for example a buffer having pH 5.0 or pH 4.5 or below, the collagen-binding domain does not bind collagen, and the composition has time to disperse systemically before it binds collagen elsewhere in the body at neutral pH. Thus systemic administration of the compositions involves administering the composition dissolved in buffer or aqueous solution at a pH lower than about 5.0 or at pH 4.5 or below. In another embodiment, systemic administration of the compositions involves administering the fusion proteins dissolved in aqueous solution at pH lower than about 6.0. Alternatively, if the skin condition is localized, the compositions described herein may be administered in a buffer with

a pH of 6.0, 6.5, 7.0, 7.5 or above in order to allow for localized delivery of the compositions to the affected area of the skin.

Pharmaceutical compositions for topical administration may also be formulated using methods and compositions such as those available to those skilled in the art. For example, gels, creams or liposome preparations may be suitable for topical delivery. These delivery vehicles may be formulated to mediate delivery to the lower layers of the skin or to allow for extended release of the pharmaceutical at the site of application.

The compositions can be administered as a single dose or as divided doses. For example, the composition may be administered two or more times separated by 4 hours, 6 hours, 8 hours, 12 hours, a day, two days, three days, four days, one week, two weeks, or by three or more weeks. Optionally, such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. The composition is expected to be more effective than a comparable or control composition comprising the therapeutic agent or a PTH/PTHrP receptor agonist that is not linked to a collagen-binding protein. In one embodiment, a smaller amount of the composition may be used or the composition may be administered less frequently than a comparable composition comprising the therapeutic agent or a PTH/PTHrP receptor agonist which is not linked to a collagen-binding protein.

The dosage amounts and frequencies of administration provided herein are encompassed by the terms therapeutically effective and prophylactically effective. The individual doses of pharmaceutical agents comprising a collagen-binding polypeptide segment linked to a therapeutic agent may be approximately the same on a molar basis as doses used for the therapeutic agent alone. It is expected that the pharmaceutical agents comprising a collagen-binding polypeptide segment linked to a therapeutic agent may be administered less frequently, because linking the agent to the collagen-binding polypeptide segment gives it much more prolonged activity in vivo.

Administration of the compositions to a subject in accordance with the invention appears to exhibit beneficial effects in a dose-dependent manner. Thus, within broad limits, administration of larger quantities of the compositions is expected to achieve increased beneficial biological effects than administration of a smaller amount. Moreover, efficacy is also contemplated at dosages below the level at which toxicity is seen.

It will be appreciated that the specific dosage administered in any given case will be adjusted in accordance with the compositions being administered, the disease to be treated or inhibited, the condition of the subject, and other relevant medical factors that may modify the activity of the agent or the response of the subject, as is well known by those skilled in the art. For example, the specific dose for a particular subject depends on age, body weight, general state of health, diet, the timing and mode of administration, the rate of excretion, medicaments used in combination and the severity of the particular disorder to which the therapy is applied. Dosages for a given patient can be determined using conventional considerations, e.g., by customary comparison of the differential activities of the compositions of the invention and of the therapeutic agent administered alone, such as by means of an appropriate conventional pharmacological or prophylactic protocol.

The maximal dosage for a subject is the highest dosage that does not cause undesirable or intolerable side effects. The number of variables in regard to an individual prophylactic or treatment regimen is large, and a considerable range of doses is expected. The route of administration will also

impact the dosage requirements. It is anticipated that dosages of the compositions will reduce symptoms of the condition being treated by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% compared to pre-treatment symptoms or symptoms is left untreated. It is specifically contemplated that pharmaceutical preparations and compositions may palliate or alleviate symptoms of the disease without providing a cure, or, in some embodiments, may be used to cure the disease or disorder.

Suitable effective dosage amounts for administering the compositions may be determined by those of skill in the art, but typically range from about 1 microgram to about 10,000 micrograms per kilogram of body weight weekly, although they are typically about 1,000 micrograms or less per kilogram of body weight weekly. In some embodiments, the effective dosage amount ranges from about 10 to about 10,000 micrograms per kilogram of body weight weekly. In another embodiment, the effective dosage amount ranges from about 50 to about 5,000 micrograms per kilogram of body weight weekly. In another embodiment, the effective dosage amount ranges from about 75 to about 1,000 micrograms per kilogram of body weight weekly. The effective dosage amounts described herein refer to total amounts administered, that is, if more than one compound is administered, the effective dosage amounts correspond to the total amount administered.

The effectiveness of the compositions described herein may be enhanced by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% relative to a control treated with the therapeutic agent alone. It will be appreciated that the effectiveness of the treatment in any given case will be enhanced variably in accordance with the specific compositions used, the type of disease being treated, the condition of the subject, the specific formulations of the compounds and other relevant medical factors that may modify the activity of the compositions or the responses of the subject as is appreciated by those of skill in the art.

The following examples are meant only to be illustrative and are not meant as limitations on the scope of the invention or of the appended claims. All references cited herein are hereby incorporated by reference in their entireties.

## EXAMPLES

### Example 1

#### CBD Targets Partially Untwisted or Undertwisted Regions of Collagen

*Clostridium histolyticum* collagenase causes extensive degradation of collagen in the connective tissue resulting in gas gangrene. The C-terminal collagen-binding domain (CBD) of these enzymes is the minimal segment required to bind to the collagen fibril. CBD binds unidirectionally to the partially untwisted C-terminus of triple helical collagen. Whether CBD could also target under-twisted regions even in the middle of the collagen triple helix was examined. Partially untwisted collagenous peptides were synthesized by introducing a Gly→Ala substitutions into the collagen  $[(\text{POG})_x\text{POA}(\text{POG})_y]_3$  where  $x+y=9$  and  $x>3$ .  $^1\text{H}$ - $^{15}\text{N}$  heteronuclear single quantum coherence nuclear magnetic resonance (HSQC NMR) titration studies with  $^{15}\text{N}$ -labeled CBD demonstrated that the untwisted mini-collagen binds to a 10 Å wide 25 Å long cleft. Six untwisted collagenous peptides each labeled with a nitroxide radical were then titrated with  $^{15}\text{N}$ -labeled CBD. The paramagnetic nuclear spin relaxation

effects showed that CBD binds close to either the Gly→Ala substitution site or to the C-terminus of each mini-collagen. Small angle X-ray scattering (SAXS) measurements revealed that CBD prefers to bind the Gly→Ala site rather than the C-terminus. The HSQC NMR spectra of <sup>15</sup>N-labeled mini-collagen and untwisted mini-collagen were unaffected by the titration of unlabeled CBD. The results imply that CBD binds to the partially unwound region of the mini-collagen but does not actively unwind the triple helix. Materials and Methods:

<sup>15</sup>N-Labeled Protein Production: The sib (Gly893-Lys1008) peptide derived from *Clostridium histolyticum* class I collagenase (ColG) was expressed as a glutathione S-transferase (GST)-fusion protein. The GST-tag was cleaved off by thrombin, and CBD was purified as described previously. Matsushita, et al., (2001) *J Biol Chem* 276, 8761-8770. Uniform <sup>15</sup>N isotope labeling was achieved using Tanaka minimal medium containing 40 mM <sup>15</sup>NH<sub>4</sub>Cl. The labeling efficiency was estimated to be 99.6% by matrix-assisted laser desorption/ionization-time of flight mass spectrometry (MALDI-TOF-MS).

Peptides: (POG)<sub>10</sub> (SEQ ID NO: 35) was purchased from Peptide Institute, Inc. (Osaka, Japan). Other peptides were constructed by a standard N-(9-fluorenyl) methoxycarbonyl (Fmoc)-based strategy on Rink-amide resins (Novabiochem, Darmstadt, Germany). N-terminal spin-labeling was performed on the resin by the treatment with 5 equivalents of 3-carboxy-PROXYL (Aldrich), 1-hydroxybenzotriazole, diisopropylcarbodiimide in N,N-dimethylformamide at room temperature for 2 hours. Peptide cleavage and de-protection steps were performed by a treatment with a standard trifluoroacetic acid (TFA) scavenger cocktail (TFA: m-cresol:thioanisole:water:triisopropylsilane=82.5:5:5:5:2.5, v/v). The spin-labeling at Cys residues was performed using 3-(2-iodoacetamido)-PROXYL (IPSL, Sigma-Aldrich). Briefly, 10 molar excess of IPSL dissolved in ethanol was added to the same volume of 10 mg/ml peptide in 0.1 M Tris-HCl (pH 8.8), 5 mM ethylenediaminetetraacetic acid. After reacting at room temperature for 1 hr, the reaction was quenched by adding excess dithiothreitol. All peptides were purified by reverse-phase HPLC using a Cosmosil 5C<sub>18</sub> AR-II column (Nacalai Tesque, Kyoto, Japan) and characterized by MALDI-TOF-MS. All the measured masses agreed with the expected values. The chemical structures of synthesized peptides are shown in FIG. 2.

Circular Dichroism Spectroscopy: The triple helical conformation and the stability of the collagenous peptides were verified using CD spectroscopy (See FIGS. 3 and 4). CD spectra were recorded with a J-820 CD spectropolarimeter (JASCO Co., Hachioji, Japan) equipped with a Peltier thermo controller, using a 0.5-mm quartz cuvette and connected to a data station for signal averaging. All peptide samples were dissolved in water (1 mg/ml), and stored at 4° C. for 24 h. The spectra are reported in terms of ellipticity units per mole of peptide residues  $[\theta]_{mwr}$ . Thermostability of the triple helix was monitored by the  $[\theta]_{225}$  values of each peptide with increasing temperature at the rate of 0.3° C./min.

NMR Spectroscopy: NMR experiments were performed on a Bruker 700 MHz spectrometer equipped with Cryo-probe™. All the NMR titration experiments were carried out at 16±0.5° C. The working temperature is lower than the melting temperatures ( $T_m$ ) of all the paramagnetic spin-labeled collagenous peptides (Table 1) used. The concentration of the protein was 0.1 mM in 50 mM Tris-HCl (pH 7.5) containing 100 mM NaCl and 20 mM CaCl<sub>2</sub>. The dilution effect on the course of titration was minimized by the titration of a highly concentrated (4 mM) peptide stock. Aliquots of collagenous peptide were added to the protein and equilibrated for 5 min before acquiring <sup>1</sup>H-<sup>15</sup>N HSQC

spectra. The pH of the NMR samples monitored during the titration exhibited no significant shift in the pH (within ±0.2 units).

TABLE 1

Melting temperatures ( $T_m$ ) of various mini-collagen peptides that were used in NMR titration and the experiments described herein.		
Peptides	$T_m$ (° C.)	SEQ ID NO:
(POG) <sub>4</sub> POA(POG) <sub>5</sub>	29	38
PROXYL-(POG) <sub>4</sub> POA(POG) <sub>5</sub>	29	38
PROXYL-(POG) <sub>3</sub> POA(POG) <sub>6</sub>	28	39
PROXYL-(POG) <sub>5</sub> POA(POG) <sub>4</sub>	28	37
PROXYL-(POG) <sub>6</sub> POA(POG) <sub>3</sub>	27	36
(POG) <sub>4</sub> POA(POG) <sub>5</sub> C-PROXYL	30	41
<sup>11</sup> PROXYL-(POG) <sub>3</sub> PCG(POG) <sub>4</sub>	28	40

Dynamic Light Scattering Experiments: The dynamic light scattering (DLS) data were collected using DynaPro-E equipped with a temperature controlled micro-sampler on the samples of CBD, collagenous peptides and CBD:mini-collagen complexes in 10 mM Tris-HCl (pH 7.5) containing 100 mM sodium chloride and 20 mM CaCl<sub>2</sub>. The protein samples were spun at 10,000 rpm for 10 min and were filtered through 0.02 μm Whatman syringe directly into a 50-μL quartz cuvette. For each experiment, 20 measurements were made. The mean hydrodynamic radius ( $R_H$ ), standard deviation, polydispersity, and percent of peak area were analyzed using Dynamics V6 (Protein Solutions). The hydrodynamic radius and molecular weight estimations were calculated from time dependent fluctuations induced by Brownian motion as described. Proteau, et al. (2010) *Curr Protoc Protein Sci* Chapter 17, Unit 17 10.

Small Angle X-Ray Solution Scattering Experiments: The small angle X-ray solution scattering (SAXS) data were collected on solutions of CBD, collagenous peptides and CBD-mini-collagen complexes in 10 mM Tris-HCl (pH 7.5), 100 mM NaCl and 20 mM CaCl<sub>2</sub> at SAXS/WAXS setup located at the 5-ID-D beamline of the DND-CAT synchrotron research center, Advanced Photon Source, Argonne National Laboratory (Argonne, Ill.). The main advantage of X-ray scattering is that it can be carried out in solution in near physiological conditions. Petoukhov et al., (2007) *Curr Opin Struct Biol* 17, 562-571. 1.2398 Å (10 keV) radiation was selected from the APS Undulator A insertion device using a Si-111 monochromator, with 1:1 horizontal focusing and higher harmonic rejection from a Rh coated mirror, and beam defining slits set at 0.3 mm vertical by 0.25 mm horizontal. A 1.6 mm diameter capillary flow-cell with a flow rate of 4 μl/sec was used to collect four frames with 10 second exposure time. The SAXS detector used was a Mar165 scintillator fiber-optic coupled CCD detector and covered the momentum transfer range 0.005<q<0.198 Å<sup>-1</sup>, where  $q=4\pi \sin \theta/\lambda$  (2θ is the scattering angle). The WAXS detector was a custom Roper scintillator fiber-optic coupled CCD detector and covered 0.191<q<1.8 Å<sup>-1</sup> S. Weigand, et al. (2009) *Advances in X-ray Analysis* 52, 58-68.

All scattering data were acquired at sample temperature of 10° C. The four scattering patterns from each detector were averaged and merged with the rejection of outlying scans. For further analysis the program IGOR Pro 5.5 A (WaveMetrics) was used. The scattering profiles of the protein, peptide and their complexes were obtained after subtracting the buffer profiles. The reduced scattering data were plotted as scattering intensity I(Q) vs. Q (FIG. 4A). The radius of gyration,  $R_g$ , was obtained from the Guinier approximation by linear least squares fitting in the  $QR_g < 1$  region, where the forward scattering intensity I(0) is proportional to the

molecular weight of the protein complex. An indirect Fourier transformation of I(Q) data using GNOM provided the particle distribution function P(r) in the real space (FIG. 4B). Svergun, D. (1992) *J Appl Crystallogr* 25, 495-503. Where P(r) intersects with x-axis represents the maximum diameter  $D_{max}$  averaged in all orientations. The molecular envelopes were constructed for all the samples based on the SAXS data after ab initio calculation with the program GASBOR. Svergun, et al. (2001) *Biophys J* 80, 2946-2953. Simulated annealing minimization of randomly distributed dummy atoms converged to the protein structure after being tested for the best fit to the I(Q) scattering data. No symmetry restraints were applied to any of the shape reconstructions. For each of the complexes, ten ab initio models were calculated with GASBOR and averaged using DAMAVER. Svergun, D. (2003) *J Appl Crystallogr* 36. The atomic models represented as a compact interconnected configuration of beads with diameter  $D_{max}$  were adjusted to fit the experimental data  $I_{exp}(s)$  to minimize error. Atomic models were docked into ab initio envelopes with the program SUBCOMB. Kozin, M. B., and Svergun, D. (2000) *J Appl Crystallogr* 33, 775-777.

Docking Model: The CBD-collagenous peptide complex is generated from Protein Data Bank entries of ColG s3b (1NQD) and partially untwisted collagenous peptide 1CAG (Ala mutation in 15<sup>th</sup> position). Other untwisted mini collagen molecules were generated by modifying 1CAG using fragments derived from [(POG)<sub>10</sub>]<sub>3</sub> structure (1K6F). To obtain the complex, the soft docking algorithm BiGGER was used. Palma, et al. (2000) *Proteins* 39, 372-384. Solutions were filtered using NMR titration data and the highest scoring model that satisfied NMR and SAXS results was chosen. The manual adjustments were aided by the use of MIFit. McRee. (1999) *J Struct Biol* 125, 156-165.

Results and Discussion:

<sup>1</sup>H-<sup>15</sup>N HSQC NMR Titration-CBD Targeting the Under-Twisted Sites in Collagen: The untwisted collagenous peptide [(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 36) that has Ala in the 21<sup>st</sup> position from the N-terminus was synthesized. This peptide was further modified to accommodate a paramagnetic spin label at the N-terminus. <sup>1</sup>H-<sup>15</sup>N HSQC NMR titrations were performed with [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 36) and <sup>15</sup>N-labeled CBD at ratios ranging from 0.02:1 to 1.5:1. As demonstrated earlier, a total of eleven residues on the collagen binding interface (S928, W956, G971, K995, Y996, L924, T957, Q972, D974, L991 and V993) either disappeared from the HSQC spectrum or exhibited significant chemical shift perturbation from their original position on the course of titration. Philominathan, et al. (2009) *J Biol Chem* 284, 10868-10876. The PROXYL group on the N-terminus of the collagenous peptides can cause a distance-dependent line broadening of the NMR

signals of CBD during the course of titration. In addition to the eleven residues, three more residues, V973, G975 and S979 exhibited appreciable line broadening and these residues eventually disappeared from the <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of CBD (FIGS. 5A and 5B). When the [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 36):CBD complex was reduced with ascorbic acid those three residues reappeared in the <sup>1</sup>H-<sup>15</sup>N HSQC spectrum. The disappearance of these three residues was consistent with the titration of [PROXYL-G(POG)<sub>7</sub>]<sub>3</sub> (SEQ ID NO: 42) (C-terminus is at 22<sup>nd</sup> position from the N-terminal PROXYL) in our earlier publication. The comparison of the two titration results demonstrates that CBD is targeting the Gly→Ala substituted site. If CBD had only bound to the C-terminus of [PROXYL-(POG)<sub>6</sub>POA(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 36) (C-terminus is at 30<sup>th</sup> position from the N-terminal PROXYL), we would expect to observe the disappearance of only one residue (V973) at the most, as in the published titration of [PROXYL-G(POG)<sub>7</sub>(PRG)]<sub>3</sub> (SEQ ID NO: 43). The disappearance of the residues (V973, G975 and S979) located at distal side from the Ca<sup>2+</sup> binding site (FIG. 5C) confirmed that CBD binds unidirectionally to untwisted collagen as well. The collagen binding surface in CBD is a 10-Å-wide and 25-Å-long cleft. The width of the binding cleft in CBD matches the diameter of the triple helix and its length could accommodate [(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 44). NMR results imply that CBD is binding to the under-twisted [(POG)<sub>2</sub>POA]<sub>3</sub> (SEQ ID NO: 45) region of the collagen.

As paramagnetic relaxation enhancement is a distance dependent phenomenon, Gly→Ala substitution made at closer to the N-terminal PROXYL group should result in the disappearance of more residues on CBD. PROXYL containing collagenous peptides, [PROXYL-(POG)<sub>5</sub>POA(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 37) (Ala at 18<sup>th</sup> position from the N-terminal PROXYL), [PROXYL-(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub> (SEQ ID NO: 38) (Ala at the 15<sup>th</sup> position from the PROXYL) and [PROXYL-(POG)<sub>3</sub>POA(POG)<sub>6</sub>]<sub>3</sub> (SEQ ID NO: 39) (Ala at the 12<sup>th</sup> position from the PROXYL) were synthesized. Just as in the previous titrations, the line broadening effects on the residues of CBD were analyzed from the changes in the <sup>1</sup>H-<sup>15</sup>N HSQC spectrum. The shorter the distance between Gly→Ala substitution site and the N-terminal PROXYL, more residues in CBD disappeared (FIG. 6 and Table 2). The magnitude of intensity drop for four amide resonances (Q972, G975, S979 and L924) of four different mini-collagen molecules was also the function of the distance (FIG. 7). The NMR results are consistent with CBD binding to the [(POG)<sub>2</sub>POA]<sub>3</sub> (SEQ ID NO: 45) region in each of the four under-twisted mini-collagen. The binding constants obtained from all the NMR titrations were <100 μM indicating a moderate binding affinity between CBD and under-twisted mini-collagen.

TABLE 2

Residues that disappear due to the presence of PROXYL either at the N-terminus, C-terminus or in the middle of the collagenous peptide sequence.			
No.	Peptides	Alanine position	Residues disappeared due to PROXYL
	Blank		
1	[(POG) <sub>10</sub> ] <sub>3</sub> (SEQ ID NO: 35)		
	PROXYL at N-terminus		
2	[PROXYL-(POG) <sub>6</sub> POA(POG) <sub>3</sub> ] <sub>3</sub> (SEQ ID NO: 36)	21	V973, G975, S979

TABLE 2-continued

Residues that disappear due to the presence of PROXYL either at the N-terminus, C-terminus or in the middle of the collagenous peptide sequence.			
No.	Peptides	Alanine position	Residues disappeared due to PROXYL
3	[PROXYL-(POG) <sub>5</sub> POA(POG) <sub>4</sub> ] <sub>3</sub> (SEQ ID NO: 37)	18	Q972, V973, G975, S979
4	[PROXYL-(POG) <sub>4</sub> POA(POG) <sub>5</sub> ] <sub>3</sub> (SEQ ID NO: 38)	15	L946, Q972, V973, G975, S979
5	[PROXYL-(POG) <sub>3</sub> POA(POG) <sub>6</sub> ] <sub>3</sub> (SEQ ID NO: 39)	12	L946, G953, Q972, V973, D974, G975, N976, V978, S979
<b>PROXYL at C-terminus</b>			
6	[(POG) <sub>4</sub> POA(POG) <sub>5</sub> -PROXYL] <sub>3</sub> (SEQ ID NO: 41)	15	S906, R929, S997, G998
<b>PROXYL in the middle</b>			
7	[11PROXYL-(POG) <sub>3</sub> PCG(POG) <sub>4</sub> ] <sub>3</sub> (SEQ ID NO: 40)		V973, G975, S979 and S906, R929, S997, G998

The helical conformation in both the [(POG)<sub>2</sub>POA]<sub>3</sub> (SEQ ID NO: 45) and the C-terminal [(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 44) are similarly under-wound. The degree of rotation about the screw axis symmetry that describes the internal triple helical twist is defined as the helical twist value  $\kappa$ . The  $\kappa$ -value oscillates around an average value of  $-103^\circ$  for [(POG)<sub>10</sub>]<sub>3</sub> (SEQ ID NO: 35). Bella (2010) *J Struct Biol* 170, 377-391. The C-terminus of a mini-collagen is under-twisted ( $\kappa$  value shifts from  $-103^\circ$  to  $-110^\circ$ ) but the N-terminus is usually over-twisted. Collagen peptides with Gly→Ala substitution in the center of the peptide sequence still form triple helices, but with an abrupt under-twisting ( $\kappa$  value shifts from  $-103^\circ$  to  $-115^\circ$ ) at the substitution site followed over-twisting to the norm. Because the [(POG)<sub>2</sub>POA]<sub>3</sub> (SEQ ID NO: 45) region is somewhat more under-twisted than C-terminal [(POG)<sub>3</sub>]<sub>3</sub> (SEQ ID NO: 44), the former could be preferentially targeted by CBD than the latter. However, CBD could still bind to the C-terminus.

CBD Also Targets the C-Terminus of the Under-Twisted Mini-Collagen: To demonstrate that CBD binds to the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) as well, a collagenous peptide [(POG)<sub>4</sub>POA(POG)<sub>5</sub>-PROXYL]<sub>3</sub> (SEQ ID NO: 38) was synthesized. [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-PROXYL]<sub>3</sub> (SEQ ID NO: 41) was titrated with <sup>15</sup>N-labeled CBD at ratios 0.02:1 to 1.5:1 with increments of 0.02, and the changes in the HSQC spectrum of CBD were monitored. When the mini-collagen was bound to the cleft, a total of eleven residues on the collagen binding interface either line broadened or showed significant chemical shift perturbation as described earlier. Philominathan, et al. (2009) *J Biol Chem* 284, 10868-10876. Four additional residues S906, R929, S997 and G998 disappeared from the HSQC spectrum due to PROXYL (FIGS. 6 A, B and D). These peaks reappeared upon addition of ascorbic acid. This phenomenon is identical to our previous titration of [GPRG(POG)<sub>7</sub>C-PROXYL]<sub>3</sub> (SEQ ID NO: 46) when CBD bound the C-terminus. If CBD were to bind only to the partially unwound Ala site, we would have observed the disappearance of fewer residues. Thus in addition to targeting the (POG)<sub>2</sub>POA (SEQ ID NO: 45) region of the collagenous peptide, CBD also binds to the C-terminal (POA)<sub>3</sub> (SEQ ID NO: 44). As described, the

helical confirmation of both the (POG)<sub>2</sub>POA (SEQ ID NO: 45) region and the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) are similarly under-twisted compared to the norm. Bella. (2010) *J Struct Biol* 170, 377-391. Our current explanation for why CBD is targeting the under-twisted regions is that the partial unwinding positions main-chain carbonyl groups to favor hydrogen-bonding interactions with the hydroxyl group of Tyr994. Tyr994 mutation to Phe resulted in 12-fold reduction in binding to mini-collagen, and the mutation to Ala lost binding capability. Wilson, et al. (2003) *EMBO J* 22, 1743-1752.

To demonstrate CBD's ability to target both the (POG)<sub>2</sub>POA (SEQ ID NO: 45) region and the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) region, a collagenous peptide [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 40) modified to accommodate PROXYL group in the middle (11<sup>th</sup> position) was synthesized. PROXYL group is covalently joined to the cysteine residue. Due to the presence of the bulky PROXYL group, this peptide is expected to be partially untwisted. The precise degree of under-twisting is not known for the peptide, but mini-collagen with GPX repeats exhibits a moderate under-twisting ( $\kappa=-105^\circ$ ). Bella. (2010) *J Struct Biol* 170, 377-391. The bulky PROXYL group will likely induce greater untwisting than  $\kappa=-105^\circ$ . In addition to the eleven amide resonances either line-broaden or shifted, <sup>1</sup>H-<sup>15</sup>N HSQC NMR titrations revealed two distinct phenomena. At lower ratio (0.2:1) amide resonances corresponding to S906, R929, S997, and G998 disappeared from the HSQC spectrum of CBD (FIGS. 8E, F and H). Then at higher ratio (0.3:1), additional amide resonances corresponding to V973, G975 and S979 disappeared from the HSQC spectrum of CBD (FIGS. 8E, F and H). In order to cause the disappearance of four residues (S906, R929, S997 and G998), CBD must initially bind to the N-terminal (POG)<sub>3</sub> (SEQ ID NO: 44). The disappearance of resonances V973, G975 and S979 can be explained if CBD binds to the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) of the mini-collagen. However the initial phenomenon signifies that CBD binds preferentially to the under-twisted mid-section to C-terminus.

## 21

To demonstrate that PROXYL caused the line broadening and Ala or Cys residues did not, three more control peptides, [(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub> (SEQ ID NO: 38), [(POG)<sub>4</sub>POA(POG)<sub>5</sub>C-carbamidomethyl]<sub>3</sub> (SEQ ID NO: 41), and [(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 40) that lack the PROXYL groups were synthesized, and NMR titrations were repeated (FIGS. 6F, 8C and 8G, respectively). The titration results were nearly identical with those of [(POG)<sub>10</sub>]<sub>3</sub> (SEQ ID NO: 35). Only the eleven amide resonances were either line broadened or shifted even at 1:1 (mini-collagen:CBD) ratio. These control peptides bound to the same cleft, and PROXYL caused the additional residues to line broaden.

To illustrate if CBD binds only to the partially untwisted site in the middle of the collagen peptide and/or to the C-terminus of mini-collagen, dynamic light scattering experiments (DLS) were performed. DLS experiments provided the stoichiometries of collagen:CBD complexes. The hydrodynamic radius of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>-PROXYL]<sub>3</sub> (SEQ ID NO: 38):CBD and [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 40):CBD was 3 nm and the apparent molecular weight of the complex was 42±1 kDa, which is similar to those observed for [(POG)<sub>10</sub>]<sub>3</sub> (SEQ ID NO: 35):CBD complex (Table 3). Other complexes also exhibited similar values. Thus far, all the mini-collagen and CBD always formed 1:1 complex. CBD binds to either one of the available sites in mini-collagen but does not occupy both sites to form a 1:2 complex.

## 22

Small Angle X-Ray Scattering Experiments (SAXS): The three dimensional molecular shapes of the CBD-collagenous peptide complexes were constructed using SAXS measurements. The main advantage of SAXS measurements is that the experiments are performed in solution under near physiological conditions. In our previous work, these three dimensional molecular envelopes were used to demonstrate asymmetric binding of CBD to the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) of mini-collagen. The molecular shapes were constructed for complexes of CBD and six different untwisted mini-collagen molecules. In all cases CBD bound to (POG)<sub>2</sub>POA (SEQ ID NO: 45) region preferentially to C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) (FIGS. 9A-F). For example the docking model for CBD:[(POG)<sub>4</sub>POA(POG)<sub>5</sub>]<sub>3</sub> (SEQ ID NO: 38) constructed using the crystal structure of CBD (pdb accession code 1NQG) interacting with (POG)<sub>2</sub>POA (SEQ ID NO: 45) region of the untwisted collagen (pdb accession code 1CAG) fit the envelope well (FIG. 9B). Although NMR results demonstrated that CBD also binds to the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) of [(POG)<sub>4</sub>POA(POG)<sub>5</sub>-PROXYL]<sub>3</sub> (SEQ ID NO: 38), CBD predominantly binds to the (POG)<sub>2</sub>POA (SEQ ID NO: 45) region of the peptide (FIGS. 9E and 9F).

Structures derived from SAXS profiles using simulated annealing calculations for [11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 40) (FIGS. 9G and 9H) indicated an additional density that could be attributed to the PROXYL group. The SAXS derived three-dimensional shape of

TABLE 3

Hydrodynamic radius (RH), apparent molecular weight (Mw), Radius of gyration (Rg) and Maximum particle diameter (Dmax) computed from Dynamic light scattering (DLS) and small angle X-ray scattering (SAXS) for various CBD:collagenous peptides complexes.					
No	Complexes	Dynamic Light Scattering (DLS)		Small Angle X-ray Scattering (SAXS)	
		Hydro-	Apparent	Radius of Gyration (Rg)	Max Diameter (Dmax)
		dynamic Radius (RH)	Molecular Weight (Mw)		
1	CBD:[(POG) <sub>10</sub> ] <sub>3</sub> (SEQ ID NO: 35)	3	43	22.62 ± 0.04	93
2	CBD:[PROXYL-(POG) <sub>6</sub> POA(POG) <sub>3</sub> ] <sub>3</sub> (SEQ ID NO: 36)	3	44	24.67 ± 0.09	87
3	CBD:[PROXYL-(POG) <sub>5</sub> POA(POG) <sub>4</sub> ] <sub>3</sub> (SEQ ID NO: 37)	3	42	21.08 ± 0.02	90
4	CBD:[PROXYL-(POG) <sub>4</sub> POA(POG) <sub>5</sub> ] <sub>3</sub> (SEQ ID NO: 38)	3	43	25.48 ± 0.08	92
	CBD:[(POG) <sub>4</sub> POA(POG) <sub>5</sub> ] <sub>3</sub> (SEQ ID NO: 38)	3	43	124.45 ± 0.14	85
5	CBD:[PROXYL-(POG) <sub>3</sub> POA(POG) <sub>6</sub> ] <sub>3</sub> (SEQ ID NO: 39)	3	42	21.97 ± 0.14	94
6	CBD:[(POG) <sub>4</sub> POA(POG) <sub>5</sub> C-PROXYL] <sub>3</sub> (SEQ ID NO: 41)	3	44	24.09 ± 0.16	85
	CBD:[(POG) <sub>4</sub> POA(POG) <sub>5</sub> ] <sub>3</sub> (SEQ ID NO: 38)	3	42	24.67 ± 0.1	84
7	CBD:[11PROXYL-(POG) <sub>3</sub> PCG(POG) <sub>4</sub> ] <sub>3</sub> (SEQ ID NO: 40)	3	42		96
	CBD:[(POG) <sub>3</sub> PCG(POG) <sub>4</sub> ] <sub>3</sub> (SEQ ID NO: 40)	3	43	23.59 ± 0.05	90



[11PROXYL-(POG)<sub>3</sub>PCG(POG)<sub>4</sub>]<sub>3</sub> (SEQ ID NO: 40):CBD complex superimposes well with either NMR derived complexes i.e., CBD binding to the N-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) or to the C-terminal (POG)<sub>3</sub> (SEQ ID NO: 44) (FIGS. 9G and 9H).

Little Structural Change of <sup>15</sup>N-Minicollagen Upon CBD Binding: The studies thus far suggest that CBD scans the collagen fibril for under-twisted regions. Upon binding to the less structured regions, does it actively unwind collagen? Active unwinding by CBD would facilitate collagenolysis. To investigate two collagenous peptides selectively labeled with <sup>15</sup>N near N- or near C-terminus of [(POG)<sub>10</sub>]<sub>3</sub> (SEQ ID NO: 35) were synthesized (Table 4, peptides A, B), and the structural changes due to the binding of unlabeled CBD were monitored using <sup>1</sup>H-<sup>15</sup>N HSQC titration.

TABLE 4

<sup>15</sup> N-Labeled Mini-collagen	SEQ ID NO:
A POGPOG*POGPOGPOGPOGPOGPOGPOGPOG	35
B POGPOGPOGPOGPOGPOGPOGPOGPOG*POG	35
C POGPOG*POGPOGPOGPOGPOGPOGPOGPOG	38
D POGPOGPOGPOGPOGPOGPOGPOGPOG*POG	38

\*indicates the <sup>15</sup>N-labeled Glycine; A indicates Gly → Ala substitution.

The <sup>15</sup>N-Gly labeled peptides exhibited two distinct cross peaks in the <sup>1</sup>H-<sup>15</sup>N HSQC spectrum (FIGS. 10A and 10B). Those cross peaks corresponded to unwound monomer and triple helical conformations assigned in earlier NMR studies. Liu, et al. (1996) *Biochemistry* 35, 4306-4313 and Li, et al. (1993) *Biochemistry* 32, 7377-7387. The Gly residue closer to the terminal triplets exhibits both monomer and trimer peaks in the HSQC spectrum, whereas the Gly residue in the middle of the triple helix exhibits a strong trimer cross peak. If CBD is to bend or to cause any unwinding of the triple helix upon binding, we expected the cross peak corresponding to the triple helix to line broaden and disappear on the course of titration, and the cross peak corresponding to the single chain to intensify. However during the course of the titration, CBD did not instigate any changes on the <sup>1</sup>H-<sup>15</sup>N HSQC spectra of the collagenous peptides. Thus CBD bound to C-terminal (POG)<sub>3</sub> (SEQ ID NO: 35) imposed little structural changes to the triple helix.

Untwisted mini-collagen molecule selectively labeled with <sup>15</sup>N-Gly either at near the N- or C-termini (Table 4C and D) was titrated with unlabeled CBD. Cross peaks corresponding to monomer and triple-helix were identified on the HSQC spectra (FIGS. 10C and 10D). The titration of unlabeled CBD induced little change in the intensity of either monomer or trimer cross peak. Even upon binding to the partially unwound mini-collagen, CBD does not initiate any further unwinding.

CBD unidirectionally binds to the under-twisted site in the triple helical collagen. CBD may help disband the collagen fibril, but does not unwind the triple helix. Targeting under-twisted regions of tropocollagen may circumvent the energy barrier required for unwinding the triple helices. When CBD is used as a drug delivery molecule, the injected molecule distributes prominently to the end plates of vertebral discs, near the growth plates of tibia and fibula, and also to skin. It could be unloading its payload to the most blood accessible collagen that is undergoing remodeling, thus rich in under-twisted regions.

### Structural Comparison of ColH and ColG Collagen-Binding Domains

The C-terminal collagen-binding domain (CBD) of collagenase is required for insoluble collagen fibril binding and for subsequent collagenolysis. The high resolution crystal structures of ColG-CBD (s3b) and ColH-CBD (s3) the molecules resemble one another closely (r.m.s.d. C<sub>α</sub>=1.5 Å), despite sharing only 30% sequence identity. Five out of six residues chelating Ca<sup>2+</sup> are conserved. The dual Ca<sup>2+</sup> binding sites in s3 are completed by a functionally equivalent aspartate. The three most critical residues for collagen interaction in s3b are conserved in s3. The general shape of the binding pocket is retained by altered loop structures and side-chain positions. Small angle X-ray scattering data revealed that s3 also binds asymmetrically to mini-collagen. Besides the calcium-binding sites and the collagen-binding pocket, architecturally important hydrophobic residues and hydrogen-bonding network around the cis-peptide bond are well-conserved in metallopeptidase subfamily M9B.

Common structural features described above and in Bauer et al. (2012) *J Bacteriol* November 9 (which is incorporated herein by reference in its entirety), enabled us to update the sequence alignment of the CBD in the M9B subfamily (FIG. 1). Conserved residues are important for one of four reasons: calcium chelation (red), cis-trans isomerization of the linker (yellow), collagen-binding (blue) or protein folding (green). FIG. 1 also indicates the strands of the structure along the top of the figure.

The dual calcium-binding site is formed by four chelating residues (Glu899, Glu901, Asn903, and Asp904) within the N-terminal linker, two chelating residues (Asp927 and Asp930) from the β-strand C and invariant Tyr1002 hydrogen-bonds and orients Asp930. Residue numbers used in this paragraph are of s3b. Likewise other supporting cast such as Gly921 is conserved in the middle of β-strand strategically placed to make room for Glu899. The dual calcium chelation site is fashioned sometimes by functionally equivalent residues. As mentioned, Asp897 of s3 acts equivalently to Asp927 of s3b. Asp897 equivalents are tentatively identified in *B. brevis* s3a and s3b, *C. botulinum* A3 s3a and *C. histolyticum* ColG s3a. Tridentate and divalent Asp and Glu residues are conserved with only *C. sordellii* s3a as the exception. The monodentate Asp904 residue is sometimes substituted by Asn. For those substituted, the net charge of the dual calcium site is neutral rather than -1.

The peptide between residues 901-902 has cis conformation in the holo state for both s3b and s3. The position 902 in other CBD molecules is Pro, Asp or Asn. Pro frequently succeeds the peptide bond to ease trans-cis isomerization. The s3 molecule has Pro. In s3b, OD of Asn902 hydrogen-bonds with the main-chain N of Asp904. The hydrogen-bond is critical for the peptide isomerization. Spiriti and van der Vaart. (2010) *Biochemistry* 49:5314-5320, which is incorporated herein by reference in its entirety. For the remainder of CBD molecules with Asp at the position, OD of Asp could play the same role as that of Asn902. Other hydrogen-bonds identified by simulation studies important in stabilizing the transition states are well conserved. These donor-acceptor pairs in s3 and s3b are tabulated (Table 5). Calcium ions could catalyze the isomerization in all the CBD molecules and their transition states and catalytic mechanism may look very similar.

TABLE 5

Hydrogen-bonds important in trans-cis peptide isomerization in s3b and their counterparts in s3.	
Important H-bonds in s3b for transition state formation	Corresponding H-bonds in s3
T910_OG1 . . . N903_NH2	S879_OG1 . . . N872_ND2
T910_OG1 . . . N900_N	S879_OG1 . . . K86_N
E899_OE1 . . . N903_ND2	E868_OE1 . . . N872_ND2
E899_OE2 . . . S922_N	E868_OE2 . . . T891_N
N902_OD1 . . . D904_N	NA (N902 replaced with P871)
D930_OD2 . . . Y1002_OH	D939_OD2 . . . Y97_OH
Y1002_OH . . . Y932_OH	NA (Y932 replaced with F901)

Non-functional residues that are important in either folding or architectural stability are conserved. Hydrophobic residues packed between the  $\beta$ -sheets are better conserved if they are located in the vicinity of functionally critical residues. For example, invariant Trp956 of strand E is packed between the  $\beta$ -sheets. The residues flanking (Thr955 & Thr957) interact with mini-collagen. Tyr932 is packed between the sheets and helps positioning Tyr1002. Residues at tight turns are conserved as well. Gly975 is well conserved to allow a type II' turn in s3b. Gly942 (Gly975 equivalent) in s3 allows Asp941 side-chain to stabilize the reverse turn. A highly conserved six-residue stretch, between residues 986 and 991, adopts a tight turn and precedes the functionally important strand H. The region is well ordered in the crystal structures with low B-factors, and is the least dynamic based on NMR and limited proteolysis MALDI-TOF MS (25). Philominathan, et al. (2009) J Biol Chem 284:10868-10876 and Sides et al. J Am Soc Mass Spectrom. (2012) 23(3):505-19 both of which are incorporated herein by reference in their entireties. The main-chain carbonyl and amino groups of Arg985 hydrogen-bond with OH of Tyr989 to stabilize the turn. Only Gly987 can make room for the bulky Tyr989 side chain. Tyr990 packs against the invariant Ala909 and conserved  $3_{10}$  helix. Ala909 is at the base of the linker that undergoes  $\alpha$ -helix $\rightarrow\beta$ -strand transformation. The tight turn may ensure that collagen interacting Leu992, Tyr994, and Tyr996 would be correctly positioned. Tyr994 is the most critical residue in interacting with collagenous peptides. Wilson, et al. (2003) EMBO J 22:1743-1752. The strands adjacent to strand H, i.e. strands C and E, are very well conserved. The three antiparallel strands mold the collagen-binding pocket. Strand F staples the  $\beta$ -sheets by interacting with both sheets. The  $\beta$ -strand first interacts in an antiparallel orientation with strand E then breaks its direction at Gly971 to interact with strand G. In place of Gly971, Ala or Pro is found at the location where the strand switches its allegiance. The dual interaction of the strand helps positioning Tyr970 to interact with mini-collagen.

Three residues shown to interact strongly with mini-collagen are conserved. The invariant Tyr994 and well conserved Tyr970 and Tyr996 constitute the "hot spot". Y994A mutation lost binding capability. Since Y994F resulted in 12-fold reduction in binding to mini-collagen, the hydroxyl group of Tyr994 may interact with collagen through a hydrogen-bond. Tyr996, which is a critical residue in binding mini-collagen, is not so well conserved. Y996A caused 40-fold reduction in binding to the mini-collagen. Y996 in s3b is replaced with Phe in s3, though both side chains have identical orientation. In other CBD molecules, an aromatic residue, such as Phe or His, is sometimes found at the site. Y970A results in 12-fold reduction in binding to mini-collagen. Thr957 was found to interact with mini-collagen by  $^{15}\text{N}$ -HSQC-NMR titration. The  $\beta$ -branched

amino acid residues or Leu are found at the positions equivalent to Thr957 in most of the CBDs. Six other residues were identified by  $^{15}\text{N}$ -HSQC-NMR titration to interact with mini-collagen are not very well conserved. Since divergent CBDs (s3 and s3b) adopted a similar saddle-shaped binding pocket, other CBDs may also adopt similar collagen-binding strategy.

Divergent CBD could target different collagen sequences and could possibly target different collagen types; however, this structural study suggest otherwise. Rather, all the CBD domains may bind similarly to an under-twisted region such as the C-terminus of a collagen fibril. The C-terminus of type I collagen is exposed in the fibril surface based on X-ray fiber diffraction experiments, and it is the most accessible site for the bacterial collagenase to initiate assaults. However CBD binding only at the C-terminal region of tropo-collagen is unfounded. Gold particle-labeled tandem ColG-CBD (s3a-s3b) labeled with gold particle bound to type I collagen fibrils exhibited no periodicity. In the collagen fibrils, the molecules are staggered from each other by about 67 nm. Therefore CBD could target partially under-twisted regions in the middle of a tropocollagen that are also vulnerable for assaults.

Much like s3b, s3 is both compact, and extremely stable in the presence of physiological  $\text{Ca}^{2+}$ . Thus, the enzyme could degrade extracellular matrix for prolonged time. The linker that induced structural transformation is a common feature found in M9B collagenase. It could act as  $\text{Ca}^{2+}$  sensor to trigger domain rearrangement as means of enzyme activation.  $\text{Ca}^{2+}$  concentration in extracellular matrix is higher than that inside a bacterium. Both s3 and s3b bind similarly to a mini-collagen, thus M9B collagenase molecules could initiate collagenolysis from analogous structural features in various collagen fibril. Fusion protein of any CBD derived from M9B collagenase and a growth factor should result in comparable clinical outcome.

Example 3

CBD-PTH Agonist Spurs Hair Growth and  
CBD-PTH Antagonist Inhibits Hair Growth

In-Vitro Characterization of CBD-Linked PTH Compounds: Collagen binding of each peptide was verified in flow-through collagen binding assays as previously described in U.S. Patent Publication No. 2010/0129341, which is incorporated herein by reference in its entirety. PTH-CBD, consisting of the first 33 amino acids of PTH linked directly to the collagen binding domain (SEQ ID NO: 1), was the most potent agonist, having a similar effect to that of PTH(1-34) (SEQ ID NO: 7) on cAMP accumulation. Ponnappakkam et al. (2011) Calcif 88:511-520. Epub 2011 April 2022. Among the antagonists, PTH(7-33)-CBD (SEQ ID NO: 10) had the best combination of low intrinsic activity and high receptor blockade (not shown), similar to those seen in other PTH antagonists, including those used in hair growth studies. Peters, et al. (2001) J Invest Dermatol 117:173-178.

In-Vivo Distribution of PTH-CBD: Tissue distribution was assessed by administering  $^{35}\text{S}$ -labelled PTH-CBD via subcutaneous injection, followed by whole mount frozen and whole-body autoradiography. PTH-CBD with a phosphorylation site between PTH(1-33) and the CBD was purified, activated and labeled with  $[\gamma\text{-}^{35}\text{S}]$  ATP as described previously. Tamai et al. (2003) Infect Immun. 71:5371-5375. Approximately 10.8 mcg of  $^{35}\text{S}$ -PTH-CBD (122 kcm/mcg) was injected subcutaneously in 7 week-old

mice (32-35 g). Mice were sacrificed at 1 hour or 12 hours post-injection, and then frozen in dry ice-acetone. Frozen sections (50  $\mu$ m) were prepared with an autocryotome, dried at  $-20^{\circ}$  C., and exposed to an image plate for 4 weeks. There appeared to be an initial distribution of  $^{35}$ S-PTH-CBD to a broad area of skin around the site of injection, followed by a rapid redistribution to the skin of the entire animal, as well as to several other tissues (i.e. bone, intestine, bladder) (FIG. 11). PTH-CBD thus showed the desired properties of distribution and retention to skin with subcutaneous administration.

#### PTH-CBD Reverses Hair Loss in Chemotherapy-Induced Alopecia in Mice:

We compared efficacy of CBD linked PTH agonists and antagonists in chemotherapy-induced alopecia, utilizing an experimental design published by Peters, et al., for non-CBD linked PTH compounds. Peters, et al. (2001) *J Invest Dermatol* 117:173-178. C57BL/6J mice (Jackson Laboratories, Bar Harbor, Me.) were depilated to synchronize the hair follicles, and cyclophosphamide (CYP, 150 mg/kg) was administered on day 9 to maximize the chemotherapy-induced damage. The agonist (PTH-CBD) and the antagonist (PTH(7-33)-CBD) were administered 2 days prior to chemotherapy, and given the long-term retention of the compounds in the skin, we administered only a single dose to cover the timing of the multiple injections of PTH agonist and antagonist in the study by Peters, et al. The administered dose of CBD-linked compounds (320 mcg/kg) is well tolerated in mice. Ponnappakkam et al. (2011) *Calcif* 88:511-520. Epub 2011 April 2022.

The results of the photodocumentation record indicate that the agonist, PTH-CBD, was far more effective at stimulating hair growth than was the antagonist (FIG. 12). Histological examination revealed morphological changes in the hair follicles after CYP therapy, which were more superficially located and exhibited clumped melanocytes around the bulb, characteristics of the dystrophic anagen and catagen phase (FIG. 13). While the antagonist PTH(7-33)-CBD had no beneficial effect, treatment with the agonist PTH-CBD led to deeper rooting and reduced melanocyte clumping, thus reversing the dystrophic changes. Counts of anagen VI hair follicles per high-powered field (HPF) were compared between groups; animals treated with PTH-CBD had a higher number of hair follicles, approaching those of animals which did not receive chemotherapy (FIG. 14), while the antagonist PTH(7-33)-CBD had no beneficial effect.

Importantly, we saw no evidence of adverse effects from PTH-CBD administration. While PTH injections are known to elevate blood calcium and can cause kidney stones, PTH-CBD had no effect on serum calcium. In addition, there was no evidence of excess hair length on the body or of excess hair growth on the ears and tail, where a full coat is normally not present. The effects of PTH-CBD on hair growth have been confirmed in models of chemotherapy-induced alopecia without depilation, which more closely mimic clinical protocols.

**Quantification of Effects of PTH-CBD in Chemotherapy-Induced Alopecia:** We followed these studies by comparing the effects of different doses of PTH-CBD in chemotherapy-induced alopecia. In these studies, we applied the injections more distally on the back and applied a gray-scale analysis to quantify the amount of hair growth. Injecting more distally in the back allows us to compare regrowth of hair after PTH-CBD treatment with less interference from the normal hair regrowth, which normally proceeds from head

to tail in mice. The results are shown in FIG. 15, indicating a dose-dependent effect on hair regrowth both qualitatively and quantitatively.

**Chemotherapy-Induced Alopecia without Depilation:** While the depilated model of chemotherapy-induced alopecia provides a uniform model for comparison of drug effects, the depilation process is known to cause hair follicle injury, and may alter the response of the animals to the PTH-CBD administration. We therefore tested the effects of PTH-CBD in another model of chemotherapy-induced alopecia, where the animals were given 3 courses of cyclophosphamide therapy (50 mg/kg/wk), similar to the usual manner in which cancer patients might be treated. In this model, it takes much longer (4-6 months) for alopecia to develop. Animals that received a single dose of PTH-CBD (320 mcg/kg subcutaneous) prior to the first cycle did not develop hair loss as shown in FIG. 16.

In a second study, we compared the effects of PTH-CBD when given prophylactically, at the time of the first cycle of chemotherapy, vs. therapeutically, after the hair loss had developed. While PTH-CBD was effective in both instances, the effects were more prominent when given prophylactically. This is evident both visually and quantitatively in FIG. 17, using the same grey scale analysis used in our dose-response study.

**Depilation Alopecia:** The agonist PTH-CBD appears to increase hair growth by increasing the number of anagen phase hair follicles. As such, there is no reason to believe that hair growth effects should be limited to the chemotherapy model. We therefore tested both PTH-CBD and antagonist compound, PTH(7-33)-CBD, after removing hair from C57/BL6J mice by waxing (FIG. 18). The results were quite interesting; agonist (PTH-CBD) treated animals had earlier anagen eruption (day 7 vs. day 9 for vehicle controls), and exhibited more complete regrowth of hair by the end of the study (day 18). Antagonist (PTH(7-33)-CBD) treated animals also had an early anagen eruption, but the hair growth which followed was markedly curtailed, and the hair cycle was arrested after this point, resulting no further observed regrowth of hair. Thus, it appears that agonist therapy is acting to promote more rapid regrowth of hair by promoting more rapid transition to the anagen phase, while the antagonist inhibited hair regrowth by blocking this transition.

PTH-CBD is a fusion protein of the first 33 amino acids of parathyroid hormone (PTH) and a bacterial collagen binding domain. The collagen binding activity causes PTH-CBD to be retained at its site of action in the dermal collagen, maximizing efficacy and reducing systemic side-effects. PTH-CBD stimulates hair growth by causing hair follicles to enter an anagen VI or growth phase, presumably by activating WNT signaling and increasing production of beta-catenin. We therefore plan to conduct the following additional studies to confirm this mechanism of action and to determine the effect of PTH-CBD in two distinct genetic mouse models with WNT signaling inhibition. These data will be used in formulating clinical trials for PTH-CBD as a therapy for alopecia.

**Alopecia Areata:** Alopecia Areata is a disease of patchy hair loss due to autoimmune destruction of the hair follicles. We tested the efficacy of PTH-CBD in promoting regrowth of hair in an animal model of alopecia areata, the engrafted C3H/HeJ mouse. In this model, hair loss develops variably over the first 2 months of life. Shown in FIG. 19 is the results of a single dose of PTH-CBD (320 mcg/kg subcutaneous) administered into the engrafted site, the center of the back, where there was maximal hair loss. Compared to vehicle

## 29

control animals, which continued to lose hair at this site, animals receiving PTH-CBD began to show regrowth of hair within the next 1-4 days. Importantly, the response was found to be sustained during the 2 month course of the experiment.

## Example 4

CBD-PTH can Prevent or Treat  
Hyperparathyroidism

In this experiment, rats had their ovaries surgically removed at age 3 months. At age 9 months, rats were injected with either a single dose of PTH-CBD (320 mcg/kg)

## 30

or vehicle control. Animals were sacrificed 6 months after therapy (age 15 months). Human intact PTH levels were measured to assess serum levels of PTH-CBD, and were found to be undetectable in both groups. Serum calcium was measured and there were no differences between groups (Vehicle: 13.5+/-1.1 vs. PTH-CBD: 14.3+/-1.1 mg/dl, NS). Rat intact PTH levels were measured to assess endogenous PTH production, and PTH-CBD suppressed the normal increase in endogenous PTH levels seen in aged, ovariectomized rats. These findings indicate that a single injection of PTH-CBD can provide long-term suppression of endogenous PTH production, preventing the normal rise seen with age in the ovariectomized rat model, and thus may serve as a therapy for hyperparathyroidism.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 47

<210> SEQ ID NO 1

<211> LENGTH: 158

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion protein containing parathyroid hormone segment and collagen-binding domain

<400> SEQUENCE: 1

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn  
1 5 10 15

Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His  
20 25 30

Asn Gly Ile Asn Ser Pro Val Tyr Pro Ile Gly Thr Glu Lys Glu Pro  
35 40 45

Asn Asn Ser Lys Glu Thr Ala Ser Gly Pro Ile Val Pro Gly Ile Pro  
50 55 60

Val Ser Gly Thr Ile Glu Asn Thr Ser Asp Gln Asp Tyr Phe Tyr Phe  
65 70 75 80

Asp Val Ile Thr Pro Gly Glu Val Lys Ile Asp Ile Asn Lys Leu Gly  
85 90 95

Tyr Gly Gly Ala Thr Trp Val Val Tyr Asp Glu Asn Asn Asn Ala Val  
100 105 110

Ser Tyr Ala Thr Asp Asp Gly Gln Asn Leu Ser Gly Lys Phe Lys Ala  
115 120 125

Asp Lys Pro Gly Arg Tyr Tyr Ile His Leu Tyr Met Phe Asn Gly Ser  
130 135 140

Tyr Met Pro Tyr Arg Ile Asn Ile Glu Gly Ser Val Gly Arg  
145 150 155

<210> SEQ ID NO 2

<211> LENGTH: 251

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion protein containing parathyroid hormone fragment and collagen-binding domain and polycystic kidney disease domain of ColH.

<400> SEQUENCE: 2

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn  
1 5 10 15

Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His  
20 25 30

-continued

---

Asn	Gly	Ile	Pro	Glu	Ile	Lys	Asp	Leu	Ser	Glu	Asn	Lys	Leu	Pro	Val
	35						40				45				
Ile	Tyr	Met	His	Val	Pro	Lys	Ser	Gly	Ala	Leu	Asn	Gln	Lys	Val	Val
	50					55					60				
Phe	Tyr	Gly	Lys	Gly	Thr	Tyr	Asp	Pro	Asp	Gly	Ser	Ile	Ala	Gly	Tyr
	65				70					75				80	
Gln	Trp	Asp	Phe	Gly	Asp	Gly	Ser	Asp	Phe	Ser	Ser	Glu	Gln	Asn	Pro
			85						90					95	
Ser	His	Val	Tyr	Thr	Lys	Lys	Gly	Glu	Tyr	Thr	Val	Thr	Leu	Arg	Val
			100					105					110		
Met	Asp	Ser	Ser	Gly	Gln	Met	Ser	Glu	Lys	Thr	Met	Lys	Ile	Lys	Ile
		115					120					125			
Thr	Asp	Pro	Val	Tyr	Pro	Ile	Gly	Thr	Glu	Lys	Glu	Pro	Asn	Asn	Ser
	130					135					140				
Lys	Glu	Thr	Ala	Ser	Gly	Pro	Ile	Val	Pro	Gly	Ile	Pro	Val	Ser	Gly
	145				150					155					160
Thr	Ile	Glu	Asn	Thr	Ser	Asp	Gln	Asp	Tyr	Phe	Tyr	Phe	Asp	Val	Ile
			165					170						175	
Thr	Pro	Gly	Glu	Val	Lys	Ile	Asp	Ile	Asn	Lys	Leu	Gly	Tyr	Gly	Gly
			180				185						190		
Ala	Thr	Trp	Val	Val	Tyr	Asp	Glu	Asn	Asn	Asn	Ala	Val	Ser	Tyr	Ala
		195					200					205			
Thr	Asp	Asp	Gly	Gln	Asn	Leu	Ser	Gly	Lys	Phe	Lys	Ala	Asp	Lys	Pro
	210					215					220				
Gly	Arg	Tyr	Tyr	Ile	His	Leu	Tyr	Met	Phe	Asn	Gly	Ser	Tyr	Met	Pro
	225				230					235					240
Tyr	Arg	Ile	Asn	Ile	Glu	Gly	Ser	Val	Gly	Arg					
			245						250						

<210> SEQ ID NO 3  
 <211> LENGTH: 5464  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic expression vector

<400> SEQUENCE: 3

agcttatcga ctgcacgggtg caccaatgct tctggcgctca ggcagccatc ggaagctgtg	60
gtatggctgt gcaggctcgt aatcactgca taattcgtgt cgctcaaggc gcaactccgt	120
tctggataat gttttttgctg ccgacatcat aacggttctgt gcaaatattc tgaaatgagc	180
tggtgacaat taatcatcgg ctgcgtataat gtgtggaatt gtgagcggat aacaatttca	240
cacaggaaac agtattcatg tccctatac taggttattg gaaaattaag ggccttggtc	300
aaccactcgt acttcttttg gaatatcttg aagaaaaata tgaagagcat ttgtatgagc	360
gcgatgaagg tgataaatgg cgaacaaaa agtttgaatt gggtttggag tttcccaatc	420
ttccttatta tattgatggt gatgttaaat taacacagtc tatggccatc atacgttata	480
tagctgacaa gcacaacatg ttgggtggtt gtccaaaaga gcgtgcagag atttcaatgc	540
ttgaaggagc gggttttgat attagatacg gtgtttcgag aattgcatat agtaaagact	600
ttgaaactct caaagttgat tttcttagca agctacctga aatgctgaaa atgttcgaag	660
atcgtttatg tcataaaaca tatttaaatg gtgatcatgt aaccatcct gacttcatgt	720
tgtatgacgc tcttgatggt gttttatata tggacccaat gtgcctggat gcgttcccaa	780
aattagtttg ttttaaaaaa cgtattgaag ctatcccaca aattgataag tacttgaaat	840

-continued

---

ccagcaagta tatagcatgg cctttgcagg gctggcaagc cacgtttggt ggtggcgacc	900
atcctccaaa atcggatctg atcgaaggtc gttctgtgag tgaataacag cttatgcata	960
acctgggaaa acatctgaac tcgatggaga gagtagaatg gctgcgtaag aagctgcagg	1020
atgtgcacaa tggaattaat tccccggtat atccaatagg cactgaaaaa gaaccaaata	1080
acagtaaaga aactgcaagt ggtccaatag taccagggtat acctgttagt ggaaccatag	1140
aaaatacaag tgatcaagat tatttctatt ttgatgttat aacaccagga gaagtaaaaa	1200
tagatataaa taaattaggg tacggaggag ctacttgggt agtatatgat gaaaataata	1260
atgcagtatc ttatgccact gatgatgggc aaaatttaag tggaagttt aaggcagata	1320
aaccaggtag atattacatc catctttaca tgtttaatgg tagttatatg ccatatagaa	1380
ttaatataga aggttcagta ggaagataat attttattag ttgaggtaac tccactcgaa	1440
ttggtcgact cgagcggcgc catcgtgact gactgacgat ctgcctcgcg cgtttcgggtg	1500
atgacggtag aaacctctga cacatgcagc tcccggagac ggtcacagct tgtctgtaag	1560
cggatgccgg gagcagacaa gcccgtcagg gcgcgtcagc ggggtgtggc ggggtgtcggg	1620
gcgcagccat gaccagtcga cgtagcgata gcggagtgtg taattcttga agacgaaagg	1680
gcctcgtgat acgcctatct ttatagggtt atgtcatgat aataatgggt tcttagacgt	1740
cagggtggcac ttttcgggga aatgtgcgcg gaacccttat ttgtttatct ttctaaatac	1800
attcaaatat gtatccgctc atgagacaat aacctgata aatgcttcaa taatattgaa	1860
aaaggaagag tatgagtatt caacatttcc gtgtcgccct tattcccttt tttgcggcat	1920
tttgcccttc tgtttttgct caccagaaa cgctggtgaa agtaaaagat gctgaagatc	1980
agttgggtgc acgagtgggt tacatcgaa cggatctcaa cagcggtaag atccttgaga	2040
gttttcgccc cgaagaacgt tttccaatga tgagcacttt taaagttctg ctatgtggcg	2100
cggattatc cctgtttgac gccgggcaag agcaactcgg tcgccgcata cactattctc	2160
agaatgactt ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcatgacag	2220
taagagaatt atgcagtgt gccataacca tgagtataa cactgcgggc aacttacttc	2280
tgacaacgat cggaggaccg aaggagctaa ccgcttttt gcacaacatg ggggatcatg	2340
taactcgct tgatcgttg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg	2400
acaccacgat gcctgcagca atggcaacaa cgttcgcaa actattaact ggcgaactac	2460
ttactctagc tccccgcaa caattaatag actggatgga ggcggataaa gttgcaggac	2520
cacttctcgc ctcggccctt ccggctgggt ggtttattgc tgataaatct ggagccgggtg	2580
agcgtgggtc tcgcggtatc attgcagcac tggggccaga tggtaaagccc tcccgtatcg	2640
tagttatcta cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg	2700
agatagggtc ctcactgatt aagcattggt aactgtcaga ccaagtttac tcatatatac	2760
tttagattga tttaaaactt catttttaat ttaaaaggat ctagggtgaag atcctttttg	2820
ataatctcat gacaaaaac ccttaacgtg agttttcggt ccaactgagcg tcagaccccg	2880
tagaaaagat caaaggatct tcttgagatc ctttttttct gcgcgtaatc tgctgcttgc	2940
aaacaaaaaa accaccgcta ccagcgggtg tttgtttgcc ggatcaagag ctaccaactc	3000
tttttcgcaa ggtaactggc ttcagcagag cgcagatacc aaatactgtc cttctagtgt	3060
agccgtagtt aggccaccac ttaagaact ctgtagcacc gcctacatac ctcgctctgc	3120
taatcctgtt accagtgggt gctgccagtg gcgataagtc gtgtcttacc gggttggact	3180
caagacgata gttaccgat aaggcgcagc ggtcgggctg aacggggggt tcgtgcacac	3240

-continued

---

agcccagctt ggagcgaaacg acctacaccg aactgagata cctacagcgt gagctatgag	3300
aaagcgccac gcttcccga gggagaaaag cggacaggta tccggaagc ggcagggctc	3360
gaacaggaga gcgcacgagg gagcttccag ggggaaacgc ctggtatctt tatagtctcg	3420
tggggtttcg ccacctctga cttgagcgtc gattttttgtg atgctcgtca ggggggcgga	3480
gcctatggaa aaacgccagc aacgcggcct ttttacggtt cctggccttt tgctggcctt	3540
ttgctcacat gttctttctc gcgttatccc ctgattctgt ggataaccgt attaccgcct	3600
ttgagtgagc tgataccgct cgcgcagcc gaacgaccga gcgcagcgag tcagtgagcg	3660
aggaagcgga agagcgctg atgcggtatt ttctccttac gcctctgtgc ggtatttcac	3720
accgcataaa ttccgacacc atcgaatggt gcaaaacctt tcgcggtatg gcctgatagc	3780
gcccgaaga gagtcaattc aggggtggtga atgtgaaacc agtaacgtta tacgatgtcg	3840
cagagtatgc cgggtgtctt tatcagaccg ttcccgcggt ggtgaaccag gccagccacg	3900
tttctgcgaa aacgcgggaa aaagtggag cggcgatggc ggagctgaat tacattccca	3960
accgcgtggc acaacaactg gcgggcaaac agtcgttgct gattggcggt gccacctcca	4020
gtctggccct gcacgcgcg tcgcaaatg tcgcggcgat taaatctcgc gccgatcaac	4080
tgggtgccag cgtggtggtg tcgatggtag aacgaagcgg cgtcgaagcc tgtaaagcgg	4140
cgggtgcacaa tcttctcgcg caacgcgtc gtgggctgat cattaactat ccgctggatg	4200
accaggatgc cattgtgtgt gaagctgcct gcaactaatg tccggcgta tttcttgatg	4260
tctctgacca gacaccctc aacagtatta ttttctccca tgaagacggt acgcgactgg	4320
gcgtggagca tctggtcgca ttgggtcacc agcaaatcgc gctgttagcg ggcccatata	4380
gttctgtctc ggcgcgctc cgtctggctg gctggcataa atatctcact cgcaatcaaa	4440
ttcagccgat agcggaaacg gaaggcgact ggagtgccat gtccggtttt caacaaacca	4500
tgcaaatgct gaatgagggc atcggtccca ctgcgatgct ggttgccaac gatcagatgg	4560
cgtctggcgc aatgcgcgc attaccgagt ccgggctcgc cgttggtcgc gatattctcg	4620
tagtgggata cgacgatac gaagacagct catgttatat cccgcggtta accaccatca	4680
aacaggattt tcgcctgctg gggcaaaacca gcgtggaccg cttgctgcaa ctctctcagg	4740
gccaggcggg gaagggaat cagctgttgc ccgtctcact ggtgaaaaga aaaaccaccc	4800
tggcgcccaa tacgaaaacc gcctctcccc gcgcgttggc cgattcatta atgcagctgg	4860
cacgacaggt ttcccgactg gaaagcgggc agtgagcga acgcaattaa tgtgagttag	4920
ctcactcatt aggcacccca ggttttacac tttatgttc cggctcgtat gttgtgtgga	4980
attgtgagcg gataacaatt tcacacagga aacagctatg accatgatta cggattcact	5040
ggcgtctggt ttacaacgct gtgactggga aaacctggc gttaccaac ttaatcgctt	5100
tgcagcacat cccctttctg ccagctggcg taatagcgaa gaggccgca ccgatcgccc	5160
ttcccaacag ttgcgcagcc tgaatggcga atggcgcttt gcctggtttc cggcaccaga	5220
agcgtgctcg gaaagctggc tggagtgcga tcttctctgag gccgatactg tcgtcgtccc	5280
ctcaaacctg cagatgcacg gttacgatgc gccatctac accaacgtaa cctatcccat	5340
tacggtcaat ccgcgcttg ttccacagga gaatccgacg ggtgtgtact cgctcacatt	5400
taatgttgat gaaagctggc tacaggaagg ccagacgcga attatttttg atggcgttgg	5460
aatt	5464

-continued

---

```

<210> SEQ ID NO 4
<211> LENGTH: 383
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic GST-PTH-CBD fusion protein

<400> SEQUENCE: 4

Met Ser Pro Ile Leu Gly Tyr Trp Lys Ile Lys Gly Leu Val Gln Pro
1          5          10          15

Thr Arg Leu Leu Leu Glu Tyr Leu Glu Glu Lys Tyr Glu Glu His Leu
20          25          30

Tyr Glu Arg Asp Glu Gly Asp Lys Trp Arg Asn Lys Lys Phe Glu Leu
35          40          45

Gly Leu Glu Phe Pro Asn Leu Pro Tyr Tyr Ile Asp Gly Asp Val Lys
50          55          60

Leu Thr Gln Ser Met Ala Ile Ile Arg Tyr Ile Ala Asp Lys His Asn
65          70          75          80

Met Leu Gly Gly Cys Pro Lys Glu Arg Ala Glu Ile Ser Met Leu Glu
85          90          95

Gly Ala Val Leu Asp Ile Arg Tyr Gly Val Ser Arg Ile Ala Tyr Ser
100         105         110

Lys Asp Phe Glu Thr Leu Lys Val Asp Phe Leu Ser Lys Leu Pro Glu
115         120         125

Met Leu Lys Met Phe Glu Asp Arg Leu Cys His Lys Thr Tyr Leu Asn
130         135         140

Gly Asp His Val Thr His Pro Asp Phe Met Leu Tyr Asp Ala Leu Asp
145         150         155         160

Val Val Leu Tyr Met Asp Pro Met Cys Leu Asp Ala Phe Pro Lys Leu
165         170         175

Val Cys Phe Lys Lys Arg Ile Glu Ala Ile Pro Gln Ile Asp Lys Tyr
180         185         190

Leu Lys Ser Ser Lys Tyr Ile Ala Trp Pro Leu Gln Gly Trp Gln Ala
195         200         205

Thr Phe Gly Gly Gly Asp His Pro Pro Lys Ser Asp Leu Ile Glu Gly
210         215         220

Arg Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu
225         230         235         240

Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val
245         250         255

His Asn Gly Ile Asn Ser Pro Val Tyr Pro Ile Gly Thr Glu Lys Glu
260         265         270

Pro Asn Asn Ser Lys Glu Thr Ala Ser Gly Pro Ile Val Pro Gly Ile
275         280         285

Pro Val Ser Gly Thr Ile Glu Asn Thr Ser Asp Gln Asp Tyr Phe Tyr
290         295         300

Phe Asp Val Ile Thr Pro Gly Glu Val Lys Ile Asp Ile Asn Lys Leu
305         310         315         320

Gly Tyr Gly Gly Ala Thr Trp Val Val Tyr Asp Glu Asn Asn Asn Ala
325         330         335

Val Ser Tyr Ala Thr Asp Asp Gly Gln Asn Leu Ser Gly Lys Phe Lys
340         345         350

```



-continued

---

Ala Asp Lys Pro Gly Arg Tyr Tyr Ile His Leu Tyr Met Phe Asn Gly  
           355                                  360                                  365

Ser Tyr Met Pro Tyr Arg Ile Asn Ile Glu Gly Ser Val Gly Arg  
       370                                  375                                  380

<210> SEQ ID NO 5  
 <211> LENGTH: 4  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic peptide: Factor Xa recognition  
                                   sequence

<400> SEQUENCE: 5

Ile Glu Gly Arg  
 1

<210> SEQ ID NO 6  
 <211> LENGTH: 1021  
 <212> TYPE: PRT  
 <213> ORGANISM: Clostridium histolyticum

<400> SEQUENCE: 6

Met Lys Arg Lys Cys Leu Ser Lys Arg Leu Met Leu Ala Ile Thr Met  
 1                  5                                  10                                  15

Ala Thr Ile Phe Thr Val Asn Ser Thr Leu Pro Ile Tyr Ala Ala Val  
                   20                                  25                                  30

Asp Lys Asn Asn Ala Thr Ala Ala Val Gln Asn Glu Ser Lys Arg Tyr  
                   35                                  40                                  45

Thr Val Ser Tyr Leu Lys Thr Leu Asn Tyr Tyr Asp Leu Val Asp Leu  
                   50                                  55                                  60

Leu Val Lys Thr Glu Ile Glu Asn Leu Pro Asp Leu Phe Gln Tyr Ser  
                   65                                  70                                  75                                  80

Ser Asp Ala Lys Glu Phe Tyr Gly Asn Lys Thr Arg Met Ser Phe Ile  
                   85                                  90                                  95

Met Asp Glu Ile Gly Arg Arg Ala Pro Gln Tyr Thr Glu Ile Asp His  
                   100                                  105                                  110

Lys Gly Ile Pro Thr Leu Val Glu Val Val Arg Ala Gly Phe Tyr Leu  
                   115                                  120                                  125

Gly Phe His Asn Lys Glu Leu Asn Glu Ile Asn Lys Arg Ser Phe Lys  
                   130                                  135                                  140

Glu Arg Val Ile Pro Ser Ile Leu Ala Ile Gln Lys Asn Pro Asn Phe  
                   145                                  150                                  155                                  160

Lys Leu Gly Thr Glu Val Gln Asp Lys Ile Val Ser Ala Thr Gly Leu  
                   165                                  170                                  175

Leu Ala Gly Asn Glu Thr Ala Pro Pro Glu Val Val Asn Asn Phe Thr  
                   180                                  185                                  190

Pro Ile Leu Gln Asp Cys Ile Lys Asn Ile Asp Arg Tyr Ala Leu Asp  
                   195                                  200                                  205

Asp Leu Lys Ser Lys Ala Leu Phe Asn Val Leu Ala Ala Pro Thr Tyr  
                   210                                  215                                  220

Asp Ile Thr Glu Tyr Leu Arg Ala Thr Lys Glu Lys Pro Glu Asn Thr  
                   225                                  230                                  235                                  240

Pro Trp Tyr Gly Lys Ile Asp Gly Phe Ile Asn Glu Leu Lys Lys Leu  
                   245                                  250                                  255

Ala Leu Tyr Gly Lys Ile Asn Asp Asn Asn Ser Trp Ile Ile Asp Asn  
                   260                                  265                                  270

Gly 275	Ile 280	Tyr 285	His 290	Ile 295	Ala 300	Pro 305	Leu 310	Gly 315	Lys 320	Leu 325	His 330	Ser 335	Asn 340	Asn 345	Lys 350
Ile 290	Gly 295	Ile 300	Glu 305	Thr 310	Leu 315	Thr 320	Glu 325	Val 330	Met 335	Lys 340	Val 345	Tyr 350	Pro 355	Tyr 360	Leu 365
Ser 305	Met 310	Gln 315	His 320	Leu 325	Gln 330	Ser 335	Ala 340	Asp 345	Gln 350	Ile 355	Lys 360	Arg 365	His 370	Tyr 375	Asp 380
Ser 315	Lys 320	Asp 325	Ala 330	Glu 335	Gly 340	Asn 345	Lys 350	Ile 355	Pro 360	Leu 365	Asp 370	Lys 375	Phe 380	Lys 385	Lys 390
Glu 325	Gly 330	Lys 335	Glu 340	Lys 345	Tyr 350	Cys 355	Pro 360	Lys 365	Thr 370	Tyr 375	Thr 380	Phe 385	Asp 390	Asp 395	Gly 400
Lys 335	Val 340	Ile 345	Ile 350	Lys 355	Ala 360	Gly 365	Ala 370	Arg 375	Val 380	Glu 385	Glu 390	Glu 395	Lys 400	Val 405	Lys 410
Arg 345	Leu 350	Tyr 355	Trp 360	Ala 365	Ser 370	Lys 375	Glu 380	Val 385	Asn 390	Ser 395	Gln 400	Phe 405	Phe 410	Arg 415	Val 420
Tyr 355	Gly 360	Ile 365	Asp 370	Lys 375	Pro 380	Leu 385	Glu 390	Glu 395	Gly 400	Asn 405	Pro 410	Asp 415	Asp 420	Ile 425	Leu 430
Thr 365	Met 370	Val 375	Ile 380	Tyr 385	Asn 390	Ser 395	Pro 400	Glu 405	Glu 410	Tyr 415	Lys 420	Leu 425	Asn 430	Ser 435	Val 440
Leu 375	Tyr 380	Gly 385	Tyr 390	Asp 395	Thr 400	Asn 405	Asn 410	Gly 415	Gly 420	Met 425	Tyr 430	Ile 435	Glu 440	Pro 445	Glu 450
Gly 385	Thr 390	Phe 395	Phe 400	Thr 405	Tyr 410	Glu 415	Arg 420	Glu 425	Ala 430	Gln 435	Glu 440	Ser 445	Thr 450	Tyr 455	Thr 460
Leu 395	Glu 400	Glu 405	Leu 410	Phe 415	Arg 420	His 425	Glu 430	Tyr 435	Thr 440	His 445	Tyr 450	Leu 455	Gln 460	Gly 465	Arg 470
Tyr 405	Ala 410	Val 415	Pro 420	Gly 425	Gln 430	Trp 435	Gly 440	Arg 445	Thr 450	Lys 455	Leu 460	Tyr 465	Asp 470	Asn 475	Asp 480
Arg 415	Leu 420	Thr 425	Trp 430	Tyr 435	Glu 440	Glu 445	Gly 450	Gly 455	Ala 460	Glu 465	Leu 470	Phe 475	Ala 480	Gly 485	Ser 490
Thr 425	Arg 430	Thr 435	Ser 440	Gly 445	Ile 450	Leu 455	Pro 460	Arg 465	Lys 470	Ser 475	Ile 480	Val 485	Ser 490	Asn 495	Ile 500
His 435	Asn 440	Thr 445	Thr 450	Arg 455	Asn 460	Asn 465	Arg 470	Tyr 475	Lys 480	Leu 485	Ser 490	Asp 495	Thr 500	Val 505	His 510
Ser 445	Lys 450	Tyr 455	Gly 460	Ala 465	Ser 470	Phe 475	Glu 480	Phe 485	Tyr 490	Asn 495	Tyr 500	Ala 505	Cys 510	Met 515	Phe 520
Met 455	Asp 460	Tyr 465	Met 470	Tyr 475	Asn 480	Lys 485	Asp 490	Met 495	Gly 500	Ile 505	Leu 510	Asn 515	Lys 520	Leu 525	Asn 530
Asp 465	Leu 470	Ala 475	Lys 480	Asn 485	Asn 490	Asp 495	Val 500	Asp 505	Gly 510	Tyr 515	Asp 520	Asn 525	Tyr 530	Ile 535	Arg 540
Asp 475	Leu 480	Ser 485	Ser 490	Asn 495	Tyr 500	Ala 505	Leu 510	Asn 515	Asp 520	Lys 525	Tyr 530	Gln 535	Asp 540	His 545	Met 550
Gln 485	Glu 490	Arg 495	Ile 500	Asp 505	Asn 510	Tyr 515	Glu 520	Asn 525	Leu 530	Thr 535	Val 540	Pro 545	Phe 550	Val 555	Ala 560
Asp 495	Asp 500	Tyr 505	Leu 510	Val 515	Arg 520	His 525	Ala 530	Tyr 535	Lys 540	Asn 545	Pro 550	Asn 555	Glu 560	Ile 565	Tyr 570
Ser 505	Glu 510	Ile 515	Ser 520	Glu 525	Val 530	Ala 535	Lys 540	Leu 545	Lys 550	Asp 555	Ala 560	Lys 565	Ser 570	Glu 575	Val 580
Lys 515	Lys 520	Ser 525	Gln 530	Tyr 535	Phe 540	Ser 545	Thr 550	Phe 555	Thr 560	Leu 565	Arg 570	Gly 575	Ser 580	Tyr 585	Thr 590
Gly 525	Gly 530	Ala 535	Ser 540	Lys 545	Gly										

-continued

---

Tyr Lys Thr Leu Thr Ala Tyr Phe Thr Asn Tyr Lys Val Asp Ser Ser  
 690 695 700  
 Asn Arg Val Thr Tyr Asp Val Val Phe His Gly Tyr Leu Pro Asn Glu  
 705 710 715 720  
 Gly Asp Ser Lys Asn Ser Leu Pro Tyr Gly Lys Ile Asn Gly Thr Tyr  
 725 730 735  
 Lys Gly Thr Glu Lys Glu Lys Ile Lys Phe Ser Ser Glu Gly Ser Phe  
 740 745 750  
 Asp Pro Asp Gly Lys Ile Val Ser Tyr Glu Trp Asp Phe Gly Asp Gly  
 755 760 765  
 Asn Lys Ser Asn Glu Glu Asn Pro Glu His Ser Tyr Asp Lys Val Gly  
 770 775 780  
 Thr Tyr Thr Val Lys Leu Lys Val Thr Asp Asp Lys Gly Glu Ser Ser  
 785 790 795 800  
 Val Ser Thr Thr Thr Ala Glu Ile Lys Asp Leu Ser Glu Asn Lys Leu  
 805 810 815  
 Pro Val Ile Tyr Met His Val Pro Lys Ser Gly Ala Leu Asn Gln Lys  
 820 825 830  
 Val Val Phe Tyr Gly Lys Gly Thr Tyr Asp Pro Asp Gly Ser Ile Ala  
 835 840 845  
 Gly Tyr Gln Trp Asp Phe Gly Asp Gly Ser Asp Phe Ser Ser Glu Gln  
 850 855 860  
 Asn Pro Ser His Val Tyr Thr Lys Lys Gly Glu Tyr Thr Val Thr Leu  
 865 870 875 880  
 Arg Val Met Asp Ser Ser Gly Gln Met Ser Glu Lys Thr Met Lys Ile  
 885 890 895  
 Lys Ile Thr Asp Pro Val Tyr Pro Ile Gly Thr Glu Lys Glu Pro Asn  
 900 905 910  
 Asn Ser Lys Glu Thr Ala Ser Gly Pro Ile Val Pro Gly Ile Pro Val  
 915 920 925  
 Ser Gly Thr Ile Glu Asn Thr Ser Asp Gln Asp Tyr Phe Tyr Phe Asp  
 930 935 940  
 Val Ile Thr Pro Gly Glu Val Lys Ile Asp Ile Asn Lys Leu Gly Tyr  
 945 950 955 960  
 Gly Gly Ala Thr Trp Val Val Tyr Asp Glu Asn Asn Asn Ala Val Ser  
 965 970 975  
 Tyr Ala Thr Asp Asp Gly Gln Asn Leu Ser Gly Lys Phe Lys Ala Asp  
 980 985 990  
 Lys Pro Gly Arg Tyr Tyr Ile His Leu Tyr Met Phe Asn Gly Ser Tyr  
 995 1000 1005  
 Met Pro Tyr Arg Ile Asn Ile Glu Gly Ser Val Gly Arg  
 1010 1015 1020  
  
 <210> SEQ ID NO 7  
 <211> LENGTH: 84  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 7  
  
 Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn  
 1 5 10 15  
 Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His  
 20 25 30  
 Asn Phe Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp Ala Gly Ser  
 35 40 45

-continued

---

Gln Arg Pro Arg Lys Lys Glu Asp Asn Val Leu Val Glu Ser His Glu  
50 55 60

Lys Ser Leu Gly Glu Ala Asp Lys Ala Asp Val Asn Val Leu Thr Lys  
65 70 75 80

Ala Lys Ser Gln

<210> SEQ ID NO 8

<211> LENGTH: 141

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile Gln  
1 5 10 15

Asp Leu Arg Arg Arg Phe Phe Leu His His Leu Ile Ala Glu Ile His  
20 25 30

Thr Ala Glu Ile Arg Ala Thr Ser Glu Val Ser Pro Asn Ser Lys Pro  
35 40 45

Ser Pro Asn Thr Lys Asn His Pro Val Arg Phe Gly Ser Asp Asp Glu  
50 55 60

Gly Arg Tyr Leu Thr Gln Glu Thr Asn Lys Val Glu Thr Tyr Lys Glu  
65 70 75 80

Gln Pro Leu Lys Thr Pro Gly Lys Lys Lys Lys Gly Lys Pro Gly Lys  
85 90 95

Arg Lys Glu Gln Glu Lys Lys Lys Arg Arg Thr Arg Ser Ala Trp Leu  
100 105 110

Asp Ser Gly Val Thr Gly Ser Gly Leu Glu Gly Asp His Leu Ser Asp  
115 120 125

Thr Ser Thr Thr Ser Leu Glu Leu Asp Ser Arg Arg His  
130 135 140

<210> SEQ ID NO 9

<211> LENGTH: 160

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Gly-Ser-PTH(1-33)-CBD fusion protein

<400> SEQUENCE: 9

Gly Ser Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His  
1 5 10 15

Leu Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp  
20 25 30

Val His Asn Gly Ile Asn Ser Pro Val Tyr Pro Ile Gly Thr Glu Lys  
35 40 45

Glu Pro Asn Asn Ser Lys Glu Thr Ala Ser Gly Pro Ile Val Pro Gly  
50 55 60

Ile Pro Val Ser Gly Thr Ile Glu Asn Thr Ser Asp Gln Asp Tyr Phe  
65 70 75 80

Tyr Phe Asp Val Ile Thr Pro Gly Glu Val Lys Ile Asp Ile Asn Lys  
85 90 95

Leu Gly Tyr Gly Gly Ala Thr Trp Val Val Tyr Asp Glu Asn Asn Asn  
100 105 110

Ala Val Ser Tyr Ala Thr Asp Asp Gly Gln Asn Leu Ser Gly Lys Phe  
115 120 125

-continued

---

Lys Ala Asp Lys Pro Gly Arg Tyr Tyr Ile His Leu Tyr Met Phe Asn  
130 135 140

Gly Ser Tyr Met Pro Tyr Arg Ile Asn Ile Glu Gly Ser Val Gly Arg  
145 150 155 160

<210> SEQ ID NO 10  
<211> LENGTH: 152  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic PTH(7-33)-CBD fusion protein

<400> SEQUENCE: 10

Leu Met His Asn Leu Gly Lys His Leu Asn Ser Met Glu Arg Val Glu  
1 5 10 15

Trp Leu Arg Lys Lys Leu Gln Asp Val His Asn Gly Ile Asn Ser Pro  
20 25 30

Val Tyr Pro Ile Gly Thr Glu Lys Glu Pro Asn Asn Ser Lys Glu Thr  
35 40 45

Ala Ser Gly Pro Ile Val Pro Gly Ile Pro Val Ser Gly Thr Ile Glu  
50 55 60

Asn Thr Ser Asp Gln Asp Tyr Phe Tyr Phe Asp Val Ile Thr Pro Gly  
65 70 75 80

Glu Val Lys Ile Asp Ile Asn Lys Leu Gly Tyr Gly Gly Ala Thr Trp  
85 90 95

Val Val Tyr Asp Glu Asn Asn Asn Ala Val Ser Tyr Ala Thr Asp Asp  
100 105 110

Gly Gln Asn Leu Ser Gly Lys Phe Lys Ala Asp Lys Pro Gly Arg Tyr  
115 120 125

Tyr Ile His Leu Tyr Met Phe Asn Gly Ser Tyr Met Pro Tyr Arg Ile  
130 135 140

Asn Ile Glu Gly Ser Val Gly Arg  
145 150

<210> SEQ ID NO 11  
<211> LENGTH: 35  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic PTH(1-33) with Gly-Ser amino terminal  
extension

<400> SEQUENCE: 11

Gly Ser Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His  
1 5 10 15

Leu Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp  
20 25 30

Val His Asn  
35

<210> SEQ ID NO 12  
<211> LENGTH: 6  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic thrombin cleavage sequence

<400> SEQUENCE: 12

Leu Val Pro Arg Gly Ser  
1 5

-continued

---

```

<210> SEQ ID NO 13
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Clostridium histolyticum

<400> SEQUENCE: 13

Leu Lys Glu Lys Glu Asn Asn Asp Ser Ser Asp Lys Ala Thr Val Ile
1          5          10          15
Pro Asn Phe Asn Thr Thr Met Gln Gly Ser Leu Leu Gly Asp Asp Ser
          20          25          30
Arg Asp Tyr Tyr Ser Phe Glu Val Lys Glu Glu Gly Glu Val Asn Ile
          35          40          45
Glu Leu Asp Lys Lys Asp Glu Phe Gly Val Thr Trp Thr Leu His Pro
          50          55          60
Glu Ser Asn Ile Asn Asp Arg Ile Thr Tyr Gly Gln Val Asp Gly Asn
65          70          75          80
Lys Val Ser Asn Lys Val Lys Leu Arg Pro Gly Lys Tyr Tyr Leu Leu
          85          90          95
Val Tyr Lys Tyr Ser Gly Ser Gly Asn Tyr Glu Leu Arg Val Asn
          100          105          110

```

```

<210> SEQ ID NO 14
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: C. sporogenes

<400> SEQUENCE: 14

Ile His Glu Lys Glu Asn Asn Asp Ser Phe Glu Ser Ala Asn Lys Ile
1          5          10          15
Val Leu Asn Ala Pro Ile Leu Gly Ser Leu Asn Gly Glu Asp Leu Arg
          20          25          30
Asp Ile Tyr Ser Phe Glu Ile Lys Glu Thr Lys Asp Leu Asn Ile Lys
          35          40          45
Leu Thr Asn Leu Asn Asn Leu Gly Leu Thr Trp Thr Leu Tyr Lys Glu
          50          55          60
Ser Asp Leu Asn Asn Tyr Ile Ala Tyr Gly Ser Lys Leu Gly Ser Thr
65          70          75          80
Ile Val Gly Asn Cys His Val Thr Pro Gly Lys Tyr Tyr Leu Tyr Val
          85          90          95
Tyr Lys Tyr Ser Gly Asn Asn Gly Asn Tyr Ser Leu Ile Ile Lys
          100          105          110

```

```

<210> SEQ ID NO 15
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: C. botulinum

<400> SEQUENCE: 15

Ile Tyr Glu Lys Glu Asn Asn Asp Ser Phe Glu Thr Ala Asn Lys Ile
1          5          10          15
Met Leu Asn Thr Thr Val Leu Gly Asn Leu Asn Gly Lys Asp Val Arg
          20          25          30
Asp Ile Tyr Ser Phe Asp Ile Lys Glu Ala Lys Asp Leu Asp Ile Lys
          35          40          45
Leu Asn Asn Leu Asn Asn Leu Gly Leu Ala Trp Asn Leu Tyr Lys Glu
          50          55          60
Ser Asp Leu Asn Asn Tyr Ile Ala Tyr Gly Ser Val Ser Gly Asn Thr
65          70          75          80

```

-continued

---

Ile Lys Gly Lys Cys Asn Val Ala Pro Gly Lys Tyr Tyr Leu Tyr Val  
85 90 95

Tyr Lys Tyr Ser Gly Asp Asn Gly Asn Tyr Ser Leu Ala Ile Lys  
100 105 110

<210> SEQ ID NO 16  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: *B. cereus*

<400> SEQUENCE: 16

Leu Thr Glu Ser Glu Pro Asn Asn Arg Pro Glu Glu Ala Asn Arg Ile  
1 5 10 15  
Gly Leu Asn Thr Thr Ile Lys Gly Ser Leu Ile Gly Gly Asp His Thr  
20 25 30  
Asp Val Tyr Thr Phe Asn Val Ala Ser Ala Lys Asn Ile Asn Ile Ser  
35 40 45  
Val Leu Asn Glu Tyr Gly Ile Gly Met Thr Trp Val Leu His His Glu  
50 55 60  
Ser Asp Met Gln Asn Tyr Ala Ala Tyr Gly Gln Val Asn Gly Asn His  
65 70 75 80  
Ile Glu Ala Asn Phe Asn Ala Lys Pro Gly Lys Tyr Tyr Leu Tyr Val  
85 90 95  
Tyr Lys Tyr Asp Asn Gly Asp Gly Thr Tyr Glu Leu Ser Val Lys  
100 105 110

<210> SEQ ID NO 17  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: *B. anthracis*

<400> SEQUENCE: 17

Leu Thr Glu Ser Glu Pro Asn Asn Arg Pro Glu Glu Ala Asn Arg Ile  
1 5 10 15  
Gly Leu Asn Thr Thr Ile Lys Gly Ser Leu Ile Gly Gly Asp His Thr  
20 25 30  
Asp Val Tyr Thr Phe Asn Val Ala Ser Ala Lys Asn Ile Asp Ile Ser  
35 40 45  
Val Leu Asn Glu Tyr Gly Ile Gly Met Thr Trp Val Leu His His Glu  
50 55 60  
Ser Asp Met Gln Asn Tyr Ala Ala Tyr Gly Gln Ala Asn Gly Asn His  
65 70 75 80  
Ile Glu Ala Asn Phe Asn Ala Lys Pro Gly Lys Tyr Tyr Leu Tyr Val  
85 90 95  
Tyr Lys Tyr Asp Asn Gly Asp Gly Thr Tyr Glu Leu Ser Val Lys  
100 105 110

<210> SEQ ID NO 18  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: *Bacillus* sp.

<400> SEQUENCE: 18

Lys Thr Glu Ile Glu Pro Asn Asn Arg Pro Glu Glu Ala Thr Met Leu  
1 5 10 15  
Pro Phe Asn Thr Pro Leu Ser Gly Ser Leu Met Glu Asp Asp His Thr  
20 25 30

-continued

---

```

Asp Val Tyr Glu Phe Asn Val Thr Ser Pro Lys Glu Ile Asp Ile Ser
   35                               40                               45

Val Leu Asn Glu Asn Gln Ile Gly Met Thr Trp Val Leu Tyr His Glu
   50                               55                               60

Ser Asp Ser Gln Asn Tyr Ala Ser Phe Gly Gln Glu Asp Gly Asn Met
   65                               70                               75                               80

Ile Asn Gly Lys Trp Asn Ala Lys Pro Gly Lys Tyr Tyr Leu Tyr Val
   85                               90                               95

Tyr Lys Phe Glu Asn Glu Asn Gly Thr Tyr Thr Val His Val Gln
   100                               105                               110

```

```

<210> SEQ ID NO 19
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: L. sphaericus

```

```

<400> SEQUENCE: 19

```

```

Lys Ala Glu Ile Glu Pro Asn Asn Arg Pro Glu Glu Ala Thr Ile Leu
  1      5      10      15

Pro Phe Asn Thr Pro Leu Lys Gly Arg Leu Met Asp Asp Asp His Thr
  20      25      30

Asp Val Tyr Glu Phe Asn Val Thr Ser Pro Lys Glu Leu Asp Ile Ser
  35      40      45

Val Leu Asn Glu Asn Arg Ile Gly Met Thr Trp Val Leu Tyr His Glu
  50      55      60

Ser Asp Ser Gln Asn Tyr Ala Ser Phe Gly Gln Glu Glu Gly Asn Met
  65      70      75      80

Ile Asn Gly Lys Leu His Ala Glu Pro Gly Lys Tyr Tyr Leu Tyr Val
  85      90      95

Tyr Lys Phe Glu Asn Glu Asn Gly Thr Tyr Thr Val Gln Val Gln
  100      105      110

```

```

<210> SEQ ID NO 20
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: C. cerus

```

```

<400> SEQUENCE: 20

```

```

Val Thr Glu Asn Glu Pro Asn Asn Glu Pro Arg Gln Ala Asn Lys Val
  1      5      10      15

Asn Phe His Thr Pro Val Lys Gly Thr Leu His Asn Ser Asp Arg Val
  20      25      30

Asp Val Phe Thr Phe Gln Ile Asp Ser Pro Glu Asn Ile Asn Ile Ser
  35      40      45

Leu Leu Asn Glu Gln Asn Ile Gly Met Thr Trp Val Leu His His Glu
  50      55      60

Ser Asp Leu Asn Asn Tyr Val Ala Tyr Gly Glu Asn Glu Gly Asn Val
  65      70      75      80

Val Lys Gly Thr Tyr Asn Ala Lys Pro Gly Lys Tyr Tyr Leu Tyr Val
  85      90      95

Tyr Lys Tyr Glu Asn Lys Asp Gly Ser Tyr Val Leu Asn Ile Lys
  100      105      110

```

```

<210> SEQ ID NO 21
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: B. mycoides

```



-continued

&lt;400&gt; SEQUENCE: 21

Ser Val Glu Lys Glu Pro Asn Asn Ser Phe Gln Thr Ala Asn Lys Leu  
 1 5 10 15  
 Gln Leu Asn Gln Leu Leu Arg Ala Ser Leu Gly Asn Gly Asp Thr Ser  
 20 25 30  
 Asp Tyr Phe Glu Ile Asn Val Glu Thr Ala Arg Asn Leu Gln Ile Asn  
 35 40 45  
 Val Thr Lys Glu Asn Asn Ile Gly Val Asn Trp Val Leu Tyr Ser Ala  
 50 55 60  
 Ala Asp Leu Asn Asn Tyr Val Thr Tyr Ala Gln Thr Gln Gly Asn Lys  
 65 70 75 80  
 Leu Val Gly Ser Tyr Asn Ala His Pro Gly Lys Tyr Tyr Leu His Val  
 85 90 95  
 Tyr Gln Tyr Gly Gly Gly Thr Gly Asn Tyr Thr Val Glu Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 112

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: B. weihensteph

&lt;400&gt; SEQUENCE: 22

Ala Val Glu Lys Glu Pro Asn Asn Ser Phe Asp Ala Ala Asn Pro Leu  
 1 5 10 15  
 Ser Leu Asn Ala Leu Leu Arg Gly Asn Leu Ser Asp Gln Asp Gln Val  
 20 25 30  
 Asp Arg Phe Val Ile Asp Val Lys Asp Pro Lys Asp Leu Gln Ile Thr  
 35 40 45  
 Val Thr Asn Glu Gln Asn Leu Gly Leu Asn Trp Val Leu Tyr Ser Glu  
 50 55 60  
 Ser Asp Leu Asn Asn Tyr Val Thr Tyr Ala Thr Lys Arg Asp Gly Asn  
 65 70 75 80  
 Lys Leu Leu Gly Asn Tyr Asn Ala Lys Pro Gly Lys Tyr Tyr Leu Ser  
 85 90 95  
 Val Tyr Lys Tyr Gly Gly Gly Thr Gly Asn Phe Thr Val Glu Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 111

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: B. brevis

&lt;400&gt; SEQUENCE: 23

Glu Lys Glu Gln Glu Pro Asn Asn Ser Phe Ser Glu Ala Asn Pro Leu  
 1 5 10 15  
 Lys Ser Asn Val Glu Leu Ser Gly Gln Thr Ser Lys Gln Asp Asp Lys  
 20 25 30  
 Asp Ile Phe Ala Leu Lys Val Leu Gly Asn Gly Thr Val Lys Ile Asn  
 35 40 45  
 Val Thr Ser Glu His Asp Thr Gly Leu Asn Trp Val Val His His Glu  
 50 55 60  
 Asp Asp Leu Asn Asn Tyr Leu Ala Tyr Pro Lys Thr Ser Gly Lys Thr  
 65 70 75 80

-continued

---

Leu Ser Gly Glu Phe Glu Ala Thr Pro Gly Thr Tyr Tyr Leu Ser Val  
85 90 95

Tyr Asn Phe Asn Gly Glu Thr Ile Pro Tyr Lys Val Thr Ala Glu  
100 105 110

<210> SEQ ID NO 24  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: B. brevis

<400> SEQUENCE: 24

Pro Thr Glu Val Glu Pro Asn Asn Ser Phe Asp Asp Ala Asn Thr Leu  
1 5 10 15

Gln Leu Gly Lys Glu Ile Ser Gly Gln Thr Asp Arg Thr Asp Asp Lys  
20 25 30

Asp Thr Tyr Met Ile Gln Val Glu Glu Glu Gly Val Ile Gln Val Thr  
35 40 45

Val Ser Ser Glu Lys Asp Glu Gly Leu Asn Trp Val Val Phe His Glu  
50 55 60

Asp Asp Leu Lys Thr Tyr Phe Ala Tyr Pro Lys Thr Thr Gly Lys Lys  
65 70 75 80

Leu Thr Gly Glu Phe Glu Ala Lys Pro Gly Lys Tyr Tyr Leu Leu Val  
85 90 95

Tyr Asn Thr Asn Asn Thr Lys Ile Pro Tyr Lys Ala Ile Val Asn  
100 105 110

<210> SEQ ID NO 25  
<211> LENGTH: 112  
<212> TYPE: PRT  
<213> ORGANISM: C. perfringens

<400> SEQUENCE: 25

Ile Lys Glu Val Glu Asn Asn Asn Asp Phe Asp Lys Ala Met Lys Val  
1 5 10 15

Asp Ser Asn Ser Lys Ile Val Gly Thr Leu Ser Asn Asp Asp Leu Lys  
20 25 30

Asp Ile Tyr Ser Ile Asp Ile Lys Asn Pro Ser Asp Leu Asn Ile Val  
35 40 45

Val Glu Asn Leu Asp Asn Ile Lys Met Asn Trp Leu Leu Tyr Ser Ala  
50 55 60

Asp Asp Leu Ser Asn Tyr Val Asp Tyr Ala Asn Ala Asp Gly Asn Lys  
65 70 75 80

Leu Ser Asn Thr Cys Lys Leu Asn Pro Gly Lys Tyr Tyr Leu Cys Val  
85 90 95

Tyr Gln Phe Glu Asn Ser Gly Thr Gly Asn Tyr Thr Val Asn Leu Gln  
100 105 110

<210> SEQ ID NO 26  
<211> LENGTH: 115  
<212> TYPE: PRT  
<213> ORGANISM: C. sporogenes

<400> SEQUENCE: 26

Ile Ser Glu Lys Glu Asp Asn Asp Ser Phe Asp Lys Ala Asn Arg Val  
1 5 10 15

Gly Lys Asn Gln Thr Val Leu Ala Thr Leu Asp Thr Lys Asp Asn Arg  
20 25 30

Val 1	Ser	Glu	Lys	Glu 5	Asp	Asn	Asn	Asp	Phe 10	Thr	Thr	Ala	Asn	Pro 15	Val
Tyr	Tyr	Lys	Asp 20	Leu	Val	Asn	Gly	Ser 25	Val	Ser	Ser	Ser	Asp 30	Asn	Lys
Asp	Thr	Phe 35	Tyr	Phe	Thr	Val	Thr 40	Lys	Pro	Ser	Asp	Ile 45	Thr	Ile	Thr
Val 50	Glu	Lys	Thr	Asn	Asn	Asp 55	Lys	Ser	Glu	Phe	Asn 60	Trp	Leu	Leu	Phe
Ser 65	Asp	Glu	Asp	Lys 70	Ser	Asn	Tyr	Met	Ala	Phe 75	Pro	Asn	Lys	Glu 80	Leu
Gly	Asn	Gln	Leu	Ser 85	Asn	Thr	Val	Lys	Ile 90	Asn	Lys	Pro	Gly	Lys 95	Tyr

-continued

---

Tyr Leu Val Ile Tyr Lys Thr Leu Gly Glu Lys Val Asp Tyr Lys Phe  
                   100                  105                  110

Ser Ile Glu  
           115

<210> SEQ ID NO 29  
 <211> LENGTH: 115  
 <212> TYPE: PRT  
 <213> ORGANISM: C. botulinum

<400> SEQUENCE: 29

Val Ser Glu Lys Glu Asn Asn Asn Asp Tyr Val Asn Ala Asn Pro Val  
 1                  5                  10                  15

Tyr Ser Lys Asp Leu Val Asn Gly Ser Val Ser Ser Ser Asp Asp Arg  
           20                  25                  30

Asp Ile Phe Tyr Phe Asn Val Thr Lys Pro Ser Asp Ile Thr Ile Asn  
           35                  40                  45

Val Glu Lys Ile Asn Lys Asp Lys Ser Glu Phe Ser Trp Leu Leu Phe  
           50                  55                  60

Ser Glu Glu Asp Lys Ser Asn Tyr Ile Thr Tyr Pro Asn Lys Glu Leu  
           65                  70                  75                  80

Glu Asn Leu Phe Tyr Ser Thr Val Lys Ile Asp Lys Pro Gly Lys Tyr  
                   85                  90                  95

Tyr Leu Val Ile Tyr Lys Val Ser Gly Glu Lys Ser Asp Tyr Arg Phe  
                   100                  105                  110

Asn Ile Glu  
           115

<210> SEQ ID NO 30  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: C. sordellii

<400> SEQUENCE: 30

Gly Val Glu Gln Glu Asp Asn Asn Ser Phe Glu Lys Ala Asn Pro Phe  
 1                  5                  10                  15

Ser Ile Asn Gln Leu Val Lys Gly Glu Leu Asp Asn Asn Lys Asp Thr  
           20                  25                  30

Ser Asp Tyr Phe Lys Phe Glu Val Lys Glu Asp Ala Gln Leu Asn Ile  
           35                  40                  45

Ser Leu Glu Lys Thr Glu Gly Asp Gly Val Asn Trp Leu Leu Phe Lys  
           50                  55                  60

Asp Ser Asp Leu Glu Asn Tyr Ile Ala Ser Pro Thr Glu Ser Ile Asp  
           65                  70                  75                  80

Asn Lys Leu Asn Gly Lys Val Asp Leu Lys Val Gly Thr Tyr Tyr Leu  
                   85                  90                  95

Glu Val Tyr Gly Tyr Gly Ser Ser Pro Val Lys Tyr Asn Phe Lys Val  
                   100                  105                  110

Thr

<210> SEQ ID NO 31  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: Clostridium histolyticum

-continued

&lt;400&gt; SEQUENCE: 31

Thr Lys Glu Met Glu Pro Asn Asp Asp Ile Lys Glu Ala Asn Gly Pro  
 1 5 10 15  
 Ile Val Glu Gly Val Thr Val Lys Gly Asp Leu Asn Gly Ser Asp Asp  
 20 25 30  
 Ala Asp Thr Phe Tyr Phe Asp Val Lys Glu Asp Gly Asp Val Thr Ile  
 35 40 45  
 Glu Leu Pro Tyr Ser Gly Ser Ser Asn Phe Thr Trp Leu Val Tyr Lys  
 50 55 60  
 Glu Gly Asp Asp Gln Asn His Ile Ala Ser Gly Ile Asp Lys Asn Asn  
 65 70 75 80  
 Ser Lys Val Gly Thr Phe Lys Ser Thr Lys Gly Arg His Tyr Val Phe  
 85 90 95  
 Ile Tyr Lys His Asp Ser Ala Ser Asn Ile Ser Tyr Ser Leu Asn Ile  
 100 105 110  
 Lys

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 113

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *C. perfringens*

&lt;400&gt; SEQUENCE: 32

Ile Asn Glu Ser Glu Pro Asn Asn Asp Phe Glu Lys Ala Asn Gln Ile  
 1 5 10 15  
 Ala Lys Ser Asn Met Leu Val Lys Gly Thr Leu Ser Glu Glu Asp Tyr  
 20 25 30  
 Ser Asp Lys Tyr Tyr Phe Asp Val Ala Lys Lys Gly Asn Val Lys Ile  
 35 40 45  
 Thr Leu Asn Asn Leu Asn Ser Val Gly Ile Thr Trp Thr Leu Tyr Lys  
 50 55 60  
 Glu Gly Asp Leu Asn Asn Tyr Val Leu Tyr Ala Thr Gly Asn Glu Gly  
 65 70 75 80  
 Thr Val Leu Lys Gly Glu Lys Thr Leu Glu Pro Gly Arg Tyr Tyr Leu  
 85 90 95  
 Ser Val Tyr Thr Tyr Asp Asn Gln Ser Gly Ala Tyr Thr Val Asn Val  
 100 105 110  
 Lys

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 113

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *C. sordellii*

&lt;400&gt; SEQUENCE: 33

Ser Gln Glu Val Gly Asn Asp Asp Thr Phe Glu Thr Ala Asn Gly Pro  
 1 5 10 15  
 Ile Lys Ile Asn Thr Asn Tyr Ser Gly Asp Leu Ser Asp Thr Asp Asn  
 20 25 30  
 Lys Asp Tyr Tyr Tyr Phe Asn Leu Asp Asn Pro Ser Asn Ile Asn Ile  
 35 40 45  
 Thr Leu Glu Asn Leu Asp Asn Lys Gly Ile Ser Trp Gln Leu Phe His  
 50 55 60  
 Glu Ser Asp Leu Asn Asn Tyr Val Ala Tyr Pro Thr Thr Ser Gly Ala  
 65 70 75 80

-continued

---

Ile Leu Asn Gly Asp Tyr Asn Ala Thr Lys Pro Gly Lys Tyr Tyr Ile  
85 90 95

Leu Val Tyr Asn His Asp Lys Ser Ile Ala Asn Tyr Asn Leu Lys Val  
100 105 110

Asn

<210> SEQ ID NO 34  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: Clostridium histolyticum

<400> SEQUENCE: 34

Gly Thr Glu Lys Glu Pro Asn Asn Ser Lys Glu Thr Ala Ser Gly Pro  
1 5 10 15

Ile Val Pro Gly Ile Pro Val Ser Gly Thr Ile Glu Asn Thr Ser Asp  
20 25 30

Gln Asp Tyr Phe Tyr Phe Asp Val Ile Thr Pro Gly Glu Val Lys Ile  
35 40 45

Asp Ile Asn Lys Leu Gly Tyr Gly Gly Ala Thr Trp Val Val Tyr Asp  
50 55 60

Glu Asn Asn Asn Ala Val Ser Tyr Ala Thr Asp Asp Gly Gln Asn Leu  
65 70 75 80

Ser Gly Lys Phe Lys Ala Asp Lys Pro Gly Arg Tyr Tyr Ile His Leu  
85 90 95

Tyr Met Phe Asn Gly Ser Tyr Met Pro Tyr Arg Ile Asn Ile Glu  
100 105 110

<210> SEQ ID NO 35  
<211> LENGTH: 30  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(30)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 35

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro  
1 5 10 15

Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly  
20 25 30

<210> SEQ ID NO 36  
<211> LENGTH: 30  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(30)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 36

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro  
1 5 10 15

Xaa Gly Pro Xaa Ala Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly  
20 25 30

-continued

---

```

<210> SEQ ID NO 37
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(30)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 37

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro
1           5           10          15

Xaa Ala Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly
          20          25          30

<210> SEQ ID NO 38
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(30)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 38

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Ala Pro
1           5           10          15

Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly
          20          25          30

<210> SEQ ID NO 39
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(30)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 39

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Ala Pro Xaa Gly Pro
1           5           10          15

Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly
          20          25          30

<210> SEQ ID NO 40
<211> LENGTH: 24
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 40

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Cys Gly Pro Xaa Gly Pro
1           5           10          15

Xaa Gly Pro Xaa Gly Pro Xaa Gly
          20

```

-continued

---

<210> SEQ ID NO 41  
<211> LENGTH: 31  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(31)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 41

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Ala Pro  
1 5 10 15

Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Cys  
20 25 30

<210> SEQ ID NO 42  
<211> LENGTH: 22  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(22)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 42

Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly  
1 5 10 15

Pro Xaa Gly Pro Xaa Gly  
20

<210> SEQ ID NO 43  
<211> LENGTH: 31  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(31)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 43

Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly  
1 5 10 15

Pro Xaa Gly Pro Xaa Gly Pro Arg Gly Pro Arg Gly Pro Arg Gly  
20 25 30

<210> SEQ ID NO 44  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic peptide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(9)  
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

<400> SEQUENCE: 44

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly  
1 5



-continued

---

```

<210> SEQ ID NO 45
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

```

```

<400> SEQUENCE: 45

```

```

Pro Xaa Gly Pro Xaa Gly Pro Xaa Ala
1             5

```

```

<210> SEQ ID NO 46
<211> LENGTH: 26
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(26)
<223> OTHER INFORMATION: Any Xaa is hydroxyproline

```

```

<400> SEQUENCE: 46

```

```

Gly Pro Arg Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly
1             5             10            15

```

```

Pro Xaa Gly Pro Xaa Gly Pro Xaa Gly Cys
20            25

```

```

<210> SEQ ID NO 47
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide: GST Tag

```

```

<400> SEQUENCE: 47

```

```

Gly Ser Pro Gly Ile Pro Gly
1             5

```

---

We claim:

1. A method of treating a subject having alopecia that is not chemotherapy-induced comprising administering a composition comprising a bacterial collagen-binding polypeptide segment linked to a PTH/PTHrP receptor agonist to the subject to increase hair growth in the subject in need thereof for treatment of alopecia that is not chemotherapy-induced, wherein the bacterial collagen-binding polypeptide segment comprises a collagen-binding polypeptide derived from an M9 peptidase from *Clostridium*, *Bacillus* and *Vibrio* selected from the group consisting of one of SEQ ID NOs: 13-34 or a fragment of at least 8 consecutive amino acids of SEQ ID NOs: 13-34, residues 34-158 of SEQ ID NO: 1, a fragment of at least 8 consecutive amino acids from residues 34-158 of SEQ ID NO: 1, or a peptide that is at least 90% identical to residues 34-158 of SEQ ID NO: 1 or SEQ ID NOs: 13-34 and wherein the PTH/PTHrP receptor agonist comprises residues 1-33 of SEQ ID NO: 1, PTH (SEQ ID NO: 7), residues 1-14 of SEQ ID NO: 1, residues 1-34 of SEQ ID NO: 7 or a fragment of at least 8 consecutive amino acids from residues 1-34 of SEQ ID NO: 7.

2. The method of claim 1, wherein the alopecia is alopecia areata or is related to male pattern baldness or polycystic ovarian syndrome.

3. The method of claim 1, wherein the composition is administered locally.

4. The method of claim 3, wherein the composition is administered topically.

5. The method of claim 1, wherein the PTH/PTHrP receptor agonist is a polypeptide and the N-terminus of the collagen-binding polypeptide segment is linked directly or through a linker polypeptide segment to the C-terminus of the PTH/PTHrP receptor agonist polypeptide or wherein the collagen-binding polypeptide segment and the therapeutic agent are chemically cross-linked to each other or are polypeptide portions of a fusion protein.

6. The method of claim 1, wherein the subject is a human.

7. The method of claim 1, wherein the composition is administered in aqueous solution at pH below about 5.0 or above about 6.0.

8. The method of claim 1, wherein the bacterial collagen-binding polypeptide segment comprises one of SEQ ID NOs: 13-34, residues 34-158 of SEQ ID NO: 1, or a peptide that is at least 90% identical to residues 34-158 of SEQ ID NO: 1 or SEQ ID NOs: 13-34.

9. The method of claim 1, wherein the PTH/PTHrP receptor agonist comprises residues 1-33 of SEQ ID NO: 1, PTH (SEQ ID NO: 7), residues 1-14 of SEQ ID NO: 1, residues 1-34 of SEQ ID NO: 7.

73

10. The method of claim 1, wherein the bacterial collagen-binding polypeptide segment linked to the PTH/PTHrP receptor agonist is a fusion protein comprising SEQ ID NO: 1.

11. The method of claim 1, wherein the bacterial collagen-binding polypeptide segment linked to the PTH/PTHrP receptor agonist is a fusion protein comprising SEQ ID NO: 2.

12. The method of claim 1, wherein the collagen-binding polypeptide segment further comprises residues 807-900 of SEQ ID NO: 6.

\* \* \* \* \*

74