

2006

Hyperinfection with the Bass Tapeworm, *Proteocephalus ambloplites* (Cestoda), in the Black Basses *Micropterus punctulatus* and *M. dolomieu* from Certain Arkansas Reservoir Lakes

James J. Daly Sr.

University of Arkansas for Medical Sciences, jamesdalysr@yahoo.com

Randal J. Keller

Murray State University

Bruce DeYoung

University of Arkansas for Medical Sciences

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Part of the [Animal Diseases Commons](#), and the [Zoology Commons](#)

Recommended Citation

Daly, James J. Sr.; Keller, Randal J.; and DeYoung, Bruce (2006) "Hyperinfection with the Bass Tapeworm, *Proteocephalus ambloplites* (Cestoda), in the Black Basses *Micropterus punctulatus* and *M. dolomieu* from Certain Arkansas Reservoir Lakes," *Journal of the Arkansas Academy of Science*: Vol. 60, Article 27. Available at: <https://scholarworks.uark.edu/jaas/vol60/iss1/27>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in *Journal of the Arkansas Academy of Science* by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

Hyperinfection with the Bass Tapeworm, *Proteocephalus ambloplites* (Cestoda), in the Black Basses *Micropterus punctulatus* and *M. dolomieu* from Certain Arkansas Reservoir Lakes

JAMES J. DALY SR.^{1,3}, RANDAL J. KELLER², AND BRUCE DEYOUNG¹

¹Department of Microbiology and Immunology, University of Arkansas for Medical Sciences, Little Rock, AR 72205

²Department of Occupational Safety and Health, Murray State University, P.O. Box 9, Murray, KY 42071

Correspondence: jjdalysr@artelco.com

Proteocephalus ambloplites is a tapeworm that lives as an adult in the intestinal tract of black bass (*Micropterus* spp.). The life cycle is complex and hosts include first a microcrustacean followed by a small fish as a second intermediate host. In the small fish the parasite transforms into a metacystode larval stage, the plerocercoid, which migrates through the gut into the coelom where it awaits ingestion by the final host. The plerocercoid form of *P. ambloplites* can be found in a number of freshwater fishes that can act as intermediate hosts (Hoffman 1999). When a bass eats the infected small fish, the plerocercoids penetrate the gut and invade parenteric organs, most importantly, the gonads which can cause reduced fertility in the host fish. Certain plerocercoids under the stimulus of a water temperature change will reinvade the gut to become tapeworm adults. (Esch and Huffines 1969, Fischer and Freeman 1969, Fischer 1973). The major importance of this parasite is the effect on bass reproduction when the bass tapeworm plerocercoids are in large numbers in gonadal tissue.

All three major *Micropterus* species (*M. salmoides*, *M. dolomieu*, and *M. punctulatus*) are known to be infected with both the adult tapeworms and their plerocercoid larvae. Preliminary data from our necropsies of bass for infections with *Clinostomum marginatum* metacercaria indicated that *P. ambloplites* infections were heavier in Arkansas than previously reported. This study was initiated to determine the extent of these infections in black bass from Arkansas lakes. Bass were collected from tournament fishermen in 1988-89 from selected Arkansas reservoir lakes, transported to the University of Arkansas for Medical Sciences and necropsied for the presence of plerocercoids of *P. ambloplites*. Definition of population characteristics followed that of Bush et al. (1997) and were as follows: Prevalence (percent of fish infected), maximum abundance (largest number of parasites in a single host), mean abundance (average number of parasites per fish) and its standard deviation. The findings of this study are summarized in Table 1. Mean intensity (average number of parasites for infected fish only) and its standard deviation are not included since this report was to compare the extent of the plerocercoid infections in the Arkansas reservoirs with older literature and in those studies these values are not given. The variance of a mean intensity reported with 100% prevalence, as seen in Table 1, is equivalent to the variance of a mean abundance of 100% prevalence.

Very few (<0.1 percent) adult tapeworms were found in

any of the collections. Also, the plerocercoid numbers in the present study may be a result of seasonal effects (late spring, summer) because the plerocercoid larvae are generally not found in bass during the winter in northern climes, but reoccur in large numbers again in the spring (Esch and Huffines 1973, Fischer 1973). However, Eure (1976) found that plerocercoids could be found year-round in a reservoir in South Carolina. Studies in Arkansas have found plerocercoids in Arkansas black bass in the winter (Cloutman 1975, Kilambi and Becker 1977), but any temporal relationship between adult tapeworms and plerocercoids in the State's reservoirs is still to be definitively determined.

Table 1 shows data from this study, data from other Arkansas waters, and other selected locales from North America that have the highest reported abundances of plerocercoid infection. The current data for spotted bass (*M. punctulatus*) and largemouth data (*M. salmoides*) are both from Lake Maumelle in Pulaski County. The infected smallmouth (*M. dolomieu*) were collected from Bull Shoals Lake. All of the data in Table 1 represent the largest mean abundances for individual collections in each study but collections of other bass hosts were also made from the same bodies of water, but with smaller abundances being reported in the other bass populations. An exception is the Gull Lake, Michigan data which included all bass collections that were made in that locale for a given year.

Table 1 shows that the infrapopulations of *P. ambloplites* plerocercoids in spotted bass from Lake Maumelle are hyperinfections and among the highest reported for bass anywhere and are the highest ever for spotted bass. The intensity of the smallmouth infection from Bull Shoals Lake, although not as high, can also be considered a hyperinfection based on comparative literature reports. Largemouth bass in Lake Maumelle have many fewer plerocercoids than seen in the other two bass. This discrepancy between largemouth and other bass has also been noted before in the Gull Lake study (Gilliland and Muzzall 2004). They found that largemouth plerocercoid prevalence was 100% with a mean intensity of 18.1 (± 12.9 SD). This agrees with the degree of infection seen in the largemouth bass from Lake Maumelle in Arkansas. It might seem that the largemouth may have a natural resistance to tapeworm infections, relative to the other 2 species. However, Szalai and Dick (1990) found infected largemouth in a reservoir in Saskatchewan, Canada to have plerocercoid numbers equivalent

James J. Daly Sr., Randal J. Keller, and Bruce DeYoung

Table 1. Population parameters of selected plerocercoid infections of the bass tapeworm, *Proteocephalus ambloplites*, in Arkansas and North America. LM = largemouth, SM = smallmouth, and SP = spotted bass. Only the parameters of prevalence, maximum abundance, and mean abundance (standard deviation) are listed. Daly et al. refers to the present study. ---- are unrecorded data.

Investigators	N	Locale	Bass	Prevalence	Max. Abundance	Abundance
Daly et al. this study	17	Arkansas	SM	100	117	35.5 (± 34)
Daly et al. this study	55	Arkansas	SP	100	200	66.3 (± 43)
Daly et al. this study	8	Arkansas	LM	75	52	12.4 (± 17)
Becker et al., 1978	347	Arkansas	SP	91.5	----	12.3 (----)
Cloutman, 1975	89	Arkansas	LM	----	----	7.4 (----)
Kilambi and Becker, 1977	12	Arkansas	SM	4	7	0.3 (± 2.02)
Gilliland and Muzzall, 2004	54	Michigan	SM	100	200	72.5 (± 44.8)**
Szalai and Dick, 1990	8*	Saskatchewan	LM	100	----	99.0 (± 116)**

*Largest bass hosts only

**Mean intensities (SD)

to those found in Arkansas and Michigan smallmouth and Arkansas spotted black bass (Table 1). The reasons for these geographical variations between the black bass species are unknown and could be due to a number of regional factors such as habitat preference or forage fish diet and perhaps even an inherent strain immunity (The Saskatchewan largemouth were hatchery-raised and introduced).

The most important pathology produced by these tapeworm larvae is destruction of the host gonadal tissue, but in Arkansas reservoirs the overall effect of such large numbers of plerocercoids on bass fecundity is yet to be ascertained.

ACKNOWLEDGMENTS.—The authors wish to thank bass tournament fishermen from the Bull Shoals' Firemen's Association, Mountain Home, Arkansas; Brady Mountain Resort, Mount Ida; and especially Roger Nesuda of Jolly Roger's Marina on Lake Maumelle. This work was done with the cooperation of the Arkansas Game and Fish Commission which provided the permits and some research support.

Literature Cited

- Becker, DA, WD Carr, DG Cloutman, WA Evans, RG Heard, PD Holmes, MD Norman, and WB Owens, Jr. 1978. Pre- and post impoundment ichthyoparasite succession in a new Arkansas Reservoir. Water Resources Research Center, Publication Number 54, University of Arkansas, Fayetteville, Arkansas.
- Bush, AO, KD Lafferty, JF Lotz and AW Shostak. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *The Journal of Parasitology* 83:575-583.
- Cloutman DG. 1975. Parasite community structure of largemouth bass, warmouth bass, and bluegill in Lake Fort Smith, Arkansas. *Transactions of the American Fisheries Society* 104: 277-283.
- Esch GW and WJ Huffines. 1973. Histopathology associated with endoparasitic helminths in bass. *The Journal of Parasitology* 59:306-313.
- Eure H. 1976. Seasonal abundance of *Proteocephalus ambloplites* (Cestoidea: Proteocephalidea) from largemouth bass living in a heated reservoir. *Parasitology* 73:205-212.
- Fischer H and RS Freeman. 1969. Penetration of parenteral plerocercoids of *Proteocephalus ambloplites* (Leidy) in the gut of smallmouth bass. *The Journal of Parasitology* 55:766-774.
- Fischer H. 1973. The role of plerocercoids in the biology of *Proteocephalus ambloplites* (Cestoda) maturing in smallmouth bass. *Canadian Journal of Zoology* 51:133-141.
- Gilliland MG and PM Muzzall. 2004. Microhabitat analysis of bass tapeworm in smallmouth bass, *Micropterus dolomieu*, and largemouth bass, *Micropterus salmoides*. *Comparative Parasitology* 71:221-225.
- Hoffman GL. 1999. *Parasites of North American Freshwater Fishes*. 2nd ed. Ithaca and London, Comstock Publishing Associates (Cornell University Press). 539p.
- Kilambi RV and D Becker. 1977. Population dynamics and species diversity of ichthyoparasitofauna of the Buffalo National River. Water Resources Research Center, Publication No. 48, University of Arkansas, Fayetteville, Arkansas.
- Szalai AJ and TA Dick. 1990. *Proteocephalus ambloplites* and *Contracaecum* sp. from largemouth bass stocked into Boundry Reservoir, Saskatchewan. *The Journal of Parasitology* 76:598-601.