

2002

Parasites of Four Endemic Plethodon from Arkansas and Oklahoma

Chris T. McAllister

Texas A&M University-Texarkana

Charles R. Bursey

Pennsylvania State University-Shenango Campus

Stanley E. Trauth

Arkansas State University

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Part of the [Animal Diseases Commons](#), and the [Zoology Commons](#)

Recommended Citation

McAllister, Chris T.; Bursey, Charles R.; and Trauth, Stanley E. (2002) "Parasites of Four Endemic Plethodon from Arkansas and Oklahoma," *Journal of the Arkansas Academy of Science*: Vol. 56, Article 36.

Available at: <https://scholarworks.uark.edu/jaas/vol56/iss1/36>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in *Journal of the Arkansas Academy of Science* by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

Parasites of Four Species of Endemic *Plethodon* from Arkansas and Oklahoma

Chris T. McAllister*

Department of Biology
Texas A&M University-Texarkana
Texarkana, TX 75505

Charles R. Bursey

Department of Biology
Pennsylvania State University-Shenango Campus
Sharon, PA 16146

Stanley E. Trauth

Department of Biological Sciences
Arkansas State University
State University, AR 72467

*Corresponding Author

The Caddo Mountain salamander, *Plethodon caddoensis* Pope and Pope, 1951, is restricted to the Caddo and Ouachita Mountains of Howard, Montgomery, Pike, and Polk Counties of western Arkansas; the Kiamichi slimy salamander, *P. kiamichi* Highton, 1989, is known only from the Kiamichi and Round Mountain outcroppings of Polk County of western Arkansas; the Rich Mountain salamander, *P. ouachitae* Dunn and Heinze, 1933, occurs on Rich Mountain and adjacent ridges of the Ouachita Mountains of Polk County, Arkansas, and eastern Oklahoma; the southern redback salamander, *P. serratus* Grobman, 1944, exists in four isolated populations, west-central Arkansas and southeastern Oklahoma, central Louisiana, central and southeastern Missouri, and the Piedmont and Blue Ridge provinces of northeastern Georgia, adjacent Alabama, eastern Tennessee and western North Carolina (Highton et al., 1989; Conant and Collins, 1998; Trauth et al., 2003). To our knowledge, there is only one report (Winter et al., 1986) of parasites from *P. caddoensis* and *P. ouachitae*, and Arkansas populations of *P. serratus*. The purpose of this note is to report additional parasites from *P. caddoensis*, *P. ouachitae*, and *P. serratus*, and for the first time, parasites from *P. kiamichi*.

Between December 1988 and February 2002, 72 plethodontid salamanders were collected by hand from several counties within the Ouachita National Forest of Arkansas and Oklahoma: *P. caddoensis* from springs or abandoned mines and *P. kiamichi*, *P. ouachitae*, and *P. serratus* from beneath decaying logs and leaf litter in seepage areas of deciduous forest habitat. Salamanders were placed in individual bags and transported on ice to the laboratory where they were killed within 48 hr of capture by prolonged immersion in a dilute chloroform solution. Methods for salamander necropsy, coccidial isolation, and preparation and staining of blood smears and helminths follow McAllister and Upton (1987) and Upton et al. (1993); preparation of mites follows McAllister et al. (1995d).

Collection sites, sample size, mean \pm 1 SD snout-vent length (SVL) in mm, and host accession numbers for voucher specimens deposited in the Arkansas State University Museum of Zoology (ASUMZ) for each species are listed in Appendix 1. Selected voucher specimens of parasites were deposited in the U.S. National Parasite Collection (UNSPC) and their accession numbers are listed

in Table 1.

Of the 72 salamanders collected, 32 (44%) harbored parasites: 11 (39%) *P. caddoensis*, four (25%) *P. kiamichi*, seven (88%) *P. ouachitae*, and 10 (50%) *P. serratus*. Blood smears were negative for intraerythrocytic hematozoa and viral and/or rickettsial inclusions. Parasites found in this study with their intensity of infection are listed in Table 1.

Cepedietta michiganensis (Woodhead, 1928) Corliss, de Puytorac, and Lom, 1965, was originally described as *Haptophyra michiganensis* by Woodhead (1928). Joy and Tucker (2001) have summarized hosts and localities. In Arkansas, it has been reported from the western slimy salamander (*P. albagula*), Fourche Mountain salamander (*P. fourchensis*), and *P. ouachitae* (Winter et al., 1986; McAllister et al., 1993). *Plethodon serratus* represents a new host record for *C. michiganensis* and Oklahoma a new locale for the parasite.

One of the *P. kiamichi* (ASUMZ 18982, male, 65 mm SVL, collected on 23 April 1993) was found to be passing eimerian oocysts in the feces. Unfortunately, only a few oocysts were present and not enough completed sporulation to allow for specific identification; however, oocysts of this isolate clearly contained four sporocysts, a taxonomic characteristic of the genus *Eimeria*. *Plethodon kiamichi* represents a new host record for *Eimeria* sp. This is only the fourth time a coccidian has ever been reported from plethodontid salamanders (Saxe, 1955; McAllister et al., 1993; Upton et al., 1993).

Cylindrotaenia idahoensis (Waitz and Mehra, 1961) Jones, 1987 was originally described from the Coeur d'Alene salamander, *Plethodon idahoensis* from Kootenai County, Idaho (Waitz and Mehra, 1961). It has been reported from Jordan's redcheek salamander, *P. jordani* from North Carolina (Dyer, 1983; Jones, 1987) and the western redback salamander, *P. vehiculum* from Oregon (Panitz, 1969). *Plethodon caddoensis*, *P. ouachitae* and *P. serratus* represent new host records for *C. idahoensis*. The Ouachita National Forest of Arkansas and Oklahoma are new locality records for *C. idahoensis*.

Batracholandrois magnavulvaris (Schad, 1960) Petter and Quentin, 1976 was originally described as *Oxyuris magnavulvaris* by Rankin (1937) from the red-spotted newt (*Notophthalmus viridescens*) and several species of plethodontid salamanders from Buncombe County, North Carolina. It

Parasites of Four Species of Endemic *Plethodon* from Arkansas and OklahomaTable 1. Parasites of endemic *Plethodon* spp. from Arkansas and Oklahoma.

Host species	<i>P. caddoensis</i>	<i>P. kiamichi</i>	<i>P. ouachitae</i>	<i>P. serratus</i>	
Number examined	(28)	(16)	(8)	(20)	
Parasite	Number of infected hosts and intensities (mean \pm 1SD, range)				USNPC Accession No.
Protista					
<i>Cepedietta michiganensis</i> **	---	---	4 (50%)	1 (5%)*	84342,92564,92565
<i>Eimeria</i> sp.	---	1 (6%)*	---	---	---
Cestoidea					
<i>Cylindrotaenia idahoensis</i> **	9 (32%)* 2.8 \pm 1.6 (2-5)	---	2 (25%)* 2.5 \pm 0.7 (2-3)	3 (16%)* 2.3 \pm 0.6 (2-3)	84300, 84339, 84340
Nematoda					
<i>Batracholandroides magnavulvaris</i> **	---	---	3 (38%) 1.0 \pm - (1)	8 (40%) 1.6 \pm 0.7 (1-3)	92042
<i>Cosmocercoides variabilis</i>	1 (4%)* 2.0 \pm - (2)	---	3 (38%)* 1.3 \pm 0.6 (1-2)	---	84256
<i>Oswaldocruzia euryceae</i>	---	2 (13%)* 3.5 \pm 2.1 (2-5)	3 (38%) 4.6 \pm 3.8 (2-9)	---	84254
Arthropoda					
<i>Hannemania</i> sp.	2 (7%)	1 (6%)*	6 (75%)	---	84341

*New host records.

**New locality record (Ouachita National Forest, Arkansas and/or Oklahoma).

has been reported from the Ouachita dusky salamander, *Desmognathus brimleyorum*, *P. caddoensis*, *P. fourchensis*, *P. ouachitae*, and *P. serratus* from Arkansas (Winter et al., 1986; McAllister et al., 1995d), and many species of salamanders from other North American locations (hosts and localities summarized by Joy and Tucker, 2001). The Ouachita National Forest of Oklahoma is a new locality record for *B. magnavulvaris*.

Cosmocercoides variabilis (Harwood, 1930) Travassos, 1931 was originally described from Woodhouse's toad, *Bufo woodhousii* collected in Texas (Harwood, 1930). It has been reported from the ringed salamander, *Ambystoma annulatum* from Arkansas (McAllister et al., 1995b) and various other hosts (see Baker, 1987). It should be noted that a similar species, *Cosmocercoides dukae*, a parasite of gastropods, has been reported from numerous amphibians (Baker, 1987). The major difference between the two species is the number of caudal papillae: *C. dukae* with 12 pairs of plectenes (Holl, 1928); *C. variabilis* with 14-20 pairs of plectenes (Harwood, 1930). Because our specimens had 16 pairs of plectenes, we have assigned them to *C. variabilis*. *Plethodon caddoensis* and *P. ouachitae* represent new host records for *C. variabilis*.

Oswaldocruzia euryceae Reiber, Byrd, and Parker, 1940 was originally described from the three-lined salamander, *Eurycea longicauda guttolineata* collected in Georgia (Reiber et

al., 1940). It has been reported from *P. caddoensis*, *P. ouachitae*, and *P. serratus* (Winter et al., 1986). *Plethodon kiamichi* represents a new host record for *O. euryceae*.

Larval intradermal mites, *Hannemania* sp. was found encapsulated in three species *P. caddoensis*, *P. ouachitae*, and *P. kiamichi*. Because only larvae were found, specific identity was not possible. *Hannemania* sp. has also been reported in Arkansas on *D. brimleyorum* (Loomis, 1956; Winter et al., 1986; McAllister et al., 1995d), graybelly salamanders, *E. multiplicata griseogaster* (McAllister et al., 1995c), and pickerel frogs, *Rana palustris* (McAllister et al., 1995a). *Plethodon kiamichi* represents a new host record for larva of *Hannemania* sp.

In summary, nine new host records and three new locality records are reported for parasites of four endemic species of *Plethodon* from the Ouachita Province of Arkansas and Oklahoma. Our survey supports Aho's (1990) suggestion of a depauperate noninteractive community structure observed in helminth communities of most amphibians and reptiles.

We wish to thank the Oklahoma Department of Wildlife Conservation and Arkansas Game and Fish Commission for Scientific Collecting Permits Nos. 3172, 1480 and 1048 to C.T.M. and S.E.T., respectively. We also thank the Arkansas State University Natural History Class (Spring 1991), Mr.

Appendix 1. Voucher specimens of host salamanders (ASUMZ accession numbers) and collection localities in Arkansas and Oklahoma.

Species	SVL \pm 1 SD mm (range)	ASUMZ	Localities and sample sizes (parentheses) counties; Township, Range, Section
<i>P. caddoensis</i>	42 \pm 4 (32-47)	18519-18520	Montgomery Co., AR; T3S, R27W, S26 (n = 2) Polk Co., AR; T4S, R29W, S6 (n = 23) T3S, R29W, S26 (n=3)
<i>P. kiamichi</i>	57 \pm 16 (27-73)	17575-17585; 17661-17663	Polk Co., AR; T1S, R32W, S31 (n = 16)
<i>P. ouachitae</i>	49 \pm 9 (29-59)	19492	Polk Co., AR; T1S, R31W, S7 (n = 8)
<i>P. serratus</i>	40 \pm 6 (27-49)	19491; 26396-26402	Hot Spring Co., AR; T5S, R20W, S31 (n = 7) Perry Co., AR; T3N, R20W, S27 (n = 1) Pike Co., AR; T6S, R25W, S14 (n = 3) Polk Co., AR; T1S, R32W, S10 (n = 3) LeFlore Co., OK; T1S, R32W, S7 (n = 2) McCurtain Co., OK; T5S, R25E, S10 (n = 4)

David Saugey (U.S. Forest Service), and Dr. David W. Allard (TAMU-T) for assistance in collecting, and Dr. Steve J. Upton (Kansas St. Univ.) for examining some salamanders for coccidia.

Literature Cited

- Aho, J. M.** 1990. Helminth communities in amphibians and reptiles: comparative approaches to understanding patterns and processes. Pp. 157-195 in G. W. Esch, A. O. Bush, and J. M. Aho, eds. *Parasite Communities: Patterns and Processes*. Chapman and Hall, New York.
- Baker, M. R.** 1987. Synopsis of the Nematoda parasitic in amphibians and reptiles. *Mem. Univ. Newfoundland Occas. Pap. Biol.* 11:1-325.
- Conant, R. and J. T. Collins.** 1998. *A field guide to reptiles and amphibians of eastern and central North America*. 3rd ed, expanded. Houghton Mifflin Co., Boston, 616 pp.
- Dyer, W. G.** 1983. A comparison of the helminth fauna of two *Plethodon jordani* populations from different altitudes in North Carolina. *Proc. Helm. Soc. Washington* 50:257-260.
- Highton, R., G. C. Maha, and L. R. Maxson.** 1989. Biochemical evolution in the slimy salamanders of the *Plethodon glutinosus* complex in the eastern United States. *Illinois Biol. Monogr.* 57:1-153.
- Jones, M. K.** 1987. A taxonomic revision of the Nematotaeniidae Lühe, 1910 (Cestoda: Cyclophyllidae). *Syst. Parasitol.* 10:165-245.
- Joy, J. E. and R. B. Tucker.** 2001. *Cepedietta michiganensis* (Protozoa) and *Batracholondros magnavulvaris* (Nematoda) from plethodontid salamanders in West Virginia, U.S.A. *Comp. Parasitol.* 68:185-189.
- Harwood, P. D.** 1930. A new species of *Oxysomatium* (Nematoda) with some remarks on the genera *Oxysomatium* and *Aplectana* and observations on life history. *J. Parasitol.* 17:61-73.
- Holl, F. J.** 1928. Two new nematode parasites. *J. Elisha Mitchell Sci. Soc.* 43:184-186.
- Loomis, R. B.** 1956. The chigger mites of Kansas (Acarina, Trombiculidae). *Univ. Kansas Sci. Bull.* 37:1-1443.
- McAllister, C. T. and S. J. Upton.** 1987. Endoparasites of the smallmouth salamander, *Ambystoma texanum* (Caudata: Ambystomatidae) from Dallas County, Texas. *Proc. Helm. Soc. Washington* 54:258-261.
- McAllister, C. T., S. J. Upton, and S. E. Trauth.** 1993. Endoparasites of western slimy salamanders, *Plethodon albagula* (Caudata: Plethodontidae), from Arkansas. *J. Helm. Soc. Washington.* 60:124-126.
- McAllister, C. T., C. R. Bursey, and S. E. Trauth.** 1995a. Parasites of the pickerel frog, *Rana palustris* (Anura: Ranidae) from the southern part of its range. *Southwest. Nat.* 40:111-116.
- McAllister, C. T., S. E. Trauth, and B. G. Cochran.** 1995b. Endoparasites of the ringed salamander, *Ambystoma annulatum* (Caudata: Ambystomatidae), from Arkansas. *Southwest. Nat.* 40:327-330.
- McAllister, C. T., S. E. Trauth, and C. R. Bursey.** 1995c. Metazoan parasites of the graybelly salamander, *Eurycea*

- multiplicata griseogaster* (Caudata: Plethodontidae), from Arkansas. J. Helm. Soc. Washington 62:66-69.
- McAllister, C. T., C. R. Bursley, S. J. Upton, S. E. Trauth, and D. B. Conn.** 1995d. Parasites of *Desmognathus brimleyorum* (Caudata: Plethodontidae) from the Ouachita Mountains of Arkansas and Oklahoma. J. Helm. Soc. Washington 62:150-156.
- Panitz, E.** 1969. Helminth parasites of salamanders of the genus *Plethodon* in western Oregon. Canadian J. Zool., 47:157-158.
- Rankin, J. S., Jr.** 1937. New helminths from North Carolina salamanders. J. Parasitol. 23:29-42.
- Reiber, R. J., E. E. Byrd, and M. W. Parker.** 1940. Certain new and already known nematodes from Amphibia and Reptilia. Lloydia 3:125-144
- Saxe, L. H.** 1955. Observations on *Eimeria* from *Ambystoma tigrinum*, with descriptions of four new species. Proc. Iowa Acad. Sci. 62:663-673.
- Trauth, S. E., H. W. Robison, and M. V. Plummer.** 2003. Amphibians and reptiles of Arkansas. Univ. Arkansas Press, Fayetteville. (In press)
- Upton, S. J., C. T. McAllister, and S. E. Trauth.** 1993. The coccidia (Apicomplexa: Eimeriidae) of Caudata (Amphibia), with descriptions of two new species from North America. Canadian J. Zool. 71:2410-2418.
- Waitz, J. A. and K. N. Mehra.** 1961. *Baerietta idahoensis* n. sp. a nematotaeniid cestode from the intestine of *Plethodon vandykei idahoensis* from northern Idaho. J. Parasitol. 47:806-808.
- Winter, D. A., W. M. Zawada, and A. A. Johnson.** 1986. Comparison of the symbiotic fauna of the family Plethodontidae in the Ouachita Mountains of western Arkansas. Proc. Arkansas Acad. Sci. 40:82-85.
- Woodhead, A. F.** 1928. *Haptophyra michiganensis* sp. nov., a protozoan parasite of the four-toed salamander. J. Parasitol., 14:177-182.