Journal of the Arkansas Academy of Science

Volume 47

Article 14

1993

Fishes of Bayou Meto and Wattensaw Bayou, Two Lowland Streams in East Central Arkansas

Walter Dean Heckathorn Jr. Arkansas Cooperative Fish and Wildlife Research Unit

Follow this and additional works at: https://scholarworks.uark.edu/jaas

Part of the Terrestrial and Aquatic Ecology Commons

Recommended Citation

Heckathorn, Walter Dean Jr. (1993) "Fishes of Bayou Meto and Wattensaw Bayou, Two Lowland Streams in East Central Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 47, Article 14. Available at: https://scholarworks.uark.edu/jaas/vol47/iss1/14

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

Journal of the Arkansas Academy of Science, Vol. 47 [1993], Art. 14 Fishes of Bayou Meto and Wattensaw Bayou, Two Lowland Streams in East Central Arkansas

> Walter Dean Heckathorn Jr. University of Arkansas Arkansas Cooperative Fish and Wildlife Research Unit Fayetteville, AR 72701

Abstract

Bayou Meto is a low-gradient, highly turbid, warm-water stream that originates in the foothills of the Interior Highlands of central Arkansas and flows southeastward 290 km to the Arkansas River. In the 1970's, Bayou Meto was contaminated with dioxins from a point source (Vertac Corp.) now recognized as a USEPA Superfund site. The present study was initiated to investigate the impact of dioxin on the fish community of Bayou Meto. Fishes were collected by backpack-electrofishing, boat-electrofishing, seines, hoopnets, minnow traps, and trot lines, at 14 sampling stations. Diversity indices (Shannon and Margalef) were used to compare diversity among sample sites. A total of 73 fish species was collected from Bayou Meto and Wattensaw Bayou (a reference stream) between May, 1991 and September, 1992. A total of 79 species had been reported from these drainages. I collected 64 species from Bayou Meto and 48 species from Wattensaw Bayou. Of the 79 species historically reported from these drainages, 17 were not collected during this study. However, of the 73 species collected, 11 (15% of the entire collection) had not been previously recorded from these drainages. There was 57% overlap in species between Bayou Meto and Wattensaw Bayou. Differences in collected species from the two drainages mostly involved rare species i.e., those species in low abundance according to the literature and/or difficult to collect. Centrarchids and castostomids dominated the fish communities of both streams. Percids were also well represented, but 50% were not previously reported from these drainages. Cyprinidae numbers were low and distributions spotty. Diversity varied among sites and was related to impacts and stream order. Diversity was highest at less impacted locations and downstream sites.

Introduction

Surveys of species are the cornerstone of ecological studies, providing basic data about what is present or absent in an ecosystem. Listing and recording the occurrence of species is indispensable to science, from the discovery of new species to affirmation of the anticipated. Species lists from geographical regions or from specific systems provide baseline data for future studies. Species and population baselines have become increasingly important to understanding the impacts of massive habitat alterations, pollution, introduction of non-native species, and recovery of endangered species.

The overall objective of this study was to describe the fish communities of Bayou Meto and Wattensaw Bayou for present and future reference. This study was part of a larger study funded by the U. S. Environmental Protection Agency and the U. S. Fish and Wildlife Service for the Arkansas Cooperative Fish and Wildlife Research Unit to investigate the impact of dioxins upon the aquatic community of Bayou Meto.

Bayou Meto is a low-gradient, highly turbid, warm-water stream that originates in the foothills of the Interior Highlands of central Arkansas (Fig. 1). Bayou Meto begins 38.8 km northeast of Jacksonville, Arkansas, and flow southeastward 290 km through the flat, fertile farmlands of the Mississippi Embayment to the Arkansas River near Reydell, Arkansas. The entire drainage area for Bayou Meto is 1,606 km² (Neely, 1985). Based upon 30 years of monitoring at the United States Geological Survey (USGS) gauging station at Lonoke, Arkansas, the average flow for Bayou Meto at Lonke is 8.2 m³/s; minimum and maximum flows were 0 and 133 m³/s, respectively (Neely, 1985).

Bayou Meto begins as an intermittent stream at an elevation of 500 m and becomes a second order stream 12.9 km above Jacksonville. The headwaters are characteristic of an upland stream with typical riffle and pool complexes. Bayou Meto becomes a third order stream 9.2 km above Jacksonville at its confluence with Bridge Creek, a fourth order stream at Bayou Meto Wildlife Refuge, and enters the Arkansas River as a fifth order stream.

A paucity of acquatic faunal studies in east-central Arkansas and the impacts of dioxins, sewage, and pesticides from industry, municipalities, and agriculture necessitated comparison of the fish fauna of Bayou Meto to a

less impacted stream to ascertain patterns of species alterations. Since 1970, dioxin-laden waste chemicals have been discharged from Vertac Chemical Corp. into Bayou Meto (ADPC&E, 1984, 1989). The Vertac site has been declared a Superfund Site by USEPA (USEPA, 1990, 1991).



Fig. 1. Sample stations on bayous Meto and Wattensaw. Number in circle indicates location and station. Station 5 was Lake Dupreee. Bayou Meto Wildlife Management area (WMA) is denoted by paraellel lines.

Wattensaw Bayou was used as a reference stream to Bayou Meto. Wattensaw Bayou begins 3 km southeast of Cabot, Arkansas, within 16 km of Bayou Meto, and flows eastward for 103.5 km before its confluence with the White River. These two streams are morphologically similar, flowing through the Mississippi Alluvial Plain of Arkansas. Both are meandering streams with large variances in seasonal flows. Deposition in both systems includes clay, sand, gravel, and detritus. The riparian zones of both streams are dominated by bald cypress (*Taxodium distichum*) and water tupelo (*Nyssa acquatica*). The fish communities of these lowland streams should be dominated by Centrarchidae (Lee et al., 1980; Robinson and Buchanan, 1988), a family of fishes particularly adapted to the slow moving and turbid waters found in bayous of east central Arkansas. The native fish species compositions of both Meto and Wattensaw bayous were hypothesized to be similar. However, there is no known point source of effluent flowing into Wattensaw Bayou. Except for agricultural practices that dominate the lower reaches of both rivers, Wattensaw Bayou is not known to be highly polluted and, therefore, useful as a control to Bayou Meto.

Physicochemical parameters were analyzed in an seperate study and these parameters followed patterns in downstream gradients as predicted by Vannote et al. (1980) and others (Heckathorn, 1993). Assumptions made of these two drainages were that all sampling stations should produce relatively similar species, richness, and diversity with trends toward increasing numbers downstream and differences among sampling stations within Bayou Meto and between bayous Meto and Wattensaw should correlate to point source inputs. The above assumptions are based upon Heckathorn (1993) and personal observations.

Materials and Methods

In May, 1991, 12 sampling stations were designated on Bayou Meto and two sampling stations on Wattensaw Bayou (Fig. 1; Table 1). Fish collection methods included backpack-electrofishing, boat-electrofishing using a 240 V, 3500 W DC electrofishing boat adjusted to five amperes, seining, hoop nets, minnow traps, and trot lines (Nielson and Johnson, 1985). Collection sites were sampled during daylight hours intermittently from May through October, 1991. Seining proved impractical at most stations because of extensive vegetation, including both dead and living trees. At the end of the 1991 field season, sampling sties were confirmed (Table 1) and a sampling regime was designed and developed for 1992. Intensive sampling occurred in May and September of 1992 at all stations. Each sampling station was measured at 75-100 m of stream length (Owens and Karr, 1978; Schlosser, 1982). The sampling regime consisted of three 20-min. boat-electrofishing runs, accompanied by one set of two 24-h 576 mm diameter hoopnet (1 baited and 1 unbaited) and one set of eight 24-h unbaited minnow trap. Both banks of each station were boat-electrofished using a 240 V, 3500 W DC electrofishing boat adjusted to five amperes, with the same pattern of 100 m downstream and then 100 m upstream of the opposite bank being followed on all three runs. All electrofishing was done during daylight hours. Immediately after electrofishing hoopnets and minnow traps were set. Hoopnets were placed with the unbaited

Proceedings Arkansas Academy of Science, Vol. 47, 1993

Table 1. Sampling stations with locations, counties, legal descriptions, distance from the river source, and stream order. BM is bayou Meto, WB is Wattensaw Bayou, and WMA is Wattensaw Wildlife Management Area. The first of two Sewage Treatment Plants discharges into Bayou Meto above Station 3, and the second plant discharges above Station 4. Rocky Branch confluence with Bayou Meto is between stations 3 and 4.

	River		Legal						
Stations	Locations	Counties		Descriptio	Km	Order			
BM-1	Macon Br.	Pulaski	3N,	R11W,	Sect. 5	26.9	2nd		
BM-2	Cato Br.	Pulaski	T3N,	R11W,	Sect. 22	35.3	3rd		
BM-4	Reeds Br.	Pulaski	T3N,	R11W,	Sect. 31	41.9	3rd		
BM-6	Broken Br.	Pulaski	T3N,	R10W,	Sect. 33	44.9	3rd		
BM-7	I-40 Br.	Pulaski	T2N,	R10W,	Sect. 14	55.8	3rd		
BM-8	HWY 15 Br.	Lonoke	T2N,	R9W,	Sect. 29	61.4	3rd		
BM-11	HWY 13 Br.	Lonoke	T1N,	R8W,	Sect. 10	107.4	3rd		
BM-12	HWY 165 Br.	Lonoke	T2S,	R6W,	Sect. 7	133.6	3rd		
BM-13	HWY 79 Br.	Arkansas	T3S,	R6W,	Sect. 11	166.3	3rd		
BM-14	HWY 152 Br.	Arkansas	T3S,	R6W,	Sect. 33	182.0	4th		
BM-15	Cox Cypress	Arkansas	T5S,	R6W,	Sect. 1	218.7	4th		
BM-16	HWY 11 Br.	Arkansas	T6S,	R5W,	Sect. 3	253.9	4th		
WB-18	HWY 13 Br.	Prairie	T3N,	R9W,	Sect. 12	60.5	2nd		
WB-20	WMA	Prairie	T3N,	R6W,	Sect. 22	90.7	4th		

NOTE: Only Cox Cypress (BM-15) was deleted from the 1992 sampling regime. Also, stations 3, 5, 9, 10, 17, and 19 were used in the sediment and invertebrate part of the overall dioxin contamination study.

net immediatly downstream of each 100 m sampling station and with the baited net immediately upstream. Minnow traps were placed randomly throughout the sampling station. Block nets were not used due to possible high levels of dioxin contamination. The sampling was repeated again in September, 1992, but hoopnets and minnow traps were not set due to poor catches during spring sampling. Fish species lists for both Bayou Meto and Wattensaw Bayou were compiled from data collected from the 1991 random sampling and the 1992 sampling regime.

Most collected fishes were fixed in 10% formalin, washed in water, preserved in 50% isopropyl alcohol, and returned to the lab for identification. Large fish were identified in the field and returned unharmed. Specimens were eventually changed to 70% ethanol and placed into the University of Arkansas Natural History Museum in Fayetteville, Arkansas. The Common and Scientific Names of Fishes published by the American Fishery Society was used to name fish species collected (Robins et al., 1991). Species identifications were confirmed in the lab by Dr. James E. Johnson, U.S. Fish and Wildlife Service.

Electrofishing, hoopnets, and minnow traps catch per unit effort (CPUE) in the 1992 sampling regime was consistent among sites and allowed for comparisons among sampling stations using diversity indices. Only fishes collected in the May and September sampling regime were used to determine and compare diversity and richness. Two diversity indices (Shannon and Margalef) were chosen for use in this study (Shannon, 1958; Margalef, 1958, 1963; Washington, 1984; Boyle et al., 1990). Diversity indices elucidate differences among sites and show trends in declines or increases in diversity. Boyle et al. (1990), showed that best results are obtained by using indices in concert, thus allowing for complementation and dampening of deficiencies.

Shannon and Margalef indices were chosen based upon wide use, acceptance in aquatic studies, and distinction in responses to variances in populations (Washington, 1984; Boyle et al., 1990).

Results

Sampling efficiency was subjectively assessed as excellent to good for the 1992 electrofishing effort. The first and second of the set of three electrofishing runs collected 81% and 91% of the species, respectively, and numbers of fish collected declined with each successive run. These data are judged adequate to make inferences about species presence or absence, richness (total number of species), and diversity.

A total of 73 fish species was collected from bayous Meto and Wattensaw between May, 1991, and September, 1992 (Table 2). Seventy-nine species had been reported from the regional area of these drainages (Lee et al., 1980; Robison and Buchanan, 1988). Sixty-four fish species were collected from Bayou Meto and 48 species from Wattensaw Bayou. Fishes collected in only one of these two drainages are shown in Table 3. Of the 79 species historically reported from these drainages, 17 (22%) were not found during this survey. Conversely, of the 73 species collected, 11 (15%) had not been previously recorded from these drainages. There was a 53% overlap in species between Bayou Meto and Wattensaw Bayou.

Centrarchids and catostomids dominated the fish communities of both streams. Polyodon spathula and Anguilla rostrata were both collected only from Wattensaw Bayou, and only in low numbers. Gars were represented in both streams at all stations by Lepisosteus oculatus, but L. platostomas was collected only in Bayou Meto and L. osseus only in Wattensaw Bayou. Amia calva was well represented in both systems. Cyprinids were represented by shiners and minnows, but their species numbers were notably low and their distributions spotty and variable. Shiners in Wattensaw had a higher diversity than in Bayou Meto. Ictalurids were represented in both streams; Ictalurus punchtatus was most abundant. Esocids were represented in both systems by Esox americanus. Aphredoderus sayanus was abundant in Bayou Meto but absent from Wattensaw Bayou. The family Cyprinodontidae was well depicted by Fundulus olivaceus in both drainages, but F. notatus was taken only from Bayou Meto, and then only in low numbers. Gambusia affinis, and at least one of two species of Atherinidae, were collected at most stations in both streams. The family Percicthyidae was represented by Morone mississippiensis in low numbers in both bayous, and M. saxatilis was collected once at mid-stream in Bayou Meto. Percids were well represented, considering their lack of susceptibility to electrofishing (Vibert, 1967, Novotny and Priegel, 1974). Percidae also had the highest percentage of unreported species, with 50% as newly described from these drainages in this work. Aplodinotus grunniens was collected at all sampling stations except Station 4 on Bayou Meto. Some of the highest dioxin contamination concentrations in Bayou Meto were reported from Station 4 (ITC, 1987, USEPA, 1990, 1991).

Species diversity varied by station (Fig. 2). However, relative diversity trends did not vary greatly between indices nor between May and September sampling dates, and congruence between these parameters ameliorates validation of sample methodologies and these data. The highest diversities from Bayou Meto were consistently found at stations 1, 2, and 12, with the lowest diversities at stations 4 and 6, where dioxin levels and sewage effluent were highest (ITC, 1987, USEPA, 1990, 1991) (Fig. 2). Sewage effluent negatively impacts fish communities immediately downstream, but further downstream beyond the septic and/or toxic zone, benefits may be derived by certain fish species due to increases in flow and nutrients (Owens and Karr, 1978). Diversity in Bayou Meto remained unpredictable low from stations 4 through 12. At stations 14 and 16, increases in width, depth, and turbidity of the stream may have reduced sampling efficiency, resulting in an underestimate of diversity and richness (Schlosser 1982; Heckathorn, 1993). Diversity in Wattensaw Bayou was relatively low at the upper station and highest at the most downstream station. Station 20 at Wattensaw had the highest diversity and richness of any station (Figs. 2 and 3).

Richness proved variable among sampling dates. Total numbers of species in May fluctuated less and averaged higher than in September. Numbers of species for both sampling dates followed the same trend as seen in the diversity indices, except that May results were not as conspicuously low. Contrasts between samples taken in May and September, were most likely due to differences in flow. For example, in September, water flows dropped to levels which prevented the sampling of Station 12. Fishes might have emmigrated out of my sampling stations to deeper waters. Lower water levels should also concentrate fishes, making them more susceptable to collecting gear. In general diversity and richness decreased at stations with decreases in water levels, but realative diversity and richness remained consistent among stations in between May and September sampling dates.



Diversity indices for Bayou Meto and Wattensaw Bayou



Diversity indices for Bayou Meto and Wattensaw Bayou

Fig. 2. Diversity by stations for May and September sampling dates. Stations 1 and 2 are upstream of dioxin and sewage impacts. Stations 18 and 20 are at unimpacted Wattensaw Bayou. We were unable to sample station 12 in September due to low flow.



Richness of Bayou Meto and Wattensaw Bayou

Fig. 3. Richness by stations for May and September sampling dates. Stations 1 and 2 are upstream of dioxins and sewage impacts. Stations 18 and 20 are at Wattensaw Bayou. Richness should increase downstream as seen from unimpacted stations 1-2 and 18-20. We were unable to sample site 12 in September.

Discussion

Differences in species collected from the two drainages mostly involved rare species (those collected in low numbers or at only one station). It is likely that these species were present in low numbers in both streams. However, 15 species were collected in substantial numbers in one drainage but not the other: Lepisosteus platostomas, Cyprinus carpio, Lythrurus umbratilis, Carpoides carpio, Ictiobus niger, Ictalurus furcatus, Aphredoderus sayanus, Fundulus notatus, Centrarchus macropterus, Micropterus punctulatus, Etheostoma chlorosomum, E. spectabile, and Percina maculata came only from Bayou Meto, and Lepisosteus osseus and Notropis texanus came only from Wattensaw Bayou (Table 3).

Many of the 17 species reported from bayou Meto and Wattensaw, but not collected during this study, are not susceptible to electrofishing, and other species, if present, would be expected in low numbers, reducing their chances of capture (Table 3). For example, Polyodon spathula was collected from Wattensaw Bayou, but only by chance. During a water quality sampling effort, not during fish collection efforts I chanced upon a fisherman, whom was catching young P. spathula on minnows. Fish like P. spathula, Scaphirhynchus platorynchus, Cycleptus elongatus, and Percina shumardi may still reside in the two streams, but are not susceptible to capture (Miller and Robison, 1973; Robison and Buchanan, 1988). Fish like Alosa chrysochloris, Hiodon alosoides, H. tergisus, Hypophthalmichthys molitrix, H. nobilis, and Erimyzon sucetta should have been present, but low numbers likely reduced chance of capture (Beckett and Pennington, 1986; Robison and Buchanan, 1988). Notropis blennius, N. venustus, N. volucellus, Pimephales notatus, Fundulus chrysotus, F. dispar, Morone chrysops, and Lepomis marginatus were expected in abundant numbers and are also susceptable to electrofishing as evidenced by the consistent capture of similar species from the same families of the above species, and their absence may indicate recent losses from these drainages (Robison and Buchanan, 1988) (Table 3). Results of Shannon and Margalef diversity indices were similar, and indicate a real change in diversity among stations. Species diversity should be maximized at mid-reach of rivers (fourth to sixth stream order) where microhabitats are most abundant. However, as expected the fish community of Bayou Meto did not recapitulate the River Continuum Concept (RCC), possibly due to contaminants from dioxins, pesticides, and sewage discharge (Platts, 1979; Vannote et al., 1980; Minshall et al., 1983; Schlosser, 1982).

In May, diversity and richness increased as expected between the first two stations at Bayou Meto, and in both May and September at Wattensaw, in accordance with the RCC. Diversity decreased at Station 4 (below dioxin and sewage effluent) and continued at a reduced level to the Table 2. Fish species composition of Bayou Meto and Wattensaw Bayou, for May, 1991, through September, 1992 (without CPUE for stations). Numbers under stations are the number of individual fish collected for all sample dates. N is the total number of individuals of a species collected from both drainages, and totals are total number of fishes collected at a sample station. Total collection included 7,465 fishes.

TAXA					STATIONS										
Family	A	bove											Wat	tensaw	
Species	v	ertac											Bayou		
	1	2	4	6	7	8	11	12	13	14	15	16	18	20	N
Polyodontidae															
Polyodon spathula														3	3
Lepisosteidae															
Lepisosteus oculatus	5	13	14	46	20	22	7	16	33	35	8	19	8	- 33	279
Lepisosteus osseus													4	1	5
Lepisosteus platostomas					1		4	4	9	6	4				28
Amiidae			-												
Amia calva	9	10	1	9	4	4	1	4	6	5		4		9	66
Anguillidae															
Anguilla rostrata														1	1
Clupeidae	00	00	10	0	10	10	10				00.		10	00	914
Dorosoma cepedianum	22	92	13	8	19	16	18	3	38	11	20a	15	13	20	314
Dorosoma petenense												25		55	18
Cyprinidae															
Campostoma anomatum	3													1	4
Cienopharyngoaen taetta	91	94		e	95	19	10	c			10	6	0		161
Cyprinus carpio	21	24	5	0	50	15	10	0	5	3	10	0	9	40	41
Hybognathus nuyt				1					1		1			30	49
Intervente formare	67			14	4	9	6	11	16		8	7		9	187
I sthrurus umbratilis	58	1	9	11	1	•	v		10					· · ·	57
Maerhybabsis storeriana	00		~		· ·									1	1
Notemigonus crysoleucas	2	5			2				1		1			3	14
Notropis amis	-	0			-	1						1		16	18
Notropis atherinoides					1	•		1	1	2		ī	2	13	21
Notropis buchanani					-									4	4
Notropis lutrensis											3				3
Notropis maculatus	4	1						1		1		2	3	14	26
Notropis shumardi														4	4
Notropis texanus														31	31
Opsopoeodus emiliae	26	18	2	8	4	7	1		5	6	4	1	8	13	103
Pimephales vigilax											1				1
Catostomidae															0
Carpiodes carpio					2	9		1	1	10	1	13			37
Erimyzon oblongus														1	1
Ictiobus bubalus	1	34	7		5	58		3	1	13	21c	9	7	5	164
Ictiobus cyprinellus	14	35	12	30	18	37	2	1	6	7	59b	3		6	230
Ictiobus niger		3	4		2	5	1				10				25
Minytrema melanops	18	11			2									5	36
Moxostoma macrolepidotum				1											1
Ictaluridae															0
Ameiurus melas		~											1		1
Ameiurus natalis	1	1								2	1.2	147		2	4
Ictalurus furcatus										2	3	5			10
Ictalurus punctatus	2	4	4	3	3	2	1	2	11	6	1	24		8	69
Noturus gyrinus	1	1													2
Fylodictis olivaris					2				1						3
Esocidae		4													10
Esox americanus	11	1	1								1			2	16
Aphredoderidae	E.	6													10
Aphreaoaerus sayanus	5	0						1							12
Evendulus notation	1	9							e						11
Fundulus notatus	84	15	8	1	1	8	- 1	9	0	1	1		11	96	159
a terratestas desducetas	0.1	10			2. 						3 4 0	9	**	AU	104

Proceedings Arkansas Academy of Science, Vol. 47, 1993

Table 2. Continued:

TAXA										STAT	IONS				
Family		Above											Wat	tensaw	
Species	i	/ertac											1	Bayou	
opecies	1	2	4	6	7	8	11	12	13	14	15	16	18	20	N
Poeciliidae		1													
Gambusia affinis	16	8	1	6	4	43	3	2	2		93	19	1	15	213
Atherinidae													~		0
Labidesthes sicculus	21	24	3	5	2	1			4	4	2		1	21	88
Menidia bervllina	2		4		-									5	11
Percicthvidae														- T	0
Morone mississibhiensis												4		3	7
Morone savatilis								1						<u></u>	1
Centrarachidae								•							ō
Centrarchus macrohterus	6	9													8
Elassoma zonatum	0	2													2
Labomis overallus	11	ŝ	5	9	1		8	8	8	9	8	1	10	1	53
Lepomis cyanettas	81	80	21	69	09	88	94	95	95	16	98	6	88	23	444
Lepomis guiosus Lebomic hymilic	9	14	7	20	45	46	95	50	88	119	38	51	89	2	548
Lepomis macrochirus	145	119	118	187	191	109	79	58	104	160	88	48	155	167	1.640
Labornie magglotie	07	109	190	189	44	19	95	99	179	78	19	50	179	89	1 180
Lepomis microlophus	15	4	8	9	7	0	8	1		8	15	50	19	9	78
Lepomis microtophus	8	6	10	•		4	16	10	16	18	6	8	84	6	140
Lepomis punctatus	•	9	10		~		10	10	10	10		9		×.	9
Microphenus bunchulatus		~	9	90					1	9	9	8			87
Micropherus salmaides	944	10	18	10	5	7	7	6	7	8	740	3	17	45	236
Pomovis appularis	19	94	9	19	90	19	94	16	26	14	44	16	6	2	253
Pomovis nioromaculatus	8	4	1	3	1	3	4	3	19	9	9	4	6	10	58
Percidae	5			5			1	5	14	-	-		U	10	0
Ammocrobia vivar										1					ĭ
Etheostoma astriaene											8			1	Â
Etheostoma chlorosomum	1	8						1	1		33			· ·	39
Etheostoma gracile		ĭ							•		00				1
Etheostoma proeliare	1	î									4		1	2	ĝ
Etheostoma spectabile	5	÷.												-	5
Etheostoma stiamaeum	1														1
Etheostoma shiphlei	î														î
Percing cabrodes								1	8		8			10	17
Parsing magulata	9		1					î		1					5
Fercina macutata	4									*					0
Ablodinotus amunnians	1	9		2	94	14	91	6	51	11	1	95	9	8	169
Aptountorus grunniens	1	4		5	24	14	21	0	51	11	1	20	-	0	105
TOTALS	754	632	386	651	504	479	293	262	659	549	595	370	548	783	7,465

a = + 263 fry.

b = 50 juveniles between 28/mm - 68/mm.

c = 18 juveniles between 28/mm - 68/mm.

d = 13 were fry. e = 71 juveniles between 33/mm - 55/mm.

Table 3. Comparison of fish species collected in 1991 and 1992 from bayous Meto and Wattensaw (without CPUE for the two drainages). Species collected in Bayou Meto and/or in Watensaw Bayou are denoted by X. Species historically reported from these drainages but not collected in this study are listed and followed by an asterisk but not marked. Rare species are those collected only once from a station or so designated in the literature, and are denoted by an X. Species difficult to capture by my sample methods are so designated; and species not recorded from these drainages but collected are designated by two asterisks.

Species	Bayou Meto	Wattensaw Bayou	Rare species	Difficult to collect
Scaphirhynchus [;atprumcjis			x	x
Polyodon spathula		x		x
Lepisosteus osseus		x		
Lepisosteus platostomus	x			
Anguilla rostrata		x	x	
Alosa chrysochloris *			x	x
Cambostoma anomalum **	x	x		
Ctenoparyngoden idella	x	~		
Cybrinus carbio	Ŷ			
Higdon alosoides *	~		x	
Hindon terrisus *			Ŷ	
Hubomathus havi **	×	v	a	
Hybophanias nayi	~	~	v	
Hypophinachichthys motific			\$	
Typophinatmichinys noonis	v		~	
Marchikatorie starmine	^	v	v	
Macryoopsis storeriana		х	x	
Notropis blennius -				
Notropis buchanani		x		
Notropis lutrensis	x	and the second se	X	
Notropis shumardi **		x	х	
Notropis texanus **		x		
Notropis venusthus *				
Notropis volucellus *				
Pimephales notatus *				
Pimephales vigilax	x		x	
Carpoides carpio	x			
Cycleptus elongatus *				
Erimyzon oblongus **		x	x	
Erimyzon sucetta		x	x	
Ictiobus niger	x			
Moxostoma macrolepidotum	x		x	
Ameiurus melas		x	x	
Ictalurus furcatus	x		x	
Noturus gyrinus	x			
Pylodictis olivaris	x			
Aphredoderus sayanus	X			
Fundulus chrysotus *				
Fundulus dispar *				
Fundulus notatus	X			
Morone chrysops *	2552			
Morone saxatilis	x		x	
Centrarchus macroblerus	x			
Elassoma zonatum	x		x	
Lehomis marginatus	•		~	
Lehomis symmetricus	x		x	
Microhterus hunchilatur	Ŷ		^	
Ammocratha vivas **	Ŷ		v	v
Fitherstown a blowstown	÷		^	Ŷ
Etheostoma chiorosomum	Ň		v	A V
Etheostoma gracite	X		X	X
Etheostoma spectabile **	X		X	X
Etheostoma stigmaeum **	X		X	X
Etheostoma whipplei **	X		x	X
Percina maculata **	х			x
				v

Proceedings Arkansas Academy of Science, Vol. 47, 1993

last station. Declines along a gradient in which diversity and richness should increase are indicative of disturbance (Diamond and Gilipin, 1980; Angermeir and Schlosser, 1989). The fish community of Bayou Meto appears to be impacted at Station 4, and community structure remained variable and unpredictable through the last sample station.

The 1992 sampling regime showed Wattensaw Bayou to have higher diversity and richness when compared to Bayou Meto. The highest diversity and most species collected for all sample dates was at Station 20 on Wattensaw Bayou. However, when examining total fish species collected for 1991 (without CPUE) and 1992 Bayou Meto has more species than Wattensaw Bayou. Bayous Meto and Wattensaw are both well represented by the total number of species. Difference in the two streams may be due to differential sampling efforts. Most likely, the total number of species in Wattensaw would have exceeded Bayou Meto if a similar sampling effort had been used (same number of sampling stations and sampling days). For example, absence of *C. carpio* from Wattensaw likely was an artifact of this differential sampling.

Bayou Meto is an impacted stream below the confluence of Rocky Branch. Dioxins and/or pesticide contamination appear to be the most likely reasons for declines in fish diversity in Bayou Meto, although sewage effluent and pesticides could also contribute significantly to these declines as a singular agent or in synergism. Fish taxa were missing from this survey, and loss of species due to the above anthropomorphic alterations is possible, and is deserving of future investigation. Wattensaw Bayou and stations 1 and 2 on Bayou Meto proved high in fish diversity and followed the RCC model.

Acknowledgements

This research has been supported by grants from the U.S. Fish and Wildlife Service and the U.S. Environmental Protection Agency, with technical support by the Arkansas Cooperative Fish and Wildlife Research Unit, the Arkansas Game and Fish Commission, and University of Arkansas at Fayetteville. Helpful manuscript suggestions were contributed by Dr. James E. Johnson, Dr. Arthur V. Brown, and Dr. William G. Layher; and invaluable field collection assistance came from Dr. James E. Johnson, Dr. William G. Layher and his field assistants at U of A Pine Bluff, Andrew L. Thompson, Steve Marchant, and Roseanne Barnhill. Thanks to my lovely wife who helped whenever possible.

Literature Cited

- ADPC&E. 1984. Arkansas Department of Pollution Control and Ecology. Monitoring Rocky Branch at East Main Street: Ref: Permit #2713 WR-1 memorandum to water files from Frank Stephens.
- ADPC&E. 1989. Monitoring Rocky Branch at East Main Street: Ref: Permit #2713 WR-1 Memorandum to Water files.
- Angermeir, P.L. and I.J. Schlosser. 1989. Species-area relationships for stream fishes. Ecology 70:1450-1462.
- Beckett, D.C. and C.H. Pennington. 1986. Water quality, macroinvertebrates, larval fishes, and fishes of the lower Mississippi River: a synthesis. Tech. Rept. E-86-12. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
- Boyle, T.P., G.M. Smillie, J.C. Anderson and D.R. Beeson. 1990. A sensitivity analysis of nine diversity and seven similarity indices. Res. J. WPCF 6:749-762.
- Diamond, J. and M.E. Gilipin. 1980. Turnover noise: contribution to variance in species number and predictions from immigration and extinction curves. Am. Nat. 115:884-889.
- Heckathorn, W.D. Jr. 1993. 2, 3, 7, 8-Tetrachlorodibenzop-dioxin in Bayou Meto and the ecology of a contaminated stream in east central Arkansas. Master Thesis. University of Arkansas.
- ITC. 1987. Sampling and analytical program plan Vertac offsite areas. International Technology Corporation. Knoxville, Tennessee. 85 pp.
- Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. Mcallister and J.R. Stauffer, Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Mus. Nat. Hist., Raleigh, North Carolina. 867 pp.
- Margalef, R. 1963. On certain unifying principles in ecology. Am. Natur. 97:357-374.
- Margalef, R. 1958. Information theory in ecology. Gen. systems 3:36-71.
- Miller, R.G. and H.W. Robison. 1973. The fishes of Oklahoma. Oklahoma State University Museum of Natural and Cultural History. Series No. 1. 246 pp.
- Minshall, G.W., R.C. Peterson, K.W. Cummins, T.L. Bott, J.R. Sedell, C.E. Cushing and R.L. Vannote. 1983. Interbiome comparison of stream ecosystem dynamics. Ecol. Monogr. 53:1-25.
- Neely, B.L. 1985. Water-surface profiles along Bayou Meto and Rocky Branch near Jacksonville, Arkansas: U.S. Geological Survey Openfile Report 85-176. 12 pp.
- Nielson, L.A. and D.L. Johnson. 1985. Fisheries techniques. American Fisheries Society Bethesda, Maryland. 468 pp.
- Novotny, D.W. and G.R. Priegel. 1974. Electrofishing boats: improved designs and operational guidelines to increase the effectiveness of boom shockers.

Wisconsin Dep. Nat. Res. Tech. Bull. 73. 48 pp.

- Owens, T.G. and J.R. Karr. 1978. Habitat structure and stream fish communities. Ecology 59:507-515.
- Platts, W.S. 1979. Relationships among stream order, fish populations, and aquatic geomorphology in an Idaho river drainage. Fisheries 4:5-9.
- Robins, R.C., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea and W.B. Scott. 1991. Common names of fishes from the United States and Canada. Fifth Edition. Am. Fisheries Soc. Spec. Pub. 20. 183 pp.
- Robison, H.W. and T.M. Buchanan. 1988. Fishes of Arkansas. The University of Arkansas Press. 563 pp.
- Schlosser, I.J. 1982. Fish community structure and function along two habitat gradients in a headwater stream. Ecol. Monogr. 52:395-414.
- Shannon, C.E. 1948. A mathematical theory of communication. Bell System Tech. J. 27:379-423, 623-656.
- **USEPA.** 1990. Quarterly status report of superfund sites in region 6, state of Arkansas. 7 pp.
- USEPA. 1991. Vertac site update: An update on activities at the Vertac Superfund site. EPA Region 6. Haz. Waste Div. 9 pp.
- Vannote, R.L., G.W. Minshall, K.W. Cummins, J.R. Sedell and C.E. Cushing. 1980. The river continuum concept. Can. J. Fish. Aquat. Sci. 37:130-137.
- Vibert, R. 1967. Fishing with electricity its applications to biology and management. Fishing News (Books) Ltd., London. 276 pp.
- Washington, H.G. 1984. Diversity, biotic and similarity indices a review with special relevance to aquatic ecosystems. Water Resour. 18:653-694.

and the second se

M. P. M. Martin, "Annual Sciences and Destination of the Action of th

Proceedings Arkansas Academy of Science, Vol. 47, 1993