

1993

## Microscale Preparation of Pyrocatechol: The Use of Sodium Percarbonate in the Dakin Reaction

T. Evans Yamazaki

*University of Arkansas at Little Rock*

Dominic T.C. Yang

*University of Arkansas at Little Rock*

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

 Part of the [Organic Chemistry Commons](#)

---

### Recommended Citation

Yamazaki, T. Evans and Yang, Dominic T.C. (1993) "Microscale Preparation of Pyrocatechol: The Use of Sodium Percarbonate in the Dakin Reaction," *Journal of the Arkansas Academy of Science*: Vol. 47 , Article 37.

Available at: <https://scholarworks.uark.edu/jaas/vol47/iss1/37>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in *Journal of the Arkansas Academy of Science* by an authorized editor of ScholarWorks@UARK. For more information, please contact [scholar@uark.edu](mailto:scholar@uark.edu).

## Microscale Preparation of Pyrocatechol: The Use of Sodium Percarbonate in the Dakin Reaction

T. Evans, F. Yamazaki, and D.T.C. Yang\*

Department of Chemistry  
University of Arkansas at Little Rock  
Little Rock, Arkansas 72204

\*To Whom correspondence should be addressed.

Sodium percarbonate ( $\text{Na}_2\text{CO}_3 \cdot 3/2\text{H}_2\text{O}_2$ ), a common household detergent, is an inexpensive, stable, and easily handled reagent that has an excellent shelf life. One equivalent of sodium percarbonate releases one and one-half equivalents of hydrogen peroxide. Sodium percarbonate (Fig. 1) has been used for the oxidation of sulfides (Ando, *Chem. Lett.*, 665, 1986), amines (Zazac, *J. Org. Chem.*, 54:2468, 1988), and organoboranes (Kabalka, *Organometallics*, 9:1316, 1990). We wish to report the use of this reagent for the preparation of pyrocatechol from salicylaldehyde (Yamazaki, M.S. Thesis, University of Arkansas at Little Rock, 1991), a Baeyer-Villiger type reaction commonly referred to as the Dakin reaction (Dakin, *Org. Synth.*, Coll. Vol. 1: 149-150, 1964).

hydroxyacetophenone in 69% yield. In closely related experiments, hydroquinone was prepared from 4-hydroxybenzaldehyde in 35% yield as well as from 4-hydroxyacetophenone in 66% yield (Evans, M.S. Thesis, University of Arkansas at Little Rock, 1991). In all cases the ir and nmr spectra of the products were identical to those of authentic samples.

### Acknowledgements

We thank UALR Cooperative Research Fund for partial support of the work.




Fig. 1. Sodium percarbonate

To a solution of salicylaldehyde (244 mg, 2 mmol) in 3 ml of 1 N sodium hydroxide was added gradually sodium percarbonate (630 mg, 6 mmol). A vigorous exothermic reaction occurred immediately with darkening of the mixture which was then stirred for 24 hours in a sandbath at 50°C. After chilling the reaction mixture in an ice-bath and neutralization with acetic acid, the solvent was completely removed under reduced pressure leaving a dry black residue. The residue was finely crushed and extracted with toluene overnight using a microscale Soxhlet extraction apparatus. The toluene solution was evaporated to yield crude pyrocatechol which was further recrystallized from toluene to afford 0.11 g of pure compound in 50% yield, mp 102-104°C (Lit. 105°C).

Similarly, pyrocatechol was also obtained from 2-