2016

An Adult Female Mosquito Survey in Southwest Missouri in 2014

O. E. Jamieson
oejamies@uark.edu

J. L. Moreland
Crowder College - Cassville Campus

D. H. Jamieson
Crowder College - Cassville Campus

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Entomology Commons](#)

Recommended Citation

Jamieson, O. E.; Moreland, J. L.; and Jamieson, D. H. (2016) "An Adult Female Mosquito Survey in Southwest Missouri in 2014," *Journal of the Arkansas Academy of Science*: Vol. 70 , Article 45.

DOI: <https://doi.org/10.54119/jaas.2016.7037>

Available at: <https://scholarworks.uark.edu/jaas/vol70/iss1/45>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

An Adult Female Mosquito Survey in Southwest Missouri in 2014

O.E. Jamieson^{1*}, J.L. Moreland², and D.H. Jamieson²

¹1259 W. Mount Comfort Road, Apt. 203, Fayetteville, AR 72701

²Department of Biological Sciences, Crowder College - Cassville Campus, 4020 North Main Street, Cassville, MO 65625

*Correspondence: oejamies@uark.edu

Running Title: An Adult Female Mosquito Survey in Southwest Missouri in 2014

In the fall of 2008 and 2009, 35 students enrolled in freshman biology courses at NorthWest Arkansas Community College participated in a landing/probing mosquito survey to determine if the Asian tiger mosquito, *Aedes albopictus* (Skuse), was the principal pest mosquito in Northwest Arkansas (Wilson and Jamieson 2010). A total of 110 mosquitoes representing four genera and seven species was collected. *A. albopictus* represented 79.1% (87 of 110) of mosquitoes collected during the study. The West Nile Fever vector *Culex pipiens* (Linnaeus) was a distant second at 6.4%.

We conducted a similar survey in the summer of 2014 in Southwest Missouri. Sampling followed the methodology of Pfizner et al. (1998), where students sat in a shaded area on their property for 20 minutes and collected any adult female mosquito attempting to take blood using a wide-mouthed vial. Students were instructed to capture the mosquito while it was probing and before it actually started taking a blood meal. All collections were done within the 2 hour period before dusk with the intent of maximizing the chances of capturing diurnal, crepuscular, and nocturnal species. Any mosquito captured was killed by being placed in a freezer overnight and subsequently identified using the keys of Darsie and Ward (2005). There were 4 collection sites, one each within the city limits of Cassville (36.6800° N, 93.8694° W), Washburn (36.5894° N, 93.9639° W), Monett (36.9218° N, 93.9259° W) and Crane (36.9039° N, 93.5711° W) Missouri. Six students made a total of 42 collections beginning on June 4th and concluding on September 25th. A total of 216 specimens was collected represented by 11 species and 5 genera (Table 1). The greater numbers and diversity in Southwest Missouri is probably explained by the fact that the collection sites were located in more rural areas than in the Northwest Arkansas survey. The rural environment offers a wider array of larval development sites than does the suburban environment sampled in the Arkansas survey. The primary oviposition sites offered to mosquitoes in

the suburban area are artificial containers such as discarded automobile tires, flower pots, house gutters and other habitats that resemble the tree holes which *A. albopictus* originally utilized as a larval habitat in its native region of Southeast Asia (Hawley 1988, Moore et al. 1988). Regardless, the Asian tiger mosquito dominated both surveys.

Table 1. Species Survey of Landing/Probing Mosquitoes in Southwest Missouri, 2014.

Species	Percentage of Total
<i>Aedes albopictus</i>	44.9
<i>A. trivittatus</i>	17.6
<i>Culex erraticus</i>	11.1
<i>A. vexans</i>	9.3
<i>C. pipiens</i>	8.8
<i>C. restuans</i>	2.8
<i>Psorophora ferox</i>	1.9
<i>O. canadensis</i>	1.4
<i>A. triseriatus</i>	1.4
<i>Anopheles quadrimaculatus</i>	0.5
<i>P. ciliata</i>	0.5

Aedes albopictus has colonized virtually all cities within the Ozark Mountains Physiographic Region and in many areas is locally the only pestiferous species (Jamieson and Olson 1995, Pfizner et al. 1998). In addition, its presence and abundance is of major concern because of its ability to vector diseases such as dengue fever and chikungunya (Miller and Loaiza 2015). The concern regarding Zika virus affecting travelers returning to the United States has made the monitoring of *A. albopictus* populations even more important. The Centers for Disease Control lists both

Aedes aegypti (Linnaeus) and *A. albopictus* as the principal vectors of this potentially devastating arbovirus (CDC – Zika 2016). *A. albopictus* is easily distinguished from native species by its small size, distinctive black and white markings and stripe on the top of the thorax (scutum) (Figure 1).

Fig.1: Distinctive markings on *Aedes albopictus*

Culex pipiens (Linnaeus) was the second most abundant mosquito collected in the Northwest Arkansas study although it only represented 6.4% of the total number of specimens collected. Wilson and Jamieson (2010) mentioned its presence as significant because of its ability to vector West Nile fever (Kilpatrick et al. 2005). In this study it was slightly more abundant (8.8% of total specimens collected) but was surpassed by another *Culex* species, *Culex erraticus* (Dyar and Knab) at 11.1%. Both are competent West Nile potential vectors because of their preference for avian blood (Bolling et al. 2005). It is interesting to note that the third author has conducted surveys, both larval and adult, across the Ozarks for the last 25 years and has never encountered *C. erraticus*. The larval production site was never found in this study, but the third author has collected *C. erraticus* in rice fields in Eastern Arkansas (Jamieson et al. 1994).

It is hoped that these data will be of value to vector control specialists with the Missouri Health Department.

Acknowledgments

We extend a special thank you to Dr. Roxanne Connelly at the University of Florida for assistance in the identification of *Culex* species.

Literature Cited

- Bolling BG, JH Kennedy and EG Zimmerman.** 2005. Seasonal dynamics of four potential West Nile vector species in north-central Texas. *Journal of Vector Ecology* 30 (2): 186-194.
- Centers for Disease Control.** 2016. Zika Virus. <http://www.cdc.gov/zika/>. Accessed 13 Feb 2016.
- Darsie RF and RA Ward.** 2005. Identification and geographical distribution of the mosquitoes of North America, North of Mexico. University Press of Florida. Gainesville, FL., 384 pp.
- Hawley WA.** 1988. The biology of *Aedes albopictus*. *Journal of the American Mosquito Control Association* 4:2-39.
- Jamieson DH and LA Olson.** 1995. Recent establishment of the Asian tiger mosquito (*Aedes albopictus*) in Independence County, Arkansas. *Proceedings of the Arkansas Academy of Science* 49:80-81.
- Jamieson DH, LA Olson, and JD Wilhide.** 1994. A larval mosquito survey in northeastern Arkansas including a new record for *Aedes albopictus*. *Journal of the American Mosquito Control Association* 10:236-239.
- Kilpatrick MA, LD Kramer, SR Campbell, EO Alleyne, AP Dobson and P Daszak.** 2005. West Nile virus risk assessment and the bridge vector paradigm. *Centers for Disease Control and Prevention* 3:425-429.
- Miller MJ and JR Loaiza.** 2015. Geographic Expansion of the Invasive Mosquito *Aedes albopictus* across Panama—Implications for Control of Dengue and Chikungunya Viruses. *PLoS Neglected Tropical Diseases* 9(1): e0003383. doi:10.1371/journal.pntd.0003383
- Moore CG, DB Francy, DA Eliason and TP Month.** 1988. *Aedes albopictus* in the United States: rapid spread of a potential disease vector. *Journal of the American Mosquito Control Association* 4:356-361.
- Pfützner S, DH Jamieson and LA Olson.** 1998. The colonization of an Ozark Mountain city by the Asian tiger mosquito (*Aedes albopictus*). *Proceedings of the Arkansas Academy of Science* 52:13.
- Wilson J and DH Jamieson.** 2010. Presence of the Asian Tiger Mosquito in Northwest Arkansas. *Journal of the Arkansas Academy of Science* 64:152.