2016

Report on an Oomycete Infection (Heterokontophyta: Oomycota) in Northern Studfish, *Fundulus catenatus* (Fundulidae) from Tenmile Creek, Saline County, Arkansas

C. T. McAllister

Eastern Oklahoma State College, cmcallister@se.edu

H. W. Robison

Southern Arkansas University

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Zoology Commons](#)

Recommended Citation

McAllister, C. T. and Robison, H. W. (2016) "Report on an Oomycete Infection (Heterokontophyta: Oomycota) in Northern Studfish, *Fundulus catenatus* (Fundulidae) from Tenmile Creek, Saline County, Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 70 , Article 47.

DOI: <https://doi.org/10.54119/jaas.2016.7039>

Available at: <https://scholarworks.uark.edu/jaas/vol70/iss1/47>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Report on an Oomycete Infection (Heterokontophyta: Oomycota) in Northern Studfish, *Fundulus catenatus* (Fundulidae) from Tenmile Creek, Saline County, Arkansas

C.T. McAllister¹ and H.W. Robison²

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

²Department of Biology, Southern Arkansas University, Magnolia, AR 71754-9354

¹Correspondence: cmcallister@se.edu

Running Title: Oomycete Infection in *Fundulus catenatus*

Cases of oomycete (water mold) infestations in fishes are relatively common (Schäperclaus 1986). Indeed, among recent emerging infectious diseases responsible for severe population declines in fish taxa, fungal and oomycete microbes have emerged as significant contributors (Gozlen et al. 2014). In addition, the oomycetes negatively impact a wide array of other taxa including insects, zooplankton, nematodes, crayfish and amphibians (Phillips et al. 2008, Beakes et al. 2012; Sarowar et al. 2014). We are not aware of any previous reports of an oomycete infection in native fishes of Arkansas. Here, we document a noteworthy oomycete infestation in Northern Studfish, *Fundulus catenatus* Storer from central Arkansas.

Fourteen adult *F. catenatus* (mean $\pm$ 1SD total length [TL] 69.9 $\pm$ 7.6, range 62-93 mm) were collected with a backpack electrofisher on 15 November 2015 from Tenmile Creek off US 70, vicinity of Lonsdale, Saline County (34.545463°N, 92.753702°W). Specimens were placed in creek water and taken to the laboratory within 24 hr for processing. Fish were killed by immersion in a concentrated chlorotone solution and measured for TL. Two (14%) of the *F. catenatus* (70 and 76 mm) were noted of having a moderate growth of what appeared to be unknown white cotton wool-like patches on a small part of their body (head and chin) (Fig. 1). This growth was sampled by removing a portion with fine forceps, placed on a microscopic slide, stained with bromphenol cotton blue, and mounted with a coverslip. The slide was examined under a light microscope and photomicrographs were obtained. Voucher specimens of *F. catenatus* are deposited in the Henderson State University (HSU) collection, Arkadelphia.

Although it was not possible with confidence to determine with certainty which genus is present in this case without culturing and DNA sequence data, microscopic examination of the infestation revealed

Figure 1. Infected *Fundulus catenatus* (TL = 76 mm). A. Oomycete growth (arrows) on top of head, mouth and chin. Note that there appears to be no other growth on remainder of body. B. Close-up of same showing filamentous mycelia growing on the mouth and underside of chin with smaller colonies on top of head.

non-septate, multinucleate and unbranched or branched hyphae (Fig. 2) similar to those of the water mold (Oomycetes) Saprolegniales species (Webster and Weber 2007). The best-studied fish pathogenic oomycetes belong to this order including the genera *Achlya*, *Aphanomyces* and *Saprolegnia*. Organisms in this order reproduce asexually by releasing biflagellated spores formed in apical slightly swollen sporangia (Schäperclaus 1986, fig. 236).

Numerous other fishes collected on the same date at the Tenmile Creek site did not harbor any similar infestation, including *Camptostoma spadiceum*, *Gambusia affinis*, *Lepomis cyanellus* and *Noturus*

lachneri. In addition, we have, over the last 40+ yr, collected many fishes from all river drainages in the

Figure 2. Oomycete infection in *Fundulus catenatus*. A. Unstained non-septate hyphae (arrow); scale bar = 50 µm. B. Three balls of encysted zoospores (arrows); scale bar = 50 µm. C. Single zoospore ball (arrow); scale bar = 25 µm. B & C stained with bromophenol cotton blue.

state and never recall seeing such a similar infection on a single fish. Therefore, we suggest this is a fairly rare event and it does not appear to pose a serious risk as a potential pathogen on Arkansas fish populations.

We document the first report of an oomycete infection in a native Arkansas fish species. Additional research on this topic, including molecular studies, will surely extend our knowledge and further hosts could be discovered having similar infestations in the state.

Acknowledgments

The Arkansas Game and Fish Commission issued a scientific collecting permit to CTM. We thank Dr. R. Gozlen (IRD, Paris, France) for information on oomycete infections in fish, Dr. R. Tumilson (HSU) for expert curatorial assistance, and the McAllister clan for help in collecting at Tenmile Creek.

Literature Cited

- Beakes GW, SL Glockling and S Sekimoto. 2012. The evolutionary phylogeny of the oomycete “fungi”. *Protoplasma* 249:3–19.
- Gozlan RW, WL Marshall, O Lilje, CN Jessop, FH Gleason and D Andreou. 2014. Current ecological understanding of fungal-like pathogens of fish: What lies beneath? *Frontiers in Microbiology* 5:1–16.
- Phillips AJ, VL Anderson, EJ Robertson, CJ Secombes and P van West. 2008. New insights into animal pathogenic oomycetes. *Trends in Microbiology* 16:13–19.
- Sandoval-Sierra JV and J Diéguez-Urbeondo. 2015. A comprehensive protocol for improving the description of Saprolegniales (Oomycota): two practical examples (*Saprolegnia aenigmatica* sp. nov. and *Saprolegnia racemosa* sp. nov.). *PLoS ONE* 10(7): e0132999. Doi:10.1371/journal.pone.0132999
- Schäperclaus W. 1986. Fish diseases, Volume 1. Amerind Publishing Company, Pvt. Ltd. (New Delhi). p 1–594.
- Sarwar NM, M Saraiva, CN Jessop, O Lilje, FH Gleason and P van West. 2014. Infection strategies of pathogenic Oomycetes in fish. In Jones EGB, KD Hyde and KL Pang, editors. *Freshwater Fungi and fungal-like organisms*, De Gruyter (Berlin). p 217–243.
- Webster J and R Weber. 2007. *Introduction to Fungi*, 3rd Edition. Cambridge University Press, New York. 867 p.