2016

An Epistylus sp. (Ciliophora: Peritrichia: Epistylididae) Infestation on Green Sunfish, *Lepomis cyanellus* (Perciformes: Centrarchidae), from Arkansas

C. T. McAllister

Eastern Oklahoma State College, cmcallister@se.edu

H. W. Robison

Southern Arkansas University

S. E. Trauth

Arkansas State University

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Zoology Commons](#)

Recommended Citation

McAllister, C. T.; Robison, H. W.; and Trauth, S. E. (2016) "An Epistylus sp. (Ciliophora: Peritrichia: Epistylididae) Infestation on Green Sunfish, *Lepomis cyanellus* (Perciformes: Centrarchidae), from Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 70 , Article 48.

DOI: <https://doi.org/10.54119/jaas.2016.7040>

Available at: <https://scholarworks.uark.edu/jaas/vol70/iss1/48>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

An *Epistylus* sp. (Ciliophora: Peritrichia: Epistylididae) Infestation on Green Sunfish, *Lepomis cyanellus* (Perciformes: Centrarchidae), from Arkansas

C.T. McAllister^{1*}, H.W. Robison² and S.E. Trauth³

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

²Department of Biology, Southern Arkansas University, Magnolia, AR 71754-9354

³Department of Biological Sciences, Arkansas State University, State University, AR 72467

*Correspondence: cmcallister@se.edu

Running Title: *Epistylis* sp. on *Lepomis cyanellus* in Arkansas

Ciliates of the genus *Epistylus* Ehrenberg, 1830 are sessile peritrichous organisms often present as a branching colony with a short oral disc and collar, and non-contractile rigid stalk (Dias et al. 2006). There are about 120 described species that generally live in freshwater environments (Lynn 2008). Epistylids are not considered true parasites but are common on crayfishes (Vogelbein and Thune 1988), the carapace of turtles (Bishop and Jahn 1941, Bovee 1976) and on the eggs, skin and gills of fishes where they may contribute to mortality from partial smothering or secondary infection (Fischthal 1949, Reichenbach-Klinke 1973, Esch et al. 1976, Miller and Chapman 1976, Crites 1977, Hazen et al. 1978, Hoffman 1999). However, some historically considered these ciliates epibionts, which colonize the surface of live substrates and serve a commensal ecological role (Wahl 1989, Fernandez-Leborans et al. 2006). Lewis et al. (1978) reported *Epistylis* on 16 of 32 fishes examined in 2 North Carolina reservoirs, mainly from centrarchids, ictalurids, and moronids. Ictalurids and salmonids seem to be especially susceptible to infestation (Hubert and Warner 1975, Hoffman 1999). Epistylids use the host as an attachment substrate, so it can feed on bacteria and suspended particles in water.

Little is known about these ciliates on Arkansas fishes. Foissner et al. (1985) reported an epizootic of *Heteropolaria colisarum* Foissner and Shubert, 1977 on the scales and fins of cultured Green Sunfish (*Lepomis cyanellus*) from the Fish Farming Experimental Station, Stuttgart, Arkansas County. Interestingly, this infection was also shown in photos on the front cover of the second edition of Hoffman's (1999) classic, *Parasites of North American Freshwater Fishes*. Lom and Dyková (1992) have synonymized *H. colisarum* with *Epistylis*. However, to our knowledge, nothing has been published on this ciliate in native fishes from natural waters in the state. Here, we present a case of *Epistylis* sp. on *L. cyanellus*

with light microscopy and scanning electron microscopy (SEM) of the infestation.

On 14 October 2015, an adult (195 mm total length) *L. cyanellus* was collected with a backpack electroshocker from the South Fork of Fourche La Fave River at Hollis, Perry County (34.8706°N, 93.109458°W). As this specimen was noted to be unusual in possessing scales and spines on the dorsal fin with some unidentified whitish growth (Fig. 1), the fish was photographed alive. No growth was noted on the gills. The specimen was immediately overdosed by immersion in tricaine methanesulfonate and several scales were removed and placed in individual vials containing 70–95% (v/v) DNA grade ethanol and 10% neutral buffered formalin (NBF). The fish was preserved in 10% formalin and later transferred to 70% (v/v) ethanol. On return to the laboratory, scrapings were taken with fine forceps from scales originally placed in ethanol. Specimens were stained with Gomori trichrome, dehydrated in 95–100% (v/v) ethanol, cleared with xylene, mounted with Canada balsam and examined using light microscopy. Photomicrographs of stereoscopic samples were taken with Canon Powershot S3IS camera fitted with a Martin Microscope adaptor (Martin Microscope, Easley, SC). Photomicrographs for light microscopy were taken with a Swift M10 Series microscope fitted with a digital camera mount. For SEM, we transferred scales and liquid suspensions in 10% NBF through a graded series of increasing ethanol solutions (70–100%). Specimens were then extracted from vials with a pipette and placed onto segments of glass coverslips (18 × 4 mm) previously coated with poly-L-lysine. An Autosamdri®-815 critical point drier (Tousimis Research Corporation, Rockville, MD; 31°C, 1072 psi, ventilation rate ~100 psi/min) was used to remove excess ethanol from cells. Dehydrated specimens on coverslips were then adhered to rectangular copper transfer boats (25 × 5 mm) with double-sided tape and

Figure 1. *Lepomis cyanellus* with whitish growth on dorsal fin and scales. Scale bar = 50 mm. Photo by Uland Thomas.

mounted onto sticky-tapped 12 mm aluminum pin stubs. Copper boats and stubs were then coated with gold using a Cressington 108 sputter coater (Cressington Scientific Instruments Ltd, Watford, UK). Specimens examined for SEM were generated at the Arkansas Nano & Bio Materials Characterization Facility (UA-Fayetteville). A host voucher specimen was deposited in the Henderson State University Collection (HSU), Arkadelphia, Arkansas; a voucher slide of the infestation was deposited in the Harold W. Manter Laboratory of Parasitology (HWML), Lincoln, Nebraska as HWML 101962.

The unusual growth on scales of *L. cyanellus* (Fig. 1) was identified as an *Epistylis* sp. Examination of colonies using stereoscopic and light microscopy (Fig. 2) revealed colonies comprising a various number of individuals, with a branched and smooth, noncontractile stalk and fully expanded zooids. When examined by SEM (Fig. 3), apical views of contracted zooids were prominent, and cilia could be seen atop them. The macronucleus is typically horseshoe shaped and transversely oriented in the middle-adoral region of the cell.

Fischthal (1949) reported an *Epistylis* sp. from a darter in a Wisconsin stream, and Rogers (1971) found it on pond fishes in the southeastern United States. Cloutman (1975) reported an *Epistylis* sp. on White Bass (*Morone chrysops*) and Striped Bass (*Morone saxatilis*) in North and South Carolina. In addition, Crites (1977) found *Epistylis niagarae* Kellicott on Smallmouth Bass, *Micropterus dolomieu*, Rock Bass, *Ambloplites rupestris* and Freshwater Drum, *Aplodinotus grunniens* from Ohio.

Other fishes ($n = 28$) collected on the same date and locality did not possess *Epistylis* as follows: 5 Greenside Darters (*Etheostoma blennioides*), 1 Western Creek Chubsucker (*Erimyzon claviformis*), 8 Slender Madtoms (*Noturus exilis*), 1 Logperch (*Percina caprodes*), 6 Longnose Darters (*Percina nasuta*), 4 Blackspotted Topminnows (*Fundulus*

olivaceus), and 3 Wedgespot Shiners (*Notropis greeniei*). In addition, we have examined over 100 *L. cyanellus* from various watersheds in Arkansas over the last decade and never have noticed any with this growth.

Figure 2. Light microscopy of *Epistylis* sp. from *Lepomis cyanellus*. A. Colonies from skin scraping, stereoscopic view; trichrome stain. B. Branched colonies showing group of zooids on noncontractile stalks; unstained. C. Zooids showing macronucleus (arrow) and daughter cells from binary fission (arrowhead); unstained. D. Two zooids showing macronuclei (arrows); trichrome stain. E. Single elongate zooid showing cilia (arrows); trichrome stain. F. Single vase-shaped zooid showing macronucleus (arrow); unstained. Scale bars: A = 200 μ m; B–F = 50 μ m.

Figure 3. Scanning electron micrographs of *Epistylis* sp. on scales of *Lepomis cyanellus*. A. Low magnification showing colonies on single scale. Bar = 500 μ m. B. Higher magnification of apical view of contracted zooids. Bar = 50 μ m. C. Another apical view showing cilia on one colony (arrow). Bar = 20 μ m. D. Close-up showing cilia (arrow). Bar = 5 μ m.

In summary, we provide the initial report of an *Epistylis* sp. from a non-cultured *L. cyanellus* and the

first time SEM photomicrographs of this form have been documented. Future studies should include molecular analyses (18S rRNA sequences) to help further provide an identity of this species.

Acknowledgments

We thank Dr. C.M. Whipps (SUNY-ESF) for help in identification of *Epistylis* sp., U. Thomas (Chicago, IL) for Fig. 1, Drs. M. Benamara and B. Martin (Arkansas Nano & Bio Materials Characterization Facility, UA-Fayetteville) for technical assistance with SEM, and S.L. Gardner (HWML) and R. Tumilson (HSU) for expert curatorial assistance. The Arkansas Game and Fish Commission issued Scientific Collecting Permits to CTM and HWR.

Literature Cited

- Bishop EL Jr and TL Jahn.** 1941. Observations on colonial peritrichs (Ciliata; Protozoa) of the Okoboji region. *Proceedings of the Iowa Academy of Science* 48:417–421.
- Bovee EC.** 1976. New epizoic peritrichs of the soft shelled turtle *Trionyx muticus*. *Transactions of the American Microscopical Society* 95:682–687.
- Cloutman DG.** 1975. *Epistylis* on White Bass and Striped Bass. *Fish Health Section of the American Fisheries Society Newsletter* 3:6.
- Crites JL.** 1977. The occurrence of *Epistylis niagarae* (Ciliophora: Peritrichida) on fishes from the island region of western Lake Erie. *The Ohio Journal of Science* 77:193–194.
- Dias RJP, S D'Ávila and M D'Agosto.** 2006. First record of epibionts, peritrichids and suctorians (Protozoa, Ciliophora) on *Pomacea lineata* (Spix, 1827). *Brazilian Archives of Biology and Technology* 49:807–812.
- Esch GW, TC Hazen, RV Dimock Jr and JW Gibbons.** 1976. Thermal effluent and the epizootiology of the ciliate *Epistylis* and the bacterium *Aeromonas* in association with centrarchid fish. *Transactions of the American Microscopical Society* 95:687–693.
- Fernandez-Leborans G, MJH Cordoba and PG Arco.** 1997. Distribution of ciliate epibionts on the portunid crab *Liocarcinus depurator* (Decapoda, Brachyura). *Invertebrate Biology* 116:171–177.
- Fischthal JH.** 1949. *Epistylis*, a peritrichous protozoan, on hatchery brook trout. *Progressive Fish-Culturalist* 11:122–124.
- Foissner W, GL Hoffman and AJ Mitchell.** 1985. *Heteropolaria colisarum* Foissner & Shubert, 1977 (Protozoa: Epistylididae) of North American freshwater fishes. *Journal of Fish Diseases* 8:145–160.
- Hazen TC, ML Raker, GW Esch and CB Fliermans.** 1979. Ultrastructure of red-sore lesions on Largemouth Bass (*Micropterus salmoides*): Association of the ciliate *Epistylis* sp. and the bacterium *Aeromonas hydrophila*. *Journal of Protozoology* 25:351–355.
- Hoffman GL.** 1999. *Parasites of North American freshwater fishes*. 2nd ed. Comstock Publishing Associates (Ithaca, NY). 539 p.
- Hubert WA and MC Warner.** 1975. Control of *Epistylis* on channel catfish in raceways. *Journal of Wildlife Diseases* 11:241–244.
- Lewis RE, JR Siler and LL Olmsted.** 1978. Prevalence and intensity of *Epistylis* on fishes from two North Carolina reservoirs. *Proceedings of the Annual Conference Southeastern Association of Fish and Wildlife Agencies* 31:372–379.
- Lom J and I Dyková.** 1992. *Protozoan parasites of fishes*. Elsevier Science Publishers (Amsterdam, Netherlands). 315 p.
- Lynn DH (ed.).** 1988. *The ciliated Protozoa: Characterization, classification, and guide to the literature*. 3rd Edition. Springer (NY). 604 p.
- Miller RW and WR Chapman.** 1976. *Epistylis* and *Aeromonas hydrophila* infections in fishes from North Carolina reservoirs. *Progressive Fish-Culturalist* 38:165–168.
- Reichenbach-Klinke HH.** 1973. *Fish pathology: A guide to the recognition and treatment of diseases and injuries of fishes, with emphasis on environmental and pollution problems*. T.F.H. Publications (Neptune City, NJ). 492 p.
- Rogers WA.** 1971. Disease in fish due to the protozoan *Epistylis* (Ciliata: Peritricha) in the Southeastern U.S. *Proceedings of the 25th Annual Conference of the Southeastern Association Game and Fish Commission* 25:493–496.
- Vogelbein WK and RL Thune.** 1988. Ultrastructural features of three ectocommensal Protozoa attached to the gills of the red swamp crayfish, *Procambarus clarkii* (Crustacea: Decapoda). *Journal of Protozoology* 35:341–348.
- Wahl M.** 1989. Marine epibiosis. I. Fouling and antifouling some basic aspects. *Marine Ecology Progress Series* 58:175–189.