

1990

## Correction: Reproductive characteristics of South Florida *Sternotherus odoratus* and *Kinosternon baurii* (Testudines: Kinosternidae)

Walter E. Meshaka Jr.

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

---

### Recommended Citation

Meshaka, Walter E. Jr. (1990) "Correction: Reproductive characteristics of South Florida *Sternotherus odoratus* and *Kinosternon baurii* (Testudines: Kinosternidae)," *Journal of the Arkansas Academy of Science*: Vol. 44 , Article 41.  
Available at: <http://scholarworks.uark.edu/jaas/vol44/iss1/41>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact [scholar@uark.edu](mailto:scholar@uark.edu), [ccmiddle@uark.edu](mailto:ccmiddle@uark.edu).

Arkansas Academy of Science

GRIGGS, G. and T. ORR. 1979. Chromosomal aberrations resulting from ultraviolet exposures to G1 *Xenopus* cells. *Photochem. and Photobiol.* 30:363-368.

GRIGGS, G. and J. PAYNE. 1981. Time course of photoreactivation of UV-induced chromosomal aberrations and lethal damage in interphase *Xenopus* cells. *Photochem. and Photobiol.* 34:57-62.

KULP, S. and G. GRIGGS. 1989. Photoreactivation of lethal damage and damage leading to chromatid deletions induced in G1 phase hamster x *Xenopus* hybrid cells by UV. *Photochem. and Photobiol.* 50:185-191.

PERRY, P. and S. WOLFF. 1974. New giemsa method for the differential staining of sister chromatids. *Nature.* 251:156-158.

WALDREN, C. and R. JOHNSON. 1974. Analysis of interphase chromosome damage by means of premature chromosome condensation after x- and ultraviolet-irradiation. *Proc. Natl. Acad. Sci.* 71:1137-1141.

ROBERT WRIGHT, STEPHEN RUBLE, and GASTON GRIGGS, Department of Biology, John Brown University, Siloam Springs, AR 72761.

**Correction** — In the article "Reproductive characteristics of south Florida *Sternotherus odoratus* and *Kinosternon baurii* (Testudines: Kinosternidae)" by Walter E. Meshaka that appeared in Volume 42 (1988) of the Proceedings of the Arkansas Academy of Science, pages 111-112, the following correction should be noted. In the third paragraph, the first sentence should read "Average carapace lengths for eighteen sexually mature females ( $77 \pm 5.69$ ; range = 62-86) and ten sexually mature males ( $68 \pm 8.79$ ; range = 52-80) support Tinkle's (1961) findings of sexual dimorphism in southern populations."