

1982

Aquatic Macroinvertebrate Taxa Present in Two Ozark Springs in Randolph County, Arkansas

G. Randall Guntharp
Arkansas State University

George L. Harp
Arkansas State University

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Part of the [Terrestrial and Aquatic Ecology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Guntharp, G. Randall and Harp, George L. (1982) "Aquatic Macroinvertebrate Taxa Present in Two Ozark Springs in Randolph County, Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 36, Article 29. Available at: <https://scholarworks.uark.edu/jaas/vol36/iss1/29>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

LITERATURE CITED

- BARR, T. C. 1963. Ecological classification of cavernicoles. *Cave Notes* 5(2):9-12.
- HAWKSLEY, O., N. W. YOUNGSTADT, and J. O. YOUNGSTADT. 1980. A sabertooth cat, *Smilodon floridanus* from northwest Arkansas. *N. S. S. Bull.* 42:8-14.
- MCDANIEL, V. R., and J. E. GARDNER. 1977. Cave fauna of Arkansas: selected vertebrate taxa. *Proc. Ark. Acad. Sci.* 31:68-71.
- MCDANIEL, V. R., K. N. PAIGE, and C. R. TURLISON. 1979. Cave fauna of Arkansas: additional invertebrate and vertebrate records. *Proc. Ark. Acad. Sci.* 33:84-85.
- JAMES D. DUNIVAN, C. RENN TURLISON and V. RICK MCDANIEL. *Dept. of Biological Sciences, Arkansas State University, State University, Arkansas 72467.*
- MCDANIEL, V. R., and K. L. SMITH. 1976. Cave fauna of Arkansas: selected invertebrate and vertebrate taxa. *Proc. Ark. Acad. Sci.* 30:57-60.
- PAIGE, K. N., C. R. TURLISON, and V. R. MCDANIEL. 1981. A second record of *Typhlichthys subterraneus* (Pisces: Amblyopidae) from Arkansas. *Southw. Nat.* 26(1):67.
- YOUNGSTADT, N. W., and J. O. YOUNGSTADT. 1980. Prehistoric bear signs of Hurricane River Cave. *N. S. S. Bull.* 42:1-7.

AQUATIC MACROINVERTEBRATE TAXA PRESENT IN TWO
OZARK SPRINGS IN RANDOLPH COUNTY, ARKANSAS

Previous studies concerning the aquatic macroinvertebrates found in the Ozark Plateau of Arkansas have dealt only with stream populations (Sublette, 1956; Van Kirk, 1962; Aggus and Warren, 1965; Robison and Harp, 1971; McGary and Harp, 1972; Cather and Harp, 1975). None have involved aquatic macroinvertebrate communities in cold water springs. The Salem Plateau section of the Ozark Plateau province supplies many cold water springs of varying size. This is largely due to the abundance of limestone and dolomite, which provides a good geologic base for the development of springs (Cronce, 1930; Thornbury, 1965). Spring basins provide a slightly different habitat for aquatic macroinvertebrate communities. Temperatures and water levels generally are confined to a more narrow range than in a stream and experience minimal seasonal variations with regard to several other physicochemical parameters (Clifford, 1966). However, temperatures are not so limiting as in cold water discharges from main stem reservoirs (McGary and Harp, 1972).

The springs that were sampled were designated Spring I and Spring II. Spring I is located in S7, T20N, R1E in Randolph County, Arkansas, and is considered to be a permanent spring. It is at an elevation of 110 m and has an average volume flow of 65.5 cc/sec that flows in a southward direction. This spring is surrounded by grassy pastureland which has not had any domestic grazing for more than eight years. The spring is shaded by one sycamore tree (*Plantanus occidentalis*) with one eastern red cedar (*Juniperus virginiana*) located about 15 m to the west. A small, spring-fed pond occurs about 100 m to the west. The spring basin was dredged over a year ago and its greatest depth is 1 m. The north end has an almost vertical drop, and this deep end covers approximately 25 percent of the basin's surface area. The rest of the basin ranges in depth from 0.08 to 0.23 m. The basin has a mean width of 2.13 m.

Spring II is located in S27, T20N, R2W, Randolph Co., Arkansas. It is a smaller spring in terms of volume flow and basin size. The volume flow ceased temporarily during the driest part of the 1980 summer drought, but the spring was still able to maintain most of its basin size. This spring lies at an elevation of 140 m and has an average volume flow of 57.8 cc/sec. Spring II is not nearly as open as Spring I, but rather is surrounded by an oak-hickory forest with thick undergrowth which provides shade throughout the entire day. It has been free of domestic animals for more than eight years and is about 200 m from a small creek. This spring varies in depth from 0.08 to 0.18 m. Its basin has a mean width of 1.22 m.

Data were collected biweekly from each spring from 26 April through 13 June 1981. Ambient and water temperatures were taken using a standard Celsius thermometer. Water was tested for pH, nitrate (NO₃), carbon dioxide, ammonium nitrogen and phosphate (PO₄) using Hach water test kits 17-N, NI-11, CA-23, NI-8 and PO-19/PO-19A, respectively. Spatial measurements were made by using a standard yardstick an converting values to metric units. Spring volume flow was measured at the outflow from the basin by a modification of the mean-sectional method (Butler, 1957). The first and third macroinvertebrate samples were collected by use of a fine mesh aquatic dip net. The second and fourth macroinvertebrate samples were obtained by use of a 12 volt-DC BioQuip blacklight collecting light. Two adult odonates were obtained using an aerial net. Samples were preserved in 70 percent ethanol. Aquatic organisms were identified according to Pennak (1978), with the exception of Gerridae which were identified according to Kittle (1980). All samples were quantified by standardizing the length of the sampling period, specifically one hour.

To minimize the effects of extrinsic factors, springs were sampled under similar logistic conditions. Water quality was quite similar for both springs, and little fluctuation occurred in the physicochemical parameters measured during the study period (Table 1). This is typical of springs (Clifford, 1966), and is derived from the ground waters flowing through similar geologic formations. Carbon dioxide, ammonium nitrate and phosphate were never present in detectable concentrations in either spring. A total of 225 organisms representing 13 taxa was collected from Spring I, while only 80 organisms of 8 taxa were collected from Spring II (Table 2). The reduced taxonomic diversity resulted primarily from the limiting factor of a rather low constant water temperature. Numerical standing crop was moderate, considering the small basin size of these two springs. The lower numerical standing crop and reduced diversity in Spring II were probably derived from its less constant discharge (which ceased during the height of the 1980 drought). Further, this spring is shaded throughout the day, thus reducing photosynthesis and therefore autochthonous food materials. The smaller size of this basin may also have been a contributing factor. *Crangonyx* was the most abundant organism in both springs, constituting 67 and 51 percent of the organisms in Springs I and II, respectively. *Isoptera* was the second most abundant organism in Spring I (11%), while *Gerris* was second most abundant in Spring II (29%). *Crangonyx* is an omnivorous, general scavenger. *Gerris* usually eat a variety of crustaceans and aquatic insects that they catch just below the water surface, but alternatively may consume terrestrial insects that fall into the water. *Isoptera* is polyphagous, but is primarily predatory on other aquatic insects (Pennak, 1978). The dominance of these organisms implies that primary production and other autochthonous food sources were minimal in these two springs.

General Notes

Table 1. Physicochemical parameters for Springs I and II, Randolph County, Arkansas (26 April-13 June 1981).

Date	Spring	Air Temp. °C	Water Temp. °C	pH	nitrate ppm
26-IV	I	31	18	7.5	20
	II	30	15	7.5	22
16-V	I	23	15	8.0	20
	II	22	14	7.7	20
23-V	I	26	17	8.0	20
	II	26	15	7.7	20
13-VI	I	27	17	8.0	20
	II	27	16	7.5	20

Table 2. Aquatic macroinvertebrates, expressed as relative abundance, for Springs I and II, Randolph County, Arkansas (26 April-13 June 1981).

TAXA	Spring I	Spring II
GASTROPODA	0	7
AMPHIPODA		
<i>Crangonyx forbesi</i> (Hubricht Mackin)	150	41
<i>Gammarus pseudolimnacus</i> Bousfield	18	0
<i>Hyalella asteca</i> (Saunders)	2	0
DECAPODA	0	1
HEMIPTERA		
<i>Corixa remigia</i> Say	8	23
<i>Sigara grossolineata</i> Hungford	1	0
<i>Microvelia americana</i> (Uhler)	0	2
<i>Microvelia australis</i> Sueno	0	4
PLECOPTERA		
<i>Perlwa placida</i> (Hagen)	1	0
<i>Isoperla</i>	25	0
EPHEMEROPTERA		
Heptageniidae	7	0
ODONATA		
<i>Calopteryx maculata</i> (Beauvois)	1	0
<i>Arista pluma</i> Calvert	1	0
<i>Platania lydia</i> Drury	0	1
COLEOPTERA		
<i>Laccophilus pictus</i> Sharp	4	0
DIPTERA		
<i>Culiseta inornata</i> (Williston)	4	1
<i>Tabanus atratus</i> Fabricius	3	0
Total	235	80

We thank Ron Smith for his assistance in identifying many of the insects. We also thank Larry Olson for his identification of the Culicidae.

G. RANDALL GUNTARP and GEORGE L. HARP, Department of Biological Sciences, Arkansas State University, State University, Arkansas 72467.

LITERATURE CITED

- AGGUS, L. R., and L. O. WARREN. 1965. Bottom organisms of the Beaver Reservoir basin: a pre-impoundment study. J. Kans. Entomol. Soc. 38:163-178.
- BUTLER, S. S. 1957. Engineering hydrology. Prentice-Hall, Englewood Cliffs, N. J. 356p.
- CATHER, M. R., and G. L. HARP. 1975. The aquatic macro-invertebrate fauna of an Ozark and a Deltaic stream. Proc. Ark. Acad. Sci. 29:30-35.
- CLIFFORD, H. F. 1966. Some limnological characteristics of six Ozark streams. Mo. Dept. of Conserv., D-J Series No. 4. 55p.
- CRONEIS, C. 1930. Geology of the Arkansas Paleozoic area. Ark. Geol. Survey Bull. 3. 457p.
- KITTLE, P. D. 1980. The water striders (Hemiptera: Gerridae) of Arkansas. Proc. Ark. Acad. Sci. 34:68-71.
- MCGARY, J. L., and G. L. HARP. 1972. The benthic macro-invertebrate community of the Greer's Ferry Reservoir cold tailwater, Little Red River, Arkansas. Proc. SE Assoc. Game and Fish Comms. 26:490-500.
- PENNAK, R. W. 1978. Fresh-water invertebrates of the United States. John Wiley & Sons, Inc., New York. 803p.
- ROBISON, H. W., and G. L. HARP. 1971. A pre-impoundment limnological investigation of the Strawberry River in northeastern Arkansas. Proc. Ark. Acad. Sci. 25:70-79.
- SUBLETTE, J. E. 1956. Seasonal changes in bottom fauna of an Ozark headwater stream (Clear Creek, Washington County, Arkansas). Southw. Nat. 1:148-156.
- THORNBURY, W. D. 1965. Regional geomorphology of the United States. John Wiley & Sons, Inc., New York. 609p.
- VAN KIRK, R. R. 1962. Some ecological and physical aspects of a cool water stream in the northwestern Arkansas with special reference to trout production. Unpub. M.S. thesis, Univ. Ark., Fayetteville. 56p.

DARK REPAIR OF LETHAL DAMAGE INDUCED IN A HYBRID MAMMALIAN TISSUE CULTURE CELL LINE BY ULTRAVIOLET LIGHT*

Dark (non-photoreactivation) intracellular mechanisms that repair ultraviolet light (UV)-induced lesions in nuclear DNA are known to contribute significantly to the resistance shown by many prokaryotic cells to such UV-induced effects as loss of colony forming ability, mutations, and division delay (Rupert and Harm, 1966; Smith, 1977). A number of studies indicate that at least two such mechanisms function in some mammalian cells (Cleaver, 1974). HeLa cancer tissue culture cells possess a mechanism that may be analogous to the "excision" repair mechanism found in many UV-resistant bacteria (Regan et al., 1968). Another mechanism has been detected in rodent cells (Cleaver, 1974) that is a post DNA replication repair process similar to "recombinational" (recom) repair found in some UV-resistant bacteria (Rupp and Howard-Flanders, 1968). A study by Meyn et al. (1974) suggests that both of these mechanisms function in many mammalian cells, but with varying efficiencies. For example, V79 hamster cells exhibit both excision and recom repair but the recom repair mechanism dominates, while HeLa cells appear to exhibit efficient exci-