

1980

Recent Mollusca of Arkansas with Annotations to Systematics and Zoogeography

Mark E. Gordon
University of Arkansas, Fayetteville

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Part of the [Terrestrial and Aquatic Ecology Commons](#)

Recommended Citation

Gordon, Mark E. (1980) "Recent Mollusca of Arkansas with Annotations to Systematics and Zoogeography," *Journal of the Arkansas Academy of Science*: Vol. 34, Article 18.

Available at: <https://scholarworks.uark.edu/jaas/vol34/iss1/18>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in *Journal of the Arkansas Academy of Science* by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

RECENT MOLLUSCA OF ARKANSAS WITH ANNOTATIONS TO SYSTEMATICS AND ZOOGEOGRAPHY

MARK E. GORDON
Department of Zoology
University of Arkansas-Fayetteville
Fayetteville, Arkansas 72701

ABSTRACT

A total of 223 taxa of Mollusca are presently known from Arkansas. The geological history and geomorphology of the region, particularly the presence of the Interior Highlands, have been responsible for the development of a diverse molluscan fauna. Thirty regionally endemic forms are included among the 107 terrestrial gastropods, 36 aquatic gastropods, 65 unionacean mussels, and 15 sphaeriacean clams.

INTRODUCTION

The first survey of Arkansas Mollusca was Sampson's (1893, 1894) report of the Gastropoda and Sphaeriidae. The Unionidae were cataloged by Call (1895). Since these initial reports, there have been few comprehensive publications concerning Arkansas mollusks. During the interim, limited species surveys, several species descriptions, and various miscellaneous studies have appeared periodically (e.g. Ortmann and Walker, 1912; Wheeler, 1914, 1918; Rehder, 1932; Kraemer, 1970; Gordon et al. 1980a). Recently, Hubricht (1972) published a fairly comprehensive list (89 taxa) of terrestrial gastropods and Gordon et al. (1980b) presented a historical review and consideration of the regional distribution patterns of the Unionacea with respect to Arkansas.

METHODS AND RESULTS

The molluscan fauna of Arkansas has been investigated by personal collecting and examination of major museum collections (see Gordon et al. 1980a, b). This has been augmented by published records which are either based on museum reference collections or may be collaborated by more than one published account. Nomenclature follows Burch (1962) and Hubricht (1972) for the terrestrial Gastropoda; Baker (1911), Goodrich (1939), Clench (1962), Clench and Fuller (1965), Clarke (1973), and Thompson (*pers. comm.*) for aquatic Gastropoda; Ortmann and Walker (1922), Clarke (1973), and Gordon et al. (1980b) for the Unionacea; and Herrington (1962) and Burch (1975) for the Sphaeriacea.

A total of 223 taxa of Mollusca are presently known from Arkansas. This represents 107 terrestrial Gastropoda (Table 1), 36 aquatic Gastropoda (Table 2), and 80 Bivalvia (Table 3). The bivalves include 65 unionacean mussels and 15 sphaeriacean clams, five of which are listed provisionally. Fifteen terrestrial gastropods, seven aquatic gastropods, and eight unionaceans are endemic within the Interior Highlands. No endemic sphaeriaceans are known. Some annotations are necessary for several species. These are as follows:

1. Subspecific nomenclature represents geographic subspecies and not ecophenotypic variation.
2. Current research and revision of the Succineidae will probably result in a reduction of species through synonymy (Wu, *pers. comm.*).
3. *Somatogyrus wheeleri* and *S. amnicoloides* appear to be known only by the type specimens. These species were collected together at the same locality. Further study may show them to be conspecific. The type locality is now receiving hypolimnetic release from DeGray Reservoir.
4. Four ecophenotypes of *Goniobasis potosiensis* have been described as subspecies. The forms known as *plebeius* Anthony,

the typical form, and *crandalli* Pilsbry, a stunted form with rounded whorls from Mammoth Springs, Fulton County, occur in Arkansas.

5. The taxonomy of *Anculosa arkansensis* is unclear. Goodrich (1939) stated that its general appearance was close to *Nitocris trilineata*. The shell of specimens I collected from tributaries of the North Fork of the White River are similar to *Nitocris*. Anatomical examination will be necessary before the generic position of this species can be confirmed. If a *Nitocris* species, this may represent a relict population of a previously described species.
6. Te (1975) has concluded that *Physa anatina* Lea is a form of *P. virgata*.
7. Clarke (1973) has concluded that *Fusconaia undata* (Barnes) is more than likely the large river ecophenotype of *F. flava*.
8. *Pleurobema cordatum* includes a complex group of ecophenotypes that have generally been given subspecific status. These include, by increasing inflatedness of the shell, *coccineum* Conrad (also known as *missouriensis* Marsh), *catillus* Conrad, *plenum* Lea, and *pyramidatum* Lea (Ortmann, 1919).
9. *Anodontia grandis* is a substantially variable species with several ecophenotypes. Synonymized here is the slough form *corpulenta* Cooper.
10. Strophitini is proposed as a new tribe within the unionid subfamily Anodontinae. The basis for this division is the digenae arrangement of the marsupium in the outer gill. Ortmann (1912) considered this to be the most advanced gill form in the Anodontinae. *Strophitus*, previously classified in the Anodontini (Clarke, 1973), is the only genus in the Unionacea that has this form of gill. Ortmann (1912) noted some similarities in the anatomy (primarily pigmentation) and the shell (beak sculpture and other undefined characters) to the *Alasmidonta*. The shell is not necessarily a good phylogenetic character (Heard and Guckert, 1970); however, the general outline of *Strophitus* often resembles that of the *Alasmidontini*. Vestigial swellings in the area of the pseudocardinals suggest an association with the *Alasmidontini* in which the pseudocardinals are single in both valves and the interdentum of the left valve is reflected into a tooth-like structure. These dental characters appear to hold true throughout the *Alasmidontini*. In *Strophitus*, the vestigial swellings correspond to the position of the single pseudocardinals and the reflected interdentum. The tribe is monogeneric — *Strophitus*.
11. *Actinonaia carinata* includes a form known as *gibba* Simpson. This is probably an ecophenotypic variation.
12. The shell of *Actinonaia ellipsiformis pleasii* is distinct from *A. e. ellipsiformis*. It is geographically restricted to the White River system of the Ozark Plateaus in which *A. e. ellipsiformis* is not distributed. *Actinonaia ellipsiformis pleasii* appears to be a true subspecies, if not a separate species.

13. *Lampsilis reeveiana* (Lea) is recognizable from its type description and lithograph. The holotype is extant in the U.S. National Museum of Natural History (USNMNH 8505). Previous difficulty in identifying this endemic Ozark Plateaus species has been partially due to Lea's (1852) mispublication of the type locality: Alexandria, Louisiana. This should be White River, Arkansas. *Reeveiana* has 35 years priority to Call's (1887) *brevicula*.
14. Most of the specimens of *Lampsilis ovata* found in Arkansas are the form known as *ventricosa* Barnes; although, some specimens are occasionally referred to as *satura* Lea. *Ventricosa* is the small river ecophenotype (Ortmann, 1919; van der Schalie, 1938). The relationship of *satura* is not presently known, it may be a southern cline of *ventricosa*. Clarke (1973) applies *ventricosa* only at an infraspecific level.
15. The taxonomy of *Lampsilis orbiculata* and *L. higginsii* is unclear. They are quite similar in appearance and have variously been considered separate species, subspecies, and the same species (Simpson, 1914; Ball, 1922; Baker, 1928; van der Schalie and van der Schalie, 1950; Parmalee, 1967). *Lampsilis higginsii* is the Mississippi River counterpart of *L. orbiculata* of the Ohio River system. Simpson (1914) identified a subspecies of *L. higginsii* that he considered intermediate between the two "species." Buchanan and Oesch (*pers. comm.*) recently have collected both forms from rivers in Missouri that flow into Arkansas (Little Black and St. Francis rivers). The presence of both forms in southern Missouri and Arkansas is quite possible when the former channels of the Mississippi and Ohio rivers are examined (see Gordon et al. 1980b). For the present, the two forms are listed as separate species.

Table 1. Arkansas Terrestrial Gastropoda.

Subclass Prosobranchia Edwards	Suborder Sigmurethra Pilsbry	Subfamily Gastrodantinae Tryon
Order Archaeogastropoda Thiele	Family Philomycidae Keferstein	<i>Ventridens demissus</i> (Binney)
Family Helicinidae Ferrussac	<i>Philomycus carolinianus</i> (Bosc)	<i>V. ligera</i> (Say)
<i>Helicina (Oligyra) orbiculata</i> (Say)	<i>Pallifera hemphilli marmorata</i> Pilsbry	<i>Zonitoides arboreus</i> (Say)
Subclass Pulmonata Cuvier	<i>P. mutabilis</i> Hubricht	<i>Striatura (Pseudohyalina) meridionalis</i>
Order Basommatophora Keferstein	<i>P. ragsdalei</i> (Webb)	(Pilsbry and Ferriss)
Family Carychiidae Jeffreys	Family Enodontidae Pilsbry	Family Haplotrematidae Baker
<i>Carychium exiguum</i> (Say)	Subfamily Enodontinae Pilsbry	<i>Haplotrema concavum</i> (Say)
Order Stylommatophora Schmidt	<i>Anguispira alternata</i> (Say)	Family Bulimulidae Pilsbry
Suborder Orthurethra Pilsbry	<i>A. strongyloides</i> (Pfeiffer)	<i>Bulimulus (Rhabdotus) dealbatus</i> (Say)
Family Cionellidae Koblet	<i>Discus patulus</i> (Deshayes)	Family Polygyridae Pilsbry
<i>Cionella lubrica</i> (Muller)	Subfamily Helicodiscinae Pilsbry	Subfamily Polygyrinae Pilsbry
Family Valloniidae Morse	<i>Helicodiscus (Helicodiscus) parallelus</i> (Say)	<i>Polygyra (Daedalochila) leporena</i> (Gould)
<i>Vallonia perspectiva</i> Sterki	<i>H. (Hebetodiscus) singleyanus</i> (Pilsbry)	<i>P. (D.) texasiana</i> (Moricand)
Family Pupillidae Turton	<i>H. (?) jacksoni</i> Hubricht	<i>P. (D.) tridentoides</i> (Bland)
Subfamily Pupillinae Pilsbry	<i>H. (?) notius</i> Hubricht	<i>P. (D.) dorfeuilliana</i> Lea
<i>Pupoides albilabris</i> (Adams)	Subfamily Punctinae Morse	<i>P. (D.) jacksoni</i> (Bland) ¹
Subfamily Gastrocoptinae Pilsbry	<i>Punctum (Punctum) minutissimum</i> (Lea)	<i>P. (D.) peregrina</i> Rehder ¹
<i>Gastrocopta (Gastrocopta) pellucida</i>	<i>P. (Toltecia) vitreum</i> Baker	<i>P. (?) lithica</i> Hubricht
<i>hordencella</i> (Pilsbry)	Family Limacidae Gray	<i>Stenotrema (Stenotrema) stenotrema</i>
<i>G. (G.) procera</i> (Gould)	Subfamily Limacinae Gray	(Pfeiffer)
<i>G. (G.) rupicola</i> (Say)	<i>Limax flavus</i> Linnaeus	<i>S. (S.) labrosum</i> (Bland) ¹
<i>G. (Albinula) armifera</i> (Say)	<i>Lehmannia poirieri</i> (Mabille)	<i>S. (S.) pilsbryi</i> (Ferriss) ¹
<i>G. (A.) contracta</i> (Say)	<i>Deroceras laeve</i> (Muller)	<i>S. (S.) blandianum</i> (Pilsbry) ¹
<i>G. (A.) holzingeri</i> (Sterki)	Subfamily Milacinae Cockerell	<i>S. (S.) unciniferum</i> (Pilsbry) ¹
<i>G. (Vertigopsis) tappaniana</i> (Adams)	<i>Milax gagates</i> (Draparnaud)	<i>S. (Euchemotrema) fraternum</i> (Say)
<i>G. (Privatula) corticaria</i> (Say)	Family Zonitidae Pilsbry	<i>S. (E.) leai</i> (Binney)
Subfamily Vertigininae Thiele	Subfamily Zonitinae Pilsbry	<i>Practicolella berlandieriana</i> (Moricand)
<i>Vertigo (Vertigo) ovata</i> Say	<i>Retinella (Glyphyalus) wheateleyi</i> (Bland)	<i>Mesodon (Mesodon) clausus</i> (Say)
<i>V. (V.) rugosula</i> Sterki	<i>R. (G.) lewisiana</i> (Clapp)	<i>M. (M.) thyroideus</i> (Say)
<i>V. (V.) tridentata</i> Wolf	<i>R. (Glyphalinia) indentata</i> (Say)	<i>M. (M.) zaitetus</i> (Binney)
<i>V. (Angustula) milium</i> (Gould)	<i>R. (G.) laticola</i> Hubricht	<i>M. (M.) elevatus</i> (Say)
<i>V. (Vertillaria) oscariana</i> Sterki	<i>R. (G.) solida</i> (Baker)	<i>M. (M.) binneyanus</i> (Pilsbry) ¹
Family Strobilopsidae Jooss	<i>Mesomphix (Omphalina) capnodes</i> (Binney)	<i>M. (M.) clenchi</i> (Rehder) ¹
<i>Strobilops (Strobilops) aenea</i> Pilsbry	<i>M. (O.) cupreus</i> (Rafinesque)	<i>M. (M.) indianorum</i> (Pilsbry) ¹
<i>S. (S.) labyrinthica</i> (Say)	<i>M. (O.) friabilis</i> (Binney)	<i>M. (M.) kiowaensis</i> (Simpson) ¹
<i>S. (S.) texasiana</i> (Pilsbry and Ferriss)	<i>M. (?) globosus</i> (Macmillan)	<i>M. (M.) roemeri</i> (Pfeiffer)
Suborder Heterurethra Pilsbry	<i>Paravitrea (Paravitrea) significans</i>	<i>M. (Patera) perigraptus</i> Pilsbry
Family Succineidae Beck	(Bland)	<i>M. (Inflctarius) inflctus</i> (Say)
<i>Succinea (Novisuccinea) ovalis</i> Say	<i>P. (P.) simpsoni</i> (Pilsbry) ¹	<i>M. (I.) magazinensis</i> (Pilsbry and Ferriss) ¹
<i>S. (Calcisuccinea) concordialis</i> Gould	<i>P. (P.) aulacogyra</i> (Pilsbry and Ferriss) ¹	Subfamily Triodopsinae Pilsbry
<i>S. (C.) luteola</i> Gould	<i>P. (P.) petrophila</i> (Bland)	<i>Triodopsis (Triodopsis) cragii</i> Call ¹
<i>S. (?) indiana</i> Pilsbry	<i>P. (Paravitreops) multidentata</i> (Binney)	<i>T. (T.) neglecta</i> (Pilsbry) ¹
<i>S. (?) witteri</i> Shimek	<i>Hawaii minuscula</i> (Binney)	<i>T. (T.) vultuosa</i> (Gould)
<i>Catinella avara</i> (Say)	Subfamily Euconulinae Baker	<i>T. (Xolotrema) fosteri</i> (Baker)
<i>C. oklahomarum</i> (Webb)	<i>Euconulus chersinus</i> (Say)	<i>T. (X.) obstricta</i> (Say)
<i>C. texana</i> Hubricht	<i>Guppya sterkii</i> (Dall)	<i>T. (Neohelix) albolabris</i> (Say)
<i>C. vermeta</i> (Say)		<i>T. (N.) divesta</i> (Gould)
<i>C. wandae</i> (Webb)		<i>T. (N.) multilineata</i> (Say)
<i>Oxyloma (Neoxyloma) salleana</i> (Pfeiffer)		

¹Endemic to the Interior Highlands Region.

Table 2. Arkansas Aquatic Gastropoda.

Subclass Prosobranchia Edwards	Subfamily Amnicolinae Tryon	Superfamily Physacea
Order Mesogastropoda Thiele	<i>Amnicola cora</i> Hubricht ¹	Family Physidae Fitzinger
Superfamily Viviparacea	Family Pomatiopsidae Stimpson	<i>Physa gyrina</i> Say
Family Viviparidae Gray	<i>Pomatiopsis lapidaria</i> (Say)	<i>P. virgata</i> Say
Subfamily Viviparinae Gill	Superfamily Cerithacea	Superfamily Planorbacea
<i>Viviparus subpurpureus</i> (Say)	Family Pleuroceridae Fischer	Family Planorbidae Rafinesque
<i>V. intertextus</i> (Say)	<i>Lithasia verrucosa</i> (Rafinesque)	Subfamily Planorbinae Rafinesque
<i>V. georgianus</i> (Lea)	<i>Pleurocera alveare</i> (Conrad)	Tribe Planorbini
Subfamily Lioplacinae Gill	<i>P. canaliculatum</i> (Say)	<i>Gyraulus parvus</i> (Say)
<i>Lioplax suculosa</i> (Menke)	<i>P. acuta</i> Rafinesque	Tribe Helisomatini
<i>Campeloma subsolidum</i> (Anthony)	<i>Goniobasis potosiensis</i> Lea ²	<i>Promenetus exacuus</i> (Say)
Superfamily Rissoacea	<i>Anculosa arkansensis</i> Hinkley ¹	<i>Menetus (Micromenetus) dilatatus</i> (Gould)
Family Hydrobiidae Troschel	Subclass Pulmonata Cuvier	<i>Helisoma (Helisoma) anceps</i> (Menke)
Subfamily Hydrobiinae Troschel	Order Basommatophora Keferstein	<i>H. (Piersoma) trivolvis</i> (Say)
<i>Cincinnatia integra</i> (Say)	Superfamily Lymnaea	Family Ancyliidae Rafinesque
<i>Probythinella binneyana</i> (Hannibal)	Family Lymnaeidae Rafinesque	<i>Laevapex fuscus</i> (Adams)
<i>Pyrgulopsis ozarkensis</i> Hinkley ¹	<i>Lymnaea (Pseudosuccinea) columella</i> (Say)	<i>L. diaphanus</i> (Haldeman)
Subfamily Lithoglyphinae Fischer	<i>L. (Fossaria) humilis</i> (Say)	<i>Ferrissia rivularis</i> (Say)
<i>Somatogyrus crassilabris</i> Walker ¹	<i>L. (F.) obrussa</i> (Say)	<i>F. walkeri</i> (Pilsbry and Ferriss)
<i>S. subglobosus</i> (Say)	<i>L. (Bakerilymnaea) bulimoides</i> (Lea)	<i>F. fragilis</i> (Tryon)
<i>S. wheeleri</i> Walker ²		
<i>S. amnicoloides</i> Walker ²		

¹Endemic Interior Highlands species restricted to the Ozark Plateaus.

²Endemic Interior Highlands species restricted to the Ouachita Mountains.

³Endemic Interior Highlands species.

Table 3. Arkansas Bivalvia.

Order Eulamellibranchia	<i>Arkansia wheleri</i> Ortmann and Walker ²	<i>Ligumia subrostrata</i> (Say)
Superfamily Unionacea	Tribe Anodontini	<i>L. recta</i> (Lamarck)
Family Margaritiferidae	<i>Anodonta (Pyganodon) grandis</i> Say	<i>Lampsilis anodontoides</i> (Lea)
Subfamily Cumberlandiinae	<i>A. (Utterbackia) imbecilis</i> Say	<i>L. radiata siliquoides</i> (Barnes)
<i>Cumberlandia monodonta</i> (Say)	<i>A. (U.) suborbiculata</i> Say	<i>L. hydiana</i> (Lea)
Family Unionidae	<i>Andontoides ferussacianus</i> (Lea)	<i>L. reeveiana</i> (Lea) ¹
Subfamily Ambleminae	<i>Simpsoniconcha ambigua</i> (Say)	<i>L. streckeri</i> Frierson
Tribe Amblemini	Tribe Strophitini n. t.	<i>L. rafinesqueana</i> Frierson ¹
<i>Fusconaia flava</i> (Rafinesque)	<i>Strophitus undulatus</i> (Say)	<i>L. ovata</i> (Say)
<i>F. ebena</i> (Lea)	Subfamily Lampsilinae	<i>L. orbiculata</i> (Hildreth)
<i>F. ozarkensis</i> (Call) ¹	Tribe Ptychogenini	<i>L. higginsii</i> (Lea)
<i>Amblema plicata</i> (Say)	<i>Ptychobranhus occidentalis</i> (Conrad) ³	<i>Dysnomia (Truncilopsis) triquetra</i> (Rafinesque)
<i>Quadrula pustulosa</i> (Lea)	Tribe Mesogenini	<i>D. (Torulosa) florentina</i> (Lea)
<i>Q. nodulata</i> (Rafinesque)	<i>Obliquaria reflexa</i> Rafinesque	<i>D. (T.) turgidula</i> (Lea)
<i>Q. quadrula</i> (Rafinesque)	<i>Cyprogenia aberti</i> (Conrad) ³	Superfamily Sphaeriacea
<i>Q. metanevra</i> (Rafinesque)	Tribe Heterogenini	Family Corbiculidae
<i>Q. cylindrica</i> (Say)	<i>Obovaria olivaria</i> (Rafinesque)	<i>Corbicula cf. fluminea</i> (Muller)
<i>Tritogonia verrucosa</i> (Rafinesque)	<i>O. jacksoniana</i> Frierson	Family Sphaeriidae
<i>Plectomerus dombeyanus</i> (Valenciennes)	<i>Actiononiais carinata</i> (Barnes)	<i>Sphaerium striatinum</i> (Lamarck)
Tribe Megalonaiaidini	<i>A. ellipsiformis ellipsiformis</i> (Conrad)	<i>Musculium lacustre</i> (Muller)
<i>Megalonaia gigantea</i> (Barnes)	<i>A. ellipsiformis pleasii</i> (Marsh) ¹	<i>M. partumeium</i> (Say)
Subfamily Unioninae	<i>Truncilla truncata</i> Rafinesque	<i>M. securis</i> (Prime)
Tribe Pleurobemini	<i>T. donaciformis</i> (Lea)	<i>M. transversum</i> (Say)
<i>Cyclonaias tuberculata</i> (Rafinesque)	<i>Plagiola lineolata</i> Rafinesque	<i>Pisidium (Cyclocalyx) casertanum</i> (Poli)
<i>Pleurobema cordatum</i> (Rafinesque)	<i>Leptodea leptodon</i> (Rafinesque)	<i>P. (C.) compressum</i> Prime
<i>Elliptio dilatatus</i> (Rafinesque)	<i>L. fragilis</i> (Rafinesque)	<i>P. (C.) fallax</i> Sterki
<i>Unio merus tetralasmus</i> (Say)	<i>Proptera laevis</i> (Lea)	<i>P. (C.) variabile</i> Prime
Subfamily Anodontinae	<i>P. purpurata</i> (Lamarck)	<i>P. (C.) adamsi</i> Stimpson (Hinkley, 1916)
Tribe Alasmidontini	<i>P. capax</i> (Green)	<i>P. (C.) nitidum</i> Jenyns (Burch, 1975)
<i>Lasmigona (Pterosyna) complanata</i> (Barnes)	<i>Carunculina parva</i> (Barnes)	<i>P. (Neopisidium) cruciatum</i> Sterki (Sterki, 1916)
<i>L. (Lasmigona) costata</i> (Rafinesque)	<i>C. texasensis</i> (Lea)	<i>P. (N.) punctatum</i> Sterki (Hinkley, 1916)
<i>Alasmidonta (Pressodonta) calceola</i> (Lea)	<i>C. glans</i> (Lea)	<i>P. (Pisidium) dubium</i> (Say) (Sampson, 1893, 1894; Hinkley, 1916)
<i>A. (Decurambis) marginata</i> Say	<i>Villosa arkansasensis</i> (Lea) ¹	
<i>Arcidens confragosus</i> (Say)	<i>V. lienosa</i> (Conrad)	
	<i>V. iris</i> (Lea)	

¹Endemic Interior Highlands species restricted to the Ozark Plateaus and some adjacent areas.

²Endemic Interior Highlands species restricted to the Ouachita Mountains.

³Endemic Interior Highlands species.

16. The first four listed *Pisidium* (see Table 3) are confirmed identifications (Gordon et al. 1980c). The remaining species are provisionally included on the authority of the references following the species name.

DISCUSSION

The geographical distribution of species richness in mollusks is similar to that of crayfish (see Hobbs in Pennak, 1978). The development of the North American molluscan fauna appears to have been centered in the highlands (Cumberland Region) of the southeastern United States (Walker, 1917; van der Schalie and van der Schalie, 1950); although, an old world derived fauna is prevalent west of the Continental Divide (Simpson, 1895; van der Schalie and van der Schalie, 1950). The number of species that have inhabited the Cumberland Region has been illustrated by Ortmann (1924, 1925) and van der Schalie (1939). Walker (1917), van der Schalie and van der Schalie (1950), Johnson (1978), and Gordon et al. (1980b) have shown that the Cumberland Region and the Interior Highlands are geologically related, have been physically connected, and were influenced by similar factors (e.g. geological uplifts, isolation, geographical shift of river systems). The past connection with the Cumberland Region, the geological age of the Interior Highlands, and their function as a biological refugium during the Cretaceous and Pleistocene (Gordon et al. 1980b) have resulted in the development of a diverse molluscan fauna in Arkansas. This fauna is distinguished by endemic species and relict populations.

The primary component of the present Arkansas mollusks is from the Interior Basin or Mississippian fauna. Endemism of the Mollusca within the Interior Highlands has been somewhat limited when compared to the Cumberland Region. However, distinct trends are represented for several groups. Fifteen terrestrial gastropods have been listed as endemic (see Table 1). Of these, the majority of species are from the family Polygyridae and are primarily restricted to the Ouachita Mountains. Species, such as *Masodon magazinensis* and *Stenotrema pilsbryi*, are restricted to single mountains (e.g. Mt. Magazine and Rich Mountain, respectively). The Hydrobiidae contains most of the endemic aquatic Gastropoda. In the Ozark Plateaus, several endemic hydrobiids have adapted to cave environments (Hubricht, 1950, 1979; Nordstrom et al. 1977). The endemic, epigeal hydrobiids appear to be known only from their type localities. At present, no endemic sphaeriids are known. However, a varied endemic unionid fauna has persisted, primarily within the Lampsiliinae. This is fairly consistent with the Cumberland fauna; although, to a lesser degree. The Cumberland association is supported by the endemic *Fusconia ozarkensis*, which Ortmann (1917) grouped with the Cumberlandian *F. barnesiana*. Other Cumberland affinities include relict populations of *Cumberlandia monodonta*, *Dyonmia triquetra*, *D. florentina* (formerly *Truncilla curtisii* Frierson and Utterback), *D. turgidula* (formerly *T. lefevrei* Utterback), and the presence of several pleurocerid species, including the endemic *Goniobasis potosiensis*. Other relicts, but of a southern origin, may include *Lampsilis streckeri* and *Arkansia wheeleri*. Needs for further study are indicated by recent faunal additions (Hubricht, 1979; Gordon et al. 1980c) and have been addressed by Gordon et al. (1980b).

ACKNOWLEDGMENTS

I would like to thank Dr. Fred G. Thompson of the Florida State Museum, University of Florida, for information concerning the taxonomy of the Hydrobiidae and Dr. Shi-Kuei Wu, University of Colorado Museum, for information regarding the revision of the Succineidae. Mr. Alan C. Buchanan, Missouri Department of Conservation, and Mr. D. Ronald Oesch, Glendale, Missouri, provided data from their collecting in Missouri. Also, I would like to thank Dr. Arthur V. Brown, Department of Zoology, University of Arkansas, for his advice and assistance.

LITERATURE CITED

- BAKER, F. C. 1911. The Lymnaeidae of North and Middle America, recent and fossil. Chi. Acad. Sci. Special Publ. 3:1-539.
- BAKER, F. C. 1928. The fresh water Mollusca of Wisconsin: Part 2. Pelecypoda. Bull. Wisc. Geol. Nat. Hist. Surv. 70:1-482.
- BALL, G. H. 1922. Variation in fresh-water mussels. Ecology 3:93-121.
- BURCH, J. B. 1962. How to know the eastern land snails. Wm. C. Brown Company Publisher, Dubuque. 214 pp.
- BURCH, J. B. 1975. Freshwater sphaeriacean clams (Mollusca: Pelecypoda) of North America. Malacological Publications, Hamburg. 96 pp.
- CALL, R. E. 1887. Descriptions of two new species of the genus *Unio*, from the Ozark region of Missouri. Proc. U.S. Natl. Mus. 10:498-500.
- CALL, R. E. 1895. A study of the Unionidae of Arkansas, with incidental references to their distribution in the Mississippi Valley. Trans. Acad. Sci. St. Louis 7:1-65.
- CLARKE, A. H. 1973. The freshwater molluscs of the Canadian Interior Basin. Malacologia 13:1-509.
- CLENCH, W. J. 1962. A catalogue of the Viviparidae of North America with notes on the distribution of *Viviparus georgianus* Lea. Occ. Pap. Moll. 2:261-287.
- CLENCH, W. J. and S. L. H. FULLER. 1965. The genus *Viviparus* (Viviparidae) in North America. Occ. Pap. Moll. 2:385-412.
- GOODRICH, C. 1939. Pleuroceridae of the Mississippi River Basin exclusive of the Ohio River system. Occ. Pap. Mus. Zool. Univ. Mich. 406:1-4.
- GORDON, M. E., A. V. BROWN and L. R. KRAEMER. 1980a. Mollusca of the Illinois River, Arkansas. Proc. Ark. Acad. Sci. 33(year 1979):35-37.
- GORDON, M. E., L. R. KRAEMER and A. V. BROWN. 1980b. Unionacea of Arkansas: historical review, checklist, and observations on distributional patterns. Bull. Am. Malac. Un. for 1979:31-37.
- GORDON, M. E., G. L. MACKIE and A. V. BROWN. 1980c. *Pisidium fallax* (Bivalvia: Sphaeriidae) in the southwestern Ozark Plateaus. Nautilus 94:121.
- HEARD, W. H. and R. H. GUCKERT. 1970. A re-evaluation of the recent Unionacea (Pelecypoda) of North America. Malacologia 10:333-355.
- HERRINGTON, H. B. 1962. A revision of the Sphaeriidae of North America (Mollusca: Pelecypoda). Misc. Publ. Mus. Zool. Univ. Mich. 118:1-74.
- HOBBS, H. H., JR. Figure 328, p. 474 in R. W. PENNAK. 1978. Fresh-water invertebrates of the United States, second edition. John Wiley and Sons, New York. 803 pp.
- HUBRICHT, L. 1950. The invertebrate fauna of Ozark caves. Bull. Natl. Speleological Soc. 12:16-17.
- HUBRICHT, L. 1972. The land snails of Arkansas. Sterkiana 46:15-17.

- HUBRICHT, L. 1979. A new species of *Amnicola* from an Ozark cave (Hydrobiidae). *Nautilus* 93:142.
- JOHNSON, R. I. 1978. Systematics and zoogeography of *Plagiola* (= *Dysnomia* = *Epioblasma*), an almost extinct genus of freshwater mussels (Bivalvia: Unionidae) from middle North America. *Bull. Mus. Comp. Zool.* 148:239-320.
- KRAEMER, L. R. 1970. The mantle flap in three species of *Lampsilis* (Pelecypoda: Unionidae). *Malacologia* 10:225-282.
- LEA, I. 1852. Observations on the genus *Unio*, etc.: vol. V. 62 pp.
- NORDSTROM, G. R., W. L. PFLIEGER, K. C. SADLER and W. H. LEWIS. 1977. Rare and endangered species of Missouri. Missouri Department of Conservation and U.S. Department of Interior Soil Conservation Service, Jefferson City. 129 pp.
- ORTMANN, A. E. 1912. Notes upon the families and genera of the naiades. *Ann. Carnegie Mus* 8:222-365.
- ORTMANN, A. E. 1917. A new type of the naiad-genus *Fusconaia*. Group of *F. barnesiana* Lea. *Nautilus* 31:58-64.
- ORTMANN, A. E. 1919. A monograph of the naiades of Pennsylvania. Part 3. Systematic account of the genera and species. *Mem. Carnegie Mus.* 8:1-384.
- ORTMANN, A. E. 1924. The naiad-fauna of the Duck River in Tennessee. *Am. Midl. Nat.* 9:18-62.
- ORTMANN, A. E. 1925. The naiad-fauna of the Tennessee River system below Walden Gorge. *Am. Midl. Nat.* 9:321-373.
- ORTMANN, A. E. and B. WALKER. 1912. A new North American naiad. *Nautilus* 25:97-100.
- ORTMANN, A. E. and B. WALKER. 1922. On the nomenclature of certain North American naiades. *Occ. Pap. Mus. Zool. Univ. Mich.* 112:1-75.
- PARMALEE, P. W. 1967. The freshwater mussels of Illinois. *Ill. State Mus. Pop. Sci. Ser.* 8:1-108.
- REHDER, H. 1932. Two new *Polygyras* from northern Arkansas. *Nautilus* 45:128-131.
- SAMPSON, F. A. 1893. Mollusca of Arkansas. *Nautilus* 7:33-35.
- SAMPSON, F. A. 1894. A preliminary list of the Mollusca of Arkansas (exclusive of the Unionidae). *Ann. Rept. Ark. State Geologist for 1891:181-199.*
- SIMPSON, C. T. 1895. The classification and geographical distribution of the pearly fresh-water mussels. *Proc. U.S. Natl. Mus.* 18:295-343.
- SIMPSON, C. T. 1914. A descriptive catalogue of the naiads or pearly freshwater mussels. Bryant Walker, Detroit. 1540 pp.
- STERKI, V. 1916. A preliminary catalogue of the North American Sphaeriidae. *Ann. Carnegie Mus.* 10:429-477.
- TE, G. A. 1975. Michigan Physidae, with systematic notes on *Physella* and *Physodon* (Basommatophora: Pulmonata). *Malac. Rev.* 8:7-30.
- VAN DER SCHALIE, H. 1938. The naiad fauna of the Huron River, in southeastern Michigan. *Misc. Publ. Mus. Zool. Univ. Mich.* 40:1-83.
- VAN DER SCHALIE, H. 1939. Additional notes on the naiads (freshwater mussels) of the lower Tennessee River. *Am. Midl. Nat.* 22:452-457.
- VAN DER SCHALIE, H. and A. VAN DER SCHALIE. 1950. The mussels of the Mississippi River. *Am. Midl. Nat.* 44:448-466.
- WALKER, B. 1917. The method of evolution in the Unionidae. *Occ. Pap. Mus. Zool. Univ. Mich.* 45:1-10.
- WHEELER, H. E. 1914. The Unione fauna of Cache River, with descriptions of a new *Fusconaia* from Arkansas. *Nautilus* 28:73-78.
- WHEELER, H. E. 1918. The Mollusca of Clark County, Arkansas. *Nautilus* 31:109-125.