## Journal of the Arkansas Academy of Science

Volume 31

Article 45

1977

# Chemical Composition of Carbonatites in Conway and Perry Counties of Arkansas

George H. Wagner University of Arkansas

Kenneth F. Steele University of Arkansas, Fayetteville

Follow this and additional works at: https://scholarworks.uark.edu/jaas

Part of the Geology Commons

### **Recommended Citation**

Wagner, George H. and Steele, Kenneth F. (1977) "Chemical Composition of Carbonatites in Conway and Perry Counties of Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 31, Article 45. Available at: https://scholarworks.uark.edu/jaas/vol31/iss1/45

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, uarepos@uark.edu.

#### 0 +0.4 Mnx 10 Six10 02 C -0.4 -0.4 -0.6 -0.8 C + Ca×102 +0.9 Cox102 -0.4 +0.6 -0.8 +0.3 0 -1.2 +0.2 -0.3 0 -0.6 0.8 0 Fex 10<sup>1</sup> -0.2 +1.0 N i × 10<sup>2</sup> -0.4 +0.5 -0.6 0 -0.8 -0.5 +6.0 -1.0 VXID Cr × 10 +4.0 +0.4 +0.2 +2.0 0 0 0104 0 10 -0.2 -2.0 0 0.3 0.6 0.9 1.2 0 0.3 0.6 0.9 1.2 1.5 1.5 WI WI

#### **Arkansas Academy of Science**

Figure 1. Change in element abundance in the weathered sample relative to the fresh sample calculated from data in Table 1. See text for definition of WI and \*. In the \*Cr and \*V plots the numbers correspond to sample numbers in Table 1. Circled points in the \*Cr plot represent minimum values.

KENNETH F. STEELE and EDWARD C. ROBISON, Department of Geology, University of Arkansas, Fayetteville, Arkansas 72701.

#### THE CHEMICAL COMPOSITION OF CARBONATITES IN CONWAY AND PERRY COUNTIES OF ARKANSAS

The geographical relationship of the Conway and Perry County Carbonatites to other igneous outcrops in Arkansas is outlined by Stone and Sterling (1964) in their summary of igneous rock occurrences in Arkansas. The Conway County outcrops were first described by Croneis and Billings in 1930. They have been called, proceeding historically, volcanic necks, sills, vein dikes, breccia and xenolithic dikes and classed as lamprophyres. The best and largest exposure, which is only a few square meters, in on the south side of the Arkansas River at Lock and Dam No. 9 near Morrilton, Arkansas. This material is a sill between shale and sandstone of probable Atokan age. With the help of Charles G. Stone of the Arkansas Geological Commission and R. R. Cohoon of Arkansas Technical University. Russellville, other exposures of similar looking material were located, and are summarized in Table 1.

The purpose of this study was to compare the chemical composition of the groundmass of these six outcrops of igenous rocks. Samples of the least weathered-looking material from each outcrop were obtained and a fresh, central portion of each, 1 cm thick X 4 cm X 10 cm was made with a water lubricated saw. Carefully avoiding large xenoliths which are plentiful. the dark gray groundmass was isolated and approximately 15 g were ground to 100 mesh by mortar and pestle. Samples were dissolved by HF and HC1 for chemical analyses. Samples were only partially soluble in HC1. The solutions were analyzed by a Perkin Elmer Model 303 Atomic Absorption Spectrophotometer using standard techniques (Anonymous, 1973). Carbon dioxide was determined by the titration method of Shapiro (1975).

Arkansas Academy of Science Proceedings, Vol. XXXI, 1977

121

#### **General Notes**

Analyses of the HF dissolved samples of the six outcrops are summarized in Table 2 and are compared with the well known Magnet Cove carbonatite, the Alno, Sweden carbonatite and the Murfreesboro kimberlite. The Morrilton Dam material has the greatest HCl solubility, 83.6%, and the Oppello Payne Farm material the least, 24.2%, with the other samples being intermediate. The differences are believed to be due to weathering, with the most soluble being the least weathered. The CO<sub>2</sub> contents parallel the calcium contents as shown in Table 2. Neglecting the badly weathered samples, Perry Mountain and Oppello Payne Farm, the other outcrop samples have similar compositions among themselves and compare well to the known carbonatites for both major and minor elements.

From the HF analyses of the whole groundmass and the HC1 solubles analyses, the composition of the HC1 insolubles were calculated. The following elements were more concentrated in the HC1 solubles: Ca. Fe, Sr, Mn, Zn, Ni and V. The following were more concentrated in the insolubles: Mg, Ti, Al, Na, K, Ba, Co, Cu and Li.

In Figures 1 and 2 the HF analyses are plotted versus the HC1 solubility of the samples for the major and minor elements, respectively. Weathering should decrease HC1 solubility. Assuming this, it is apparent from Figure 1 and 2 that the samples originally had similar compositions but experienced different degrees of weathering. As weathering proceeded, the content of the more soluble cations (alkaline earths) decreased and that of the refractory metal oxides (iron, aluminum, titanium, vanadium) either increased or stayed about constant. Silicification of the Perry Mountain B sample has caused even the refractory oxides to decrease in concentration.

We conclude that the igneous outcroppings are from the same or similar magma source and differ only in the degree of alteration. They are carbonatites befitting Heinrich's (1966) definition: "A carbonatite is here defined as a carbonate-rich rock of apparent magmatic derivation or descent."

#### LITERATURE CITED

ANON. 1973. Analytical methods for atomic absorption spectrophotometry, Perkin Elmer, Norwalk, Conn., U.S.A.

- CRONEIS, GAREY G., and MARLAND P. BILLINGS. 1930, Igneous rocks in Central Arkansas. In: Croneis, Carey (Editor), Geology of The Arkansas Paleozoic Area, Arkansas Geol. Survey Bull. 3, pp. 457.
- ERICKSON, R.L., and L.V. BLADE. 1963. Geochemistry and petrology of the alkali igneous complex at Magnet Cove, Arkansas: U.S. Geol. Survey Prof. Paper 425, 95 p.
- HEINRICH, E. WM. 1966. The geology of carbonatites. Rand McNally. Chicago, 555 p.
- MISER, H.D. and C.S. ROSS. 1923. Diamond-bearing peridotite in Pike County, Arkansas: U.S. Geol. Survey Bull. 735, p. 279-322.
- SHAPIRO, LEONARD. 1975. Rapid analyses of silicate, carbonate, and phosphate rocks — revised edition. U.S. Geol. Survey Bulletin 1401, 76 p.
- STONE, CHARLES G., and PHILIP J. STERLING. 1964. Relationship of igneous activity to mineral deposits in Arkansas. Arkansas Geological Comm. report, 56 p.

Name of Carbonatite	County	Section	Township	Range
Morrilton Dam	Conway	SWŁ,SWŁ,Sec.34	T6N	R17W
Oppello Dump	Conway	SW1, NE1, Sec. 1	T5N	R17W
Opello Boat Launch	Conway	NEL, NWL, Sec. 17	T5N	R16W
Opello Payne Farm	Conway	NWŁ,SEŁ,Sec.2	T5N	R17W
Perry Mt. Breccia	Perry	SW1, NE1, Sec. 3	T4N	R17W
Brazil Branch	Perry	SW1, SE1, Sec.28	T4N	R17W

Table 1. Location of Carbonatites in Conway and Perry Counties in Arkansas

Carbonatites	VT. 1							ppm									
	Ca	Mg	Fe	TI	Al	Na	ĸ	co2	Ba	Sr	Mn	Zn	Co	NI	Cu	v	LI
Morrilton Dam A	30.88	1.99	5.30	0.91	1.72	0.19	0.48	34.3	901	5714	3540	142	32	<7	30	277	27
Oppello Dump	29.78	2.98	6.15	0.74	1.68	0,44	0.43	32.7	1762	3305	2350	163	<5	5	6	164	24
Oppello Boat Launch	26.70	4.24	7.07	0.60	1.30	0.15	0.10	30.7	1939	2823	4780	190	19	21	5	147	5
Brazil Branch	24.36	2.58	6.92	0.64	3.06	1.07	2.23	28.8	1363	3884	4780	190	37	10	12	147	-11
Perry Mt. A	11.60	0.37	5.49	0.44	5.22	3.09	1.96	13.5	227	479	3230	136	30	36	17	130	4
Oppello Payne Farm	6.96	1.35	6.15	1.06	6.95	3.24	2.32	7.3	748	790	1920	182	38	58	24	202	9
Magnet Cove <sup>1</sup>	38.14	0.63	0.54	0.06	0.17	0.00	0.13	39.4	1000	-	5000	-	0	0	10	200	
Alnö, Sweden <sup>2</sup>	25.80	3.49	5.22	0.44	1.74	0.31	1.31	28.5	3600	3900	5300	-	17	8	2		
Kimberlite <sup>3</sup>	2.77	15.88	4.20	0.53	3.62	0.58	2.12	- 1	-	-	-	-		-	-	-	-

<sup>1</sup> analysis of sample L-304, Erickson and Blade (1963)

<sup>2</sup> from p 222 and 225, Heinrich (1966)

<sup>3</sup> from Miser and Ross, (1923)

Table 2. Groundmass Analyses of Carbonatites of Conway and Perry Counties. Arkansas (HF dissolved)

#### Arkansas Academy of Science Proceedings, Vol. XXXI, 1977



#### 123

#### Arkansas Academy of Science Proceedings, Vol. XXXI, 1977

Published by Arkansas Academy of Science, 1977