2018

Winter Surveys of *Cotinus obovatus* (American smoketree) in the Ozark Mountains

Gary R. Graves

National Museum of Natural History; University of Copenhagen, gravesg@si.edu

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Part of the [Botany Commons](#)

Recommended Citation

Graves, Gary R. (2018) "Winter Surveys of *Cotinus obovatus* (American smoketree) in the Ozark Mountains," *Journal of the Arkansas Academy of Science*: Vol. 72 , Article 15.

DOI: <https://doi.org/10.54119/jaas.2018.7201>

Available at: <https://scholarworks.uark.edu/jaas/vol72/iss1/15>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Winter Surveys of *Cotinus obovatus* (American smoketree) in the Ozark Mountains

Cover Page Footnote

I thank the Ozark National Forest (USDA) for issuing research permits (OZF-FW-FY17-03) and two anonymous reviewers for comments on the manuscript. I acknowledge the continuing support of the Alexander Wetmore Fund (Smithsonian Institution) and the Smoketree Trust.

Winter surveys of *Cotinus obovatus* (American smoketree) in the Ozark Mountains

G.R. Graves

Department of Vertebrate Zoology, MRC-116, National Museum of Natural History, Smithsonian Institution, P.O. Box 37012, Washington, D.C. 20013

Center for Macroecology, Evolution and Climate, University of Copenhagen, 2100 Copenhagen Ø, Denmark

Correspondence: gravesg@si.edu

Running Title: Winter Smoketree Surveys

Abstract

Cotinus obovatus (American smoketree) is a rare deciduous tree with a relictual distribution in southeastern North America. Efforts to map its fine-scale geographic distribution in the Ozark Mountains have been limited to the growing season when the distinctive blooming panicles and foliage facilitate detection in hardwood-cedar woodlands. I describe the physiognomic traits of leafless *C. obovatus* that permit effective population mapping in winter landscapes. Clumped growth and diagonally leaning stems facilitate detection at a distance. Bark texture, twig morphology, and sap odor confirm the identity of the tree at close range.

Introduction

Cotinus obovatus Raf. (Anacardiaceae: American smoketree), the sole member of the genus in the Western Hemisphere, has intrigued naturalists since Thomas Nuttall discovered it in 1819 on a bluff above the Grand River in eastern Oklahoma (Buckley 1881; Little 1942; Long 1970; Mohr 1882; Nuttall 1821; Palmer 1921; Sargent 1885, 1892; Weber 1990). Natural populations of this suffusively rare deciduous tree have been documented in 41 counties scattered in three disjunct regions: (1) Ozark Plateau in Arkansas and Missouri with a few scattered stations along bluffs of the Arkansas River and its tributaries in eastern Oklahoma, (2) southern Cumberland Plateau in northeastern Alabama and adjacent Tennessee and Georgia, and (3) the Edwards Plateau in west-central Texas. *Cotinus obovatus* is currently considered globally secure (G4) owing to protected populations on national forest lands in the Ozark Mountains of Arkansas and Missouri, but it is listed as threatened in Alabama (S2), Oklahoma (S2), and Tennessee (S2) and as critically imperiled (S1) in Georgia (NatureServe, 2017).

The center of abundance of *C. obovatus* on the

Ozark Plateau occurs in four counties that border the Arkansas-Missouri state line (Baxter, Marion, Taney, Ozark). Factors that restrict the tree's geographic distribution are unknown, but all well-documented sites occur on soils derived from calcareous bedrock. An effort to map its fine-scale geographic distribution in relation to the soil chemistry and geology of the Ozark Plateau has been ongoing for the past decade. Population mapping has traditionally been conducted during the growing season when the distinctive blooming panicles (late April through mid-May) and foliage (April-September) are easily detectable in roadside surveys (Davis and Graves 2016). However, dense understory foliage may conceal isolated specimens of *C. obovatus* in the interior of hardwood-cedar woodlands. If the tree can be identified at modest distances (~10–40 m) in leafless winter landscapes, detection efficiency during walking surveys may actually increase relative to growing season efforts. Here I describe the physiognomic traits that facilitate winter surveys of *C. obovatus*. These can be grouped into characters that form the search image necessary to detect trees at distance (growth form) and those that confirm identity at close range (bark, twigs, sap).

Methods

Study area

Winter surveys of *C. obovatus* were conducted 28 February–14 March 2017 and 4–20 February 2018 in the Sylamore Ranger District of the Ozark National Forest, centered on the Norfork and Norfork SE topographic quadrangles (7.5-minute series, U.S. Geological Survey) in Baxter County, Arkansas. This region is topographically complex (100–429 m), dissected by deep valleys that drain into the White River and its major tributary, the Buffalo River. Current land cover in the national forest is a patchwork of secondary hardwood forest, *Juniperus virginiana* (eastern redcedar), and *Pinus echinata* (shortleaf pine). Glades are relatively

Figure 1. Typical growth form of *Cotinus obovatus* (American smoketree) in the Ozark National Forest, Baxter County, Arkansas. Multiple stems emerge from a single root crown in many specimens.

rare and those that do occur are relatively small. Old-growth trees and elevational bands of timber that exhibit no evidence of logging are found on steep, rocky slopes. Surface rocks over most of the study area are composed of interbedded limestone, dolostone, and sandy

dolostone of the Everton Formation (middle Ordovician) which is overlain by thin beds of St. Peter sandstone. The highest ridges and knobs are capped by cherty limestones of the Boone formation, (Haley et al., 1993). Data reported in this study were obtained on off-road surveys aimed at documenting the geographic coordinates of witness specimens of *C. obovatus* at a 200 m grain size.

Data collection

I photographed specimens of *C. obovatus* ($n = 177$) to document growth form and stem count. Selected trees met the following criteria: (i) the base of the tree was unobscured by undergrowth; (ii) the number of stems (> 5 cm DBH) emerging from the root crown could be counted in the photograph; and (iii) the stems were mature enough to have attained the flaky bark characteristic of older trees. I walked around each tree until the principal axis of a focal stem, usually the thickest for multi-stemmed specimens, was perpendicular to the plane of the camera. From enlarged photographs, I measured the angle of the focal stem (from ground level) along its lower side with a protractor. Landmarks for angle measurement were stem base at ground level and a point on the stem ~ 1.5 m from the base. Only a single stem per specimen was measured. Summary statistics (mean $\pm$ standard deviation) are presented for stem angle and stem count.

Results

Growth form

Sargent (1892:3) offered a brief description of *C. obovatus* physiognomy: "A small tree, twenty-five to thirty-five feet in height, with a straight trunk occasionally twelve or fourteen inches in diameter, usually dividing, twelve or fourteen feet from the ground, into several erect stems which separate into wide-spreading, often slightly pendulous branches." Individuals of this diameter are rare in the Ozark Mountains, but more importantly, a majority of *C. obovatus* trees are composed of multiple stems that emerge from the root crown (Fig. 1).

In the sample of photographed trees (Fig. 2), the number of stems per root crown varied from 1 to 15 ($\bar{x} = 2.6 \pm 2.5$). Stems have a tendency to lean or sprawl diagonally from the base. Stem angle (from horizontal) varied from 9° to 90° ($\bar{x} = 62.7 \pm 18.6^\circ$; $n = 177$). Solitary stems tend to be more vertical ($\bar{x} = 68.2 \pm 16.6^\circ$; $n = 77$) than stems randomly chosen from multi-stemmed specimens ($\bar{x} = 58.4 \pm 18.9^\circ$; $n = 100$). In any event, diagonally leaning stems and clumped growth are

Winter Smoketree Surveys

Figure 2. Histogram of stem lean angle (90° = vertical) and number of stems per root crown of *Cotinus obovatus* (American smoketree) in the Ozark National Forest, Baxter Country, Arkansas. Y-axis represents the number of specimens in each bin.

hallmarks of mature *C. obovatus* in open and closed canopy hardwood-cedar woodland in the Ozark Mountains.

Bark

Standard floras (Mohr 1901; Steyermark 1963) of states that harbor populations of *C. obovatus* omit descriptions of bark, and the accounts and illustrations in field guides (Leopold *et al.* 1998; Samuelson and Hogan 2003; Sibley 2009; Simpson 1988) are too brief or idiosyncratic to be useful for distinguishing the bark of *C. obovatus* from that of other small forest trees with rough bark. Buckley (1881: 125) observed, "...trunks and larger limbs [of *C. obovatus*] coated with light gray and deeply-furrowed bark resembling the bark of the larger trees of the common sassafras (*S. officinale*).” Perhaps the most accurate description was penned soon after the rediscovery of the tree in Alabama in 1881 (Mohr 1882: 218), “The bark is rough, covered with a whitish gray epidermis of a deep chestnut-brown underneath, and exfoliating in oblong square scales of uniform size.”

In the Ozark Mountains, the smooth gray bark of young *C. obovatus* begins to crack and crenellate when

the stems reach a diameter of 4–7 cm, although some sprouts as large as 10 cm have immature bark. The bark of mature stems is relatively thin and fractured into rounded rectangular or oval scales that are usually 1.8–3.5 times longer than wide (Fig. 3). Scale tips are frequently curled or curved away from the stem. Some specimens possess imbricated bark scales with curved tips pointing downward. Scale size and shape tend to be relatively uniform within individuals but vary significantly among trees. The bark is pale to medium gray when dry and dark gray to black when wet. The diagnostic combination of bark texture and color distinguishes *C. obovatus* from other small trees with craquelure bark in the Ozark Mountains (e.g., eastern redbud, *Cercis canadensis*; flowering dogwood, *Cornus florida*; sassafras, *Sassafras albidum*; and gum bumelia, *Sideroxylon lanuginosum*). Large specimens of *C. obovatus* frequently exhibit cambial dieback, which can affect as much as 80% of the stem girth. Exposed wood weathers to grayish black. The crown foliage of many older specimens is nourished by a meandering band of cambium and phloem winding up the stem. The cause of cambial dieback is unknown.

Twigs

Sargent (1892: 3) provided a concise description of the twigs of *C. obovatus*, “The young shoots are purple at first, but soon become green; during the first winter they are bright red-brown and are covered with small white lenticular spots and marked by large prominent leaf-scars...The winter-buds are acuminate and an eighth of an inch long, and are covered with thin dark red-brown scales.” During the late winter in the Ozark Mountains, the reddish-brown color, moderate diameter (3–5 mm), and absence of knobby or obliquely projecting buds give twigs a smooth silhouette and distinctive appearance (Fig. 4) that help confirm the identity of *C. obovatus* at moderate distances.

Sap

Buckley (1881:3) noted that broken branches of *C. obovatus*, “emitted a yellowish sap, the odor of which was highly aromatic; to me very unpleasant.” Mohr (1882: 218) observed that the bark, when bruised, exuded “a resinous sap of a heavy, disagreeable terebinthinous odor.”

I found the pungent odor to be reminiscent of the sap of *Rhus copallinum* (winged sumac) and *R. glabra* (smooth sumac), both of which are widespread in the Ozarks, and nearly identical to that of *Cotinus coggygria* (Eurasian smoketree), which is occasionally used in urban landscaping in the Ozark region. Broken

twigs of *C. obovatus* are odorous, even in cool or cold weather (0–5° C). The odor persists for years in dead twigs. The aromatic constituents of *C. obovatus* sap

have not been analyzed but intensive analyses of *C. coggygia* have revealed that the wood and bark are rich in monoterpenes, sesquiterpenes, and phenols (Matić *et al.* 2016). As a matter of practice, all witness trees encountered during winter surveys in areas where *C. obovatus* trees occurred at low density were confirmed through the examination of growth form, bark, twigs, and sap odor.

Figure 3. Bark of mature stems of *Cotinus obovatus* (American smoketree).

Figure 4. Twigs of the American smoketree (*Cotinus obovatus*).

Winter Smoketree Surveys

Discussion

Winter survey efficiency has not been quantified although it certainly varies with topography, understory visibility, and the density of *Cotinus obovatus* trees. Once the search image has been formed, observers should be able to detect most if not all mature specimens within 20 m of the survey path at slow walking speed (1-3 km/hr). In relatively open understories, the clustered, leaning stems of *C. obovatus* may be apparent to the unaided eye at distances as great as 50 m. Binoculars are indispensable for scanning prospective specimens, and I never attempted a cross-country survey without them.

Acknowledgments

I thank the Ozark National Forest (USDA) for issuing research permits (OZF-FW-FY17-03) and two anonymous reviewers for comments on the manuscript. I acknowledge the continuing support of the Alexander Wetmore Fund (Smithsonian Institution) and the Smoketree Trust.

Literature cited

- Buckley SB.** 1881. *Rhus cotinoides*, Nutt. Proceedings of the Academy of Natural Sciences of Philadelphia 1881:125.
- Davis DR and GR Graves.** 2016. A new leafmining moth (*Cameraria cotinivora*, Lepidoptera: Gracillariidae) of the American smoketree (*Cotinus obovatus*). Proceedings of the Entomological Society of Washington 118:244-253.
- Haley BR, EE Glick, WV Bush, BF Clardy, CG Stone, MB Woodward, and DL Zachry.** 1993. Geologic Map of Arkansas. Arkansas Geological Commission and U.S. Geological Survey, 1:500,000 scale. 1 sheet.
- Leopold DJ, WC McComb, and RN Muller.** 1998. Trees of the central hardwood forests of North America. Timber Press, Portland (OR), 469 p.
- Little EL.** 1942. American smoketree (*Cotinus obovatus* Raf.), one of Oklahoma's rarest tree species. Proceedings of the Oklahoma Academy of Sciences 23:21-23.
- Long AW.** 1970. Ecological factors affecting the distribution of woody vegetation near the Arkansas River, Tulsa County with special reference to the Smoke-tree *Cotinus obovatus* [MS thesis]. Tulsa (OK): University of Tulsa.
- Matić S, S Stanić, M Mihailović, and D Bogojević.** 2016. *Cotinus coggygria* Scop.: An overview of its chemical constituents, pharmacological and toxicological potential. Saudi Journal of Biological Sciences 23:452-461.
- Mohr C.** 1882. *Rhus Cotinoides*, Nutt. Proceedings of the Academy of Natural Sciences of Philadelphia 34:217-220.
- Mohr C.** 1901. Plant Life of Alabama: An account of the distribution, modes of association, and adaptations of the flora of Alabama, together with a systematic catalogue of the plants growing in the state. USDA. Division of Botany (Washington, D.C.) 921 p.
- Natureserve.** 2017. American Smoketree (*Cotinus obovatus*). NatureServe Explorer: An online encyclopedia of life. Version 7.0. NatureServe, Arlington, VA. U.S.A. Accessed: 20 March 2017. http://explorer.natureserve.org/servlet/NatureServe?sourceTemplate=tabular_report.wmt&loadTemplate=species_RptComprehensive.wmt&selectedReport=RptComprehensive.wmt&summaryView=tabular_report.wmt&elKey=153598&paging=home&save=true&startIndex=1&nextStartIndex=1&reset=false&offPageSelectedElKey=155038&offPageSelectedElType=species&offPageYesNo=true&post_processes=&radiobutton=radiobutton&selectedIndexes=155038&selectedIndexes=153598
- Nuttall T.** 1821. Journal of travels into the Arkansas Territory, during the year 1819: with occasional observations of the manners of the aborigines: illustrated by a map and other engravings. Thos. H. Palmer (Philadelphia PA). 296 p.
- Palmer EJ.** 1921. The forest flora of the Ozark region. Journal of the Arnold Arboretum 2:216-232.
- Samuelson LJ and ME Hogan.** 2003. Forest trees: A guide to the southeastern and mid-Atlantic regions of the United States. Prentice Hall (Upper Saddle River NJ). 429 p.
- Sargent CS.** 1885. The woods of the United States: with an account of their structure, qualities, and uses: with geographical and other notes upon the trees which produce them. D. Appleton and Company (New York, NY). 203 p.
- Sargent CS.** 1892. Silva North America: A description of the trees which grow naturally in North America exclusive of Mexico. Vol. 3. Anacardiaceae-Leguminosae. Houghton, Mifflin and Company (Boston, MA and New York, NY). 141 p.
- Sibley DA.** 2009. The Sibley guide to trees. Knopf (New York, NY). 426 p.

- Simpson BJ.** 1988. A field guide to Texas trees. Texas Monthly Press (Austin, TX). 372 p.
- Steyermark J.** 1963. Flora of Missouri. Iowa State University Press (Ames, IA). 1728 p.
- Weber SF.** 1990. The population structure of *Cotinus obovatus* Raf., an ecologically narrow, endemic tree of cedar woodlands of the southern Cumberland Plateau [MS thesis]. Huntsville (AL): University of Alabama, Huntsville. 90 p.