2020

Energy Content of Seeds of Common Sunflowers (*Helianthus annuus*) in the Diet of Scaled Quail (*Callipepla squamata*) in Southeastern New Mexico

John L. Hunt

University of Arkansas at Monticello, huntj@uamont.edu

Matthew E. Grilliot

Auburn University--Montgomery, mgrillio@aum.edu

Troy L. Best

Auburn University, besttro@auburn.edu

Dixie Lozano-Lopez

University of Arkansas at Monticello, DXL20176@uamont.edu

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>

Emily R. Neilson

University of Arkansas at Monticello, em22641@uamont.edu
Part of the [Biology Commons](#), [Botany Commons](#), [Desert Ecology Commons](#), [Ornithology Commons](#),
and the [Other Nutrition Commons](#)

See next page for additional authors

Recommended Citation

Hunt, John L.; Grilliot, Matthew E.; Best, Troy L.; Lozano-Lopez, Dixie; Neilson, Emily R.; and Castillo, Isaac C. (2020) "Energy Content of Seeds of Common Sunflowers (*Helianthus annuus*) in the Diet of Scaled Quail (*Callipepla squamata*) in Southeastern New Mexico," *Journal of the Arkansas Academy of Science*: Vol. 74 , Article 11.

DOI: <https://doi.org/10.54119/jaas.2020.7401>

Available at: <https://scholarworks.uark.edu/jaas/vol74/iss1/11>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Energy Content of Seeds of Common Sunflowers (*Helianthus annuus*) in the Diet of Scaled Quail (*Callipepla squamata*) in Southeastern New Mexico

Authors

John L. Hunt, Matthew E. Grilliot, Troy L. Best, Dixie Lozano-Lopez, Emily R. Neilson, and Isaac C. Castillo

Energy Content of Seeds of Common Sunflowers (*Helianthus annuus*) in the Diet of Scaled Quail (*Callipepla squamata*) in Southeastern New Mexico

J.L. Hunt^{1*}, M.E. Grilliot², T.L. Best³, D. Lozano-Lopez¹, E.R. Neilson¹, and I.C. Castillo¹

¹University of Arkansas at Monticello, School of Mathematical and Natural Sciences, 397 University Drive, Monticello, AR 71656

²Auburn University—Montgomery, Biology Department, College of Arts and Sciences, P.O. Box 244023, Montgomery, AL 36124

³Auburn University, Department of Biological Sciences, 331 Funchess Hall, Auburn University, AL 36849-5414

*Correspondence: huntj@uamont.edu

Running Title: Energy Content of Seeds of Common Sunflowers in the Diet of Scaled Quail

Abstract

We analyzed the energy content of seeds of common sunflowers (*Helianthus annuus*) obtained from the crops of scaled quail (*Callipepla squamata*) collected from plains-mesa sand-scrub habitat in Eddy and Lea counties, New Mexico. Seeds were dried for 48 hours at 60°C to remove moisture and then analyzed for gross caloric value (i.e., energy content) in an oxygen bomb calorimeter. Energy content of seeds of common sunflowers from New Mexico was greater than that of many seeds previously reported from the diet of scaled quail and other granivorous birds and comparable to previous measurements of seeds of the same species made in Kansas.

Introduction

Conservation and management of birds requires knowledge of their food preferences, and of the energy content of their preferred foods. Feeding habits of scaled quail (*Callipepla squamata*) are well-studied (Lehman and Ward 1941; Schemnitz 1961; Ault 1981; Rollins 1981; Ault and Stormer 1983; Campbell-Kissock *et al.* 1985; Medina 1988), including several studies conducted in southeastern New Mexico (Davis and Banks 1973; Davis *et al.* 1975; Griffing and Davis 1976; Best and Smartt 1985; Hunt and Best 2001b). No study has measured the energy content of food of free-living scaled quail, although one study (Saunders and Parrish 1987) conducted in Kansas measured the assimilated energy of some potential food items by captive scaled quail. Studies have measured energy content of some known and potential food items of other birds such as mourning doves (*Zenaida macroura*—Schmid 1965; Shuman *et al.* 1988) in conjunction with determination of how well captive mourning doves metabolized various food items; some of the items measured are potential food for scaled

quail. No study of energy content of foods of scaled quail has been conducted with birds from sand-scrub habitat of New Mexico.

A study of feeding habits of scaled quail in southeastern New Mexico determined that seeds of common sunflowers (*Helianthus annuus*) made up the largest portion (14.3%) of the total mass of crop contents, and were present in 36.9% of the crops of mourning doves (Hunt and Best 2001b). Common sunflowers are also reported to be an important food item of other birds, such as northern bobwhites (*Colinus virginianus*—Hunt and Best 2001a) and mourning doves (Hunt 1999). Although energy content of seeds of wild common sunflowers from Kansas have been measured (Robel and Harper 1965), no such measurements have been conducted on seeds from sand-scrub habitat of New Mexico. We used an oxygen bomb calorimeter to determine the energy content of seeds of common sunflowers from Eddy and Lea counties in New Mexico.

Methods and Materials

Scaled quail were collected at the Waste Isolation Pilot Plant site in New Mexico in conjunction with long-term studies of lead poisoning of game birds (Best *et al.* 1992a; 1992b) and studies of feeding habits of game birds in southeastern New Mexico (Hunt 1999; Hunt and Best 2001a; Hunt and Best 2001b). Most of the study area is in eastern Eddy County, but it also extends into western Lea County. All scaled quail were collected in uncultivated, shinnery oak-honey mesquite (*Quercus havardii*-*Prosopis glandulosa*) habitat, part of the plains-mesa sand-scrub vegetation type (Dick-Peddie 1993). Several studies of the feeding ecology of scaled quail have been conducted in this area (Davis and Banks 1973; Davis *et al.* 1975; Griffing and Davis 1976; Best and Smartt 1985; Hunt and Best 2001b). There are several human-made stock

tanks on the study site, which is heavily grazed by cattle.

In late summer and autumn 1981, 58 scaled quail were collected by shooting as encountered. Collected birds were placed on ice within 10 minutes of shooting to minimize effects of post-mortem digestion (Dillery 1965; Farner 1960; Sedinger 1986); no effect of digestion on crop contents was observed. Crops were removed, placed into plastic vials, and frozen. Contents of crops were later thawed, separated by type of food, and placed into envelopes for drying. Food items were dried for 48 hours at 60°C to remove moisture. Food items were identified by comparison with samples of plants collected at the study site, and by using identification manuals (Davis 1993; Martin and Barkley 1961). Seeds used in this analysis were unbroken and unhulled.

Samples of seeds of *Helianthus annuus* were analyzed for gross caloric value (i.e., energy content) in an oxygen bomb calorimeter (Model 1341, Parr Instrument Company, Moline, Illinois). Samples of seeds from 13 individual scaled quail with crops that contained enough seeds for analysis were selected; each sample weighed 0.5-1.0 g. Seeds were combusted in the oxygen bomb; after combustion, the bomb was washed and bomb washings were titrated with sodium carbonate to allow adjustment of results for nitrate content. Results are reported in J/kg; kcal/g are given in parentheses for comparisons with previous studies.

Results

The 13 samples analyzed contained an average of 23.8 J/kg [5.7 kcal/g] — range, 21.3-24.7 J/kg [5.1-5.9 kcal/g]; standard deviation, 0.8 J/kg [0.2 kcal/g] (Table 1). This figure is similar to that previously obtained for *Helianthus annuus* in Kansas (Robel and Harper 1965) and greater than many tested food items in the diet of other seed-eating game birds.

Discussion

Although no study has demonstrated that scaled quail are selective in food choices, other species of granivorous birds, including quail, are known to be selective (Schmid 1965; Conley and Blem 1978; Shuman *et al.* 1988; Hayslette and Mirarchi 2001; Larson *et al.* 2012), although criteria for their selection are imperfectly understood. Among suggested criteria are shape, color, fiber content (Conley and Blem 1978), protein content (Larson *et al.* 2012), nutrient content (Hayslette and Mirarchi 2001), and energy

Table 1. Gross caloric value (energy content) of seeds of common sunflowers (*Helianthus annuus*) in the crops of scaled quail (*Callipepla squamata*) collected from Eddy and Lea counties, New Mexico, summer and autumn, 1981.

<u>Sample No.</u>	<u>Energy in J/kg (kcal/g)</u>
SQ002-81	23.8 (5.7)
SQ003-81	24.3 (5.8)
SQ008-81	24.7 (5.9)
SQ009-81	23.0 (5.5)
SQ015-81	23.4 (5.6)
SQ016-81	23.4 (5.6)
SQ017-81	23.0 (5.5)
SQ020-81	23.0 (5.5)
SQ021-81	24.7 (5.9)
SQ029-81	23.8 (5.7)
SQ031-81	21.3 (5.1)
SQ041-81	24.7 (5.9)
SQ042-81	24.7 (5.9)

content (Schmid 1965; Shuman *et al.* 1988). Although no study has demonstrated that scaled quail preferentially select common sunflowers over other food items, Davison (1958) categorized common sunflowers as a “choice” food plant for northern bobwhites, meaning it was digestible, nutritious, and readily eaten when encountered.

Our study demonstrates that seeds of common sunflowers have an energy content comparable to or greater than food items from previous studies. For example, Robel and Harper reported an average of 24.7 J/kg (5.9 kcal/g) for seeds of common sunflowers, and 23.0 J/kg (5.5 kcal/g) for seeds of giant ragweeds (*Ambrosia trifida*) collected in Kansas. A study of potential food items for greater prairie-chickens (*Tympanuchus cupido*—Heffron and Parrish 2005) listed 14 different commercial feeds and seeds that had a lower energy content than that which we measured for common sunflowers; the greatest energy content in that study was for hulled domestic sunflower-seed chips (23.0 J/kg [5.5 kcal/g]). A study of seeds of Texas doveweeds (*Croton texensis*) in crops of mourning doves conducted at the same study site as the current study (Hunt *et al.* 2019) reported an average energy content of 21.8 J/kg (5.2 kcal/g). In a study of 9 food items collected from crops of mourning doves in North Dakota, Schmid (1965) found only 2 that had a greater energy content—seeds of flax (*Linum usitatissimum*, 26.4 J/kg [6.3 kcal/g]) and field mustard

Energy Content of Seeds of Common Sunflowers in the Diet of Scaled Quail

(*Sinapis arvensis*, 25.1 J/kg [6.0 kcal/g]). Other seeds tested from North Dakota had less energy content; examples include wild plants such as green foxtail (*Setaria viridis*, 18.4 J/kg [4.4 kcal/g]) and cultivated crops such as corn (*Zea mays*, 17.1 J/kg [4.1 kcal/g]) and wheat (*Triticum aestivum*, 16.7 J/kg [4.0 kcal/g]). Likewise, Shuman *et al.* (1988) tested 8 varieties of seeds that were considered to be potential food items for mourning doves in Kansas, and found only 1 that had greater energy content—thistle (*Cirsium*, 25.9 J/kg [6.2 kcal/g]). Other seeds analyzed in Kansas had less energy content; examples include Maximilian sunflower (*Helianthus maximiliani*, 23.4 J/kg [5.6 kcal/g]), proso millet (*Panicum millaceum*, 18.8 J/kg [4.5 kcal/g]), and timothy (*Phleum pratense*, 19.7 J/kg [4.7 kcal/g]).

Common sunflowers and other sunflowers are associated with disturbance (Whitson *et al.* 1999). Much of southeastern New Mexico is heavily grazed by cattle, and much of the landscape has been highly modified by usage for extraction of petroleum and natural gas (Hunt 2004), so that common sunflowers grow abundantly. Abundance of common sunflowers, coupled with the relatively great energy content of the seeds, helps explain their prevalence in the diet of scaled quail (Hunt and Best 2001b) and other granivorous birds of the area.

Acknowledgments

The authors thank A. L. Williams and J. M. Bramlett for technical support. Funding for equipment was provided by Arkansas INBRE and by the School of Mathematical and Natural Sciences at the University of Arkansas at Monticello. Thanks also to Marvin Fawley for help in procuring funding.

Literature Cited

- Ault SC.** 1981. Food selection by scaled quail in Northwest Texas. [MS thesis]. Lubbock (TX): Texas Tech University. 75pp.
- Ault SC and FA Stormer.** 1983. Seasonal food selection by scaled quail in Northwest Texas. *The Journal of Wildlife Management* 47:222-228.
- Best TL, TE Garrison, and CG Schmitt.** 1992a. Availability and ingestion of lead shot by mourning doves (*Zenaida macroura*) in southeastern New Mexico. *The Southwestern Naturalist* 37:287-292.
- Best TL, TE Garrison, and CG Schmitt.** 1992b. Ingestion of lead pellets by scaled quail and bobwhite quail in southeastern New Mexico. *The Texas Journal of Science* 44:99-107.
- Best TL and RA Smartt.** 1985. Foods of scaled quail (*Callipepla squamata*) in southeastern New Mexico. *Texas Journal of Science* 37:155-162.
- Campbell-Kissock L, LH Blankenship, and JW Stewart.** 1985. Plant and animal foods of bobwhite and scaled quail in Southwest Texas. *The Southwestern Naturalist* 30:543-553.
- Conley JB and CR Blem.** 1978. Seed selection by Japanese quail, *Coturnix coturnix japonica*. *The American Midland Naturalist* 100:135-140.
- Davis CA and RL Banks.** 1973. Some food habits of scaled quail in southeastern New Mexico. New Mexico State University Agricultural Experiment Station Research Report 270:1-5.
- Davis CA, RC Barkley, and WC Haussamen.** 1975. Scaled quail foods in southeastern New Mexico. *The Journal of Wildlife Management* 39:496-502.
- Davis LW.** 1993. Weed seeds of the Great Plains. University Press of Kansas, Lawrence, KS. 145p.
- Davison VE.** 1958. A summary and reclassification of bobwhite foods. *The Journal of Wildlife Management* 22:437-438.
- Dick-Peddie WA.** 1993. New Mexico vegetation: past, present, and future. University of New Mexico Press, Albuquerque (NM). 244p.
- Dillery DG.** 1965. Post-mortem digestion of stomach contents in the savannah sparrow. *The Auk* 82:281.
- Farner DS.** 1960. Digestion and the digestive system. In: Marshall AJ, editor. *Biology and comparative physiology of birds*. Academic Press (NY). p 411-451.
- Griffing JP and CA Davis.** 1976. Comparative foods of sympatric scaled quail and mourning doves. *The Southwestern Naturalist* 21:248-249.
- Hayslette SE and RE Mirarchi.** 2001. Patterns of food preferences in mourning doves. *The Journal of Wildlife Management* 65:816-827.
- Heffron MB and JW Parrish, Jr.** 2005. Apparent metabolizable energy of seeds in greater prairie-chickens. *Transactions of the Kansas Academy of Science* 108:93-98.
- Hunt JL.** 1999. Dietary overlap and potential competition among scaled quail, mourning doves, and northern bobwhites in southeastern New Mexico [MS thesis]. Auburn (AL): Auburn University. 72p.

- Hunt JL.** 2004. Investigation into the decline of populations of the lesser prairie-chicken (*Tympanuchus pallidicinctus* Ridgway) in southeastern New Mexico [dissertation]. Auburn (AL): Auburn University. 294p.
- Hunt JL and TL Best.** 2001a. Foods of northern bobwhites (*Colinus virginianus*) in southeastern New Mexico. The Southwestern Naturalist 46:239-243.
- Hunt JL and TL Best.** 2001b. Foods of scaled quail (*Callipepla squamata*) in southeastern New Mexico. The Texas Journal of Science 53:147-156.
- Hunt JL, ME Grilliot, TL Best, CS Deen, D Lozano-Lopez, ER Neilson, and TR Schlegel-Ridgway.** 2019. Energy content of seeds of Texas doveweed (*Croton texensis*) from the diet of mourning doves (*Zenaidura macroura*) from southeastern New Mexico. Journal of the Arkansas Academy of Science 73:18-20.
- Larson JA, TE Fulbright, LA Brennan, F Hernandez, and FC Bryant.** 2012. Selection of seeds of an exotic and three native grasses by northern bobwhites (*Colinus virginianus*). The Southwestern Naturalist 57:319-322.
- Lehmann VW and H Ward.** 1941. Some plants valuable to quail in southwestern Texas. The Journal of Wildlife Management 5:131-135.
- Martin AC and WD Barkley.** 1961. Seed identification manual. University of California Press, Berkeley, California. 221p.
- Medina AL.** 1988. Diets of scaled quail in southern Arizona. The Journal of Wildlife Management 52:753-757.
- Robel RJ and WR Harper.** 1965. Energy content and retention by ragweed and sunflower seeds during fall and winter. Transactions of the Kansas Academy of Science 68:401-405.
- Rollins D.** 1981. Diets of sympatric bobwhite and scaled quail in Oklahoma. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies 35:239-249.
- Saunders DK and JW Parrish.** 1987. Assimilated energy of seeds consumed by scaled quail in Kansas. The Journal of Wildlife Management 51:787-790.
- Schenmitz SD.** 1961. Ecology of the scaled quail in the Oklahoma Panhandle. Wildlife Monographs 8:1-47.
- Schmid WD.** 1965. Energy intake of the mourning dove *Zenaidura macroura marginella*. Science, New Series 150:1171-1172.
- Sedinger SD.** 1986. Biases in comparison of proventricular and esophageal food samples from cackling Canada Geese. The Journal of Wildlife Management 34:739-746.
- Shuman TW, RJ Robel, AD Dayton, and JL Zimmerman.** 1988. Apparent metabolizable energy content of foods used by mourning doves. The Journal of Wildlife Management 52:481-483.
- Whitson TD, LC Burrill, SA Dewey, DW Cudney, BE Nelson, RD Lee, and R Parker.** 1999. Weeds of the West. University of Wyoming, Jackson Hole (WY). 630 pp.