

2021

## Remote Ornithology: Studying Nesting Behavior of Bermuda Petrels via Live Webcam

Ragupathy Kannan

University of Arkansas–Fort Smith, ragupathy.kannan@uafs.edu

Bala Chander

Jack Jackson

University of Arkansas–Fort Smith, Jack.Jackson@uafs.edu

Tara Chander

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Part of the [Biodiversity Commons](#), and the [Education Commons](#)

---

### Recommended Citation

Kannan, Ragupathy; Chander, Bala; Jackson, Jack; and Chander, Tara (2021) "Remote Ornithology: Studying Nesting Behavior of Bermuda Petrels via Live Webcam," *Journal of the Arkansas Academy of Science*: Vol. 75 , Article 18.

DOI: <https://doi.org/10.54119/jaas.2021.7512>

Available at: <https://scholarworks.uark.edu/jaas/vol75/iss1/18>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact [scholar@uark.edu](mailto:scholar@uark.edu).

## Remote Ornithology: Studying Nesting Behavior of Bermuda Petrels via Live Webcam

R. Kannan<sup>1\*</sup>, B. Chander<sup>2</sup>, J.L. Jackson II<sup>1</sup>, and T. Chander<sup>3</sup>

<sup>1</sup>University of Arkansas – Fort Smith, Fort Smith, AR 72913, U.S.A.

<sup>2</sup>501 Quaker Road, Chappaqua, NY 10514, U.S.A.

<sup>3</sup>Horace Greeley High School, Chappaqua, NY 10514, U.S.A.

\*Corresponding author: Ragupathy.Kannan@uafs.edu

Running Title: Webcam Study of Nesting Bermuda Petrels

The Bermuda Petrel *Pterodroma cahow* (hereafter, Cahow) is one of the rarest seabirds in the world (Fitzpatrick 2019; Brinkley and Sutherland 2020). After being presumed extinct for 300 years, the species was rediscovered in 1951 (Murphy and Mowbray 1951). Since then, an aggressive restoration program has increased the population (Madeiros *et al.* 2012; Brinkley and Sutherland 2020). A relocation scheme using artificial concrete nest burrows has succeeded in re-establishing a breeding population in Nonsuch Island, off the main island of Bermuda (Carlile *et al.* 2012).

Incubation behavior of *Pterodroma* petrels is difficult to study because of their burrowing and nocturnal behavior (Warham 1990). Although a lot of information is available about petrel breeding biology in general, little is known about the behavior of incubating parents. The advent of webcams has afforded an opportunity to fill this gap in knowledge.

In 2011, Nonsuch Expeditions, a Bermuda-based tour company, installed a live web camera inside an artificial nest burrow in Nonsuch Island, Bermuda, to livestream infrared video and audio. In 2016, they collaborated with the Bird Cams project at the Cornell Lab of Ornithology, resulting in 20 million minutes of footage from three seasons (Cornell Lab Bird Cams 2019). Jeremy Madeiros, Senior Conservation Officer (Terrestrial), in Bermuda's Department of Environment and Natural Resources, made periodic health checks of the nest and posted public updates on Twitter or YouTube (on egg and parental mass, etc.). In this note, we supplemented our own observations with those updates. We also included some observations from a second nest nearby with a webcam.

We observed the Cahow nest (Colony A, nest #831) via webcam for 167.3 hours in 2019, noting behaviors and involuntary movements of parents incubating a single egg. This is the first time an entire study has been done from remote via Webcam. Both parents were moved to the larger and more elevated

Nonsuch Island by biologists in 2006 as nearly fledged nestlings from two separate nesting islands nearby. This parental pair has been together since at least 2009. They had produced a nestling successfully for 5 years in a row since 2014 (J. Madeiros via video posted 21 March 2019). The website provided 4 hours of recording at any given time, enabling us to backtrack and make up to 4 hours of observation per access of camera feed. We monitored the nest for all but 9 days of the 55-day observation period. We missed those 9 days due to time conflicts. To ensure a balanced round-the-clock coverage, we monitored our coverage of each hour of the 24-hr clock throughout the study. We coordinated our observation efforts to cover all hours of the day and night. Although some time periods of the 24-hr clock were opportunistically covered better (e.g., 1000-1100 hrs Bermuda time) than others (0200-0300 hrs Bermuda time), we accomplished our goal to observe the nest at all times of the day and night. The percentage of total time observed ranged from 2% at 0200 h to 7% at 1100 h (Bermuda time) (Fig. 1). Almost equal time was spent observing the nest during day (81.4 hours) and night (85.9 hours).

Sexes look alike in Cahows, making sex identification via the webcam feed impossible. So, we based sex identifications on online postings made by Jeremy Madeiros. He reported using external cloacal examination to sex the birds within three weeks of egg-




Figure 1. Time of observations.




Figure 2. Sex of incubating parent vs. percentage of observation time.

laying. Outside this period, he found that nesting birds consistently showed a significant difference in mass and bill length, which are greater in adult males than adult females (as cited as a personal communication in Brinkley and Sutherland 2020).

We tracked the relative roles played by each parent in incubation (Fig. 2). Both parents contributed equally to incubation. Each was observed an equal proportion of observation time when they were solo in nest. The nest was left unattended only 1.5% of the observed time (Fig. 2).

The female (weighing 359g) laid a single egg (weighing 59g) on 10 January 2019. Our observations started on 11 January. An onsite health check on 17 January recorded the male parent's mass as 397g. Our observations were terminated on 6 March on the 55th day, 5 days after the known incubating time of 50 days from the same parents from the 2018 season. The mean incubation period reported for Cahows is 52.4 days (Warham 1990) and 53 days (Madeiros *et al.* 2012). The first Nonsuch Island chick hatched on 3 March 2019, in another nest. A health check of our nest on 6 March showed that the male had lost 85g since the last check, down to 312g. On 11 March 2019, the 60th day since laying, the egg was examined and found to be not viable. The parent continued incubation. Cahows are known to incubate for "up to a month" after failed incubation (J. Madeiros via video posted 21 March 2019).

We tracked several parental incubation behaviors (with codes used and percentage of observed time in parentheses). Sedentary behaviors of resting (R, 56%) and sleeping with head tucked back (S, 31%) accounted for most of the observed behaviors. We categorized behavior as "resting" when parents' head was upright with eyes open. Wheelwright and Boersma (1979) found that incubating Fork-tailed Storm Petrels spend majority of time sleeping, "often tucking bill under scapulars". Warham (1990) indicated that petrels

might sleep with head erect but eyes covered by nictitating membrane. "Comfort movements" (Warham 1990), i.e., preening (P, 5%) and nest maintenance (N, 3%) were also observed. Nest maintenance (rearranging fibers and/or digging ground) was observed 153 times with a mean time of 2.06 minutes per observation. Vigorous nest maintenance of 4–13.5 minutes was observed 22 times. Both parents were observed together in 21 different intervals. During 19 of these, allopreening (PA) was observed constituting 8% of the time while both parents were together. Moving (M) and shuffling around occupied only 1% of observed time. In addition, we noted the following other (O) less commonly observed behaviors: "yawning", wing stretching, head scratching, and adjusting egg with bill, together accounting for 0.5% of observed time. We counted 102 "yawns" (parent momentarily opening mouth wide), 42 wing stretches (parent extending one wing), 45 head scratches (all done over wing), and 41 egg adjustments (parent touching egg with bill to adjust its position) during the observation period. On 5 occasions the pair called to each other while both were in the nest, and on 1 occasion apparent copulation was observed. The attentive period, defined as proportion of time spent incubating was 98.5%; i.e. both parents were away (A) with the egg unattended for only 1.5% of observed time (Fig. 3). This high attentiveness is supported by studies on petrels in general (Warham 1990). We found no evidence of nest-ventilating behavior, wherein parent rises to its feet to expose egg (Grant *et al.* 1982).

Orientation of incubating parent on the nest was noted breaking directions down into 12 equal sectors as in the numbers on a clock, with the camera at 6:00 and 12:00 facing directly away from the camera. The camera was mounted on the side, facing so that the entrance was located at 11:00. The incubating parent




Figure 3. Behaviors of incubating parent as percentage of observation time (R=Resting, S=Sleeping with head tucked back, P=Preening, N=Nest maintenance, A=Away, M=Moving, PA=Allopreening, O=Other)

## Webcam Study of Nesting Bermuda Petrels

faced the entrance of the nesting burrow, orienting within  $30^\circ$  of the opening, 49% of the time. The exact opposite orientation was also common, with the parent facing within  $30^\circ$  of opposite the opening 28% of the time. Orientation perpendicular to the opening was rare (Fig. 4). These observations are supported by Warham (1990), who found that, generally, burrowing petrels face the entrance, apparently to greet a partner or deter intruders.

The presence of a rigidly stationary camera gives the opportunity to observe behaviors and involuntary movements that would be impossible to obtain otherwise. For the first time, breathing rates and headshaking rates were quantified in a seabird. We opportunistically quantified breathing rates (in breaths per minute) of the incubating parent for 189 times during the observation period, by counting the rhythmic heaving movements of the body for a minute. The breath rate of the male was significantly higher than that of the female (Table 1 and Fig. 5; 2-sample 2-tailed  $t$ -test,  $p < 0.0001$ ,  $t = 4.12$ ). A 95% confidence interval for the difference (male – female) in breath rate was [1.3, 3.7] breaths per minute.




Figure 4. Orientation of incubating parent.

Tube-nosed seabirds are known for their head-shaking behavior while at nest to remove salty secretions from their nostrils. When we observed these head shakes, we recorded the rate of the shakes. Male parent shook head significantly more than the female (Table 1 and Fig. 5; 2-sample 1-tailed  $t$ -test,  $p = 0.016$ ,  $t = 2.44$ ). A 95% confidence interval for the difference (male – female) in number of shakes was [1.4, 14.0] shakes per minute. It is not clear why this difference was observed, since both parents spent equal proportion of time incubating, and presumably, equal amounts of time foraging and getting exposed to salt. Excel was used to create graphs and calculate basic statistics.

There were 5 incubation shifts (parental “changing of the guard”) during our study. This agrees with typical frequency of incubation shifts in other petrels (Brown 1966; Warham *et al.* 1982; Thomas *et al.* 1983; Jouventin *et al.* 1985). Our observations, plus information provided by J. Madeiros on when the parents exchanged places, enabled us to track which days the parents were on the nest. The male was observed at the nest with the female absent for stretches of up to 13 days. Similarly, the female went up to 11 days without relief from her mate (Table 2). The female’s return on 22 January, after being away for 8 days, was a surprise. She was expected to be away for 2-3 weeks, leading to a tweet from J. Madeiros




Figure 5. Involuntary movements\*

Table 1. Involuntary movements

	Breathing Rates (breaths per minute)			Head Shakes (shakes per minute)		
	Male	Female	Combined*	Male	Female	Combined*
Number of Observations	62	65	189	49	37	96
Median	12	10	10	30	19	26.5
Mean	12.5	10.0	10.7	33.1	25.4	29.6
Standard Deviation	4.4	1.9	1.8	16.0	13.3	14.8

\*Combined includes observations when sex of the parent was unknown.



## Webcam Study of Nesting Bermuda Petrels

burrow post-season (17 May), when it was empty. For five years in a row since 2017, another webcam in the island recorded a Leach's Storm-Petrel (*Oceanodroma leucorhoa*) visiting and staying in an active Cahow nest burrow (Cornell Lab Bird Cams 2021). The list of intruders also includes an aggressive young Cahow prospecting for a nest site (Cornell Lab Bird Cams 2017).

This study would not have been possible without the collaboration between The Cornell Lab of Ornithology, Nonsuch Expeditions, and Jeremy Madeiros of Bermuda's Department of Environment and Natural Resources. Technology such as this opens many new avenues of data collection resulting in observations previously deemed impossible or impractical. Furthermore, making this video stream public allows for remote observations from around the world. The authors were able to perform these observations from Arkansas and New York without having to visit Bermuda. This study is especially unique since the authors have never seen a Cahow in person, yet they were able to make observations from the comforts of their offices or homes thousands of kilometers away. We encourage similar efforts to further the ability to do research of this kind and make the information widely available. It will also help surpass logistic and financial hurdles in conducting field studies and encourage the future use of webcams by teachers for collaborative citizen science investigations in their classrooms. The study was initiated as a special high school science project by one of the authors (TC).

### Acknowledgments

Two anonymous reviewers made useful comments on the manuscript.

### Literature Cited

- Boersma PD** and **MC Silva**. 2001. Fork-tailed Storm Petrel (*Oceanodroma furcata*). The Birds of North America, No. 569 (Poole A & F Gill, editors.). The Birds of North America, Inc. (Philadelphia, PA).
- Brinkley ES** and **K Sutherland**. 2020. Bermuda Petrel (*Pterodroma cahow*), version 2.0. In: Birds of the World (Schulenberg T, BK Keeney, and SM Billerman, editors). Cornell Lab of Ornithology (Ithaca, NY).
- Brown DA**. 1966. Breeding biology of the Snow Petrel *Pagodroma nivea* (Forster). Australian National Antarctic Research Expedition Science (Rep B) (1)

Zoology 89: 1-63.

- Carlile N, D Priddel, and J Madeiros**. 2012. Establishment of a new, secure colony of endangered Bermuda Petrel *Pterodroma cahow* by translocation of near-fledged nestlings. Bird Conservation International 22: 46-58.
- Cornell Lab Bird Cams**. 2017. Bermuda Petrel Intruder in the Burrow (edited) – Mar. 6, 2017. Youtube video [https://www.youtube.com/watch?v=KPe\\_KgSxeIc](https://www.youtube.com/watch?v=KPe_KgSxeIc). Accessed 12 August 2021.
- Cornell Lab Bird Cams**. 2019. AllAboutBirds.org/cahows. Accessed 8 April 2021.
- Cornell Lab Bird Cams**. 2021. Leach's Storm-Petrel "Stormy" visits Cahow family in nesting burrow – May 11, 2021. Youtube video <https://www.youtube.com/watch?v=8bdNj2EtG04>. Accessed 9 August 2021.
- Fitzpatrick J**. 2019. View From Sapsucker Woods: The comeback story of the Bermuda Petrel. Living Bird, Spring 2019.
- Grant SG, TN Pettit, H Rahn, GC Whittow, and CV Paganelli**. 1982. Water loss from Laysan and Black-footed Albatross eggs. Physiological Zoology 55:405-414.
- Jouventin P, JL Mouglin, JL Stahl, and H Weimerskirch**. 1985. Comparative biology of the burrowing petrels of the Crozet Islands. Notornis 32:157-220.
- Madeiros J, N Carlile, and D Priddle**. 2012. Breeding biology and population increase of the endangered Bermuda Petrel *Pterodroma cahow*. Bird Conservation International 22:35-45.
- Murphy RC** and **LS Mowbray**. 1951. New light on the Cahow, *Pterodroma cahow*. The Auk 68:266-280.
- Thomas G, JP Croxall, and PA Prince**. 1983. Breeding biology of the Light-mantled Sooty Albatross (*Phoebastria palpebrata*) at South Georgia. Journal of Zoology London 199:123-135.
- Warham J**. 1990. The Petrels: Their Ecology and Breeding Systems. Academic Press, London.
- Warham J, GJ Wilson, and BR Keeley**. 1982. The annual cycle of the Sooty Shearwater *Puffinus griseus* at the Snares Islands, New Zealand. Notornis 29:269-292.
- Wheelwright NT** and **PD Boersma**. 1979. Egg chilling and thermal environment of the Fork-tailed Storm Petrel (*Oceanodroma furcata*) nest. Physiological Zoology. 52:231-239.