

2021

Eimeria lancasterensis (Apicomplexa: Eimeriidae), Three Nematodes (Heligmosomoidea: Boehmiellidae, Heligmonellidae), and a Flea (Siphonaptera: Ceratophyllidae) from the Eastern Fox Squirrel, *Sciurus niger* (Rodentia: Sciuridae) in Arkansas

Chris T. McAllister

Eastern Oklahoma St. College, cmcallister@se.edu

John A. Hnida

Midwestern University, jhnida@midwestern.edu

Henry W. Robison

Retired, hwrobison@yahoo.com

Lance A. Durden

Georgia Southern University, ldurden@georgiasouthern.edu

Follow this and additional works at: <https://scholarworks.uark.edu/jaas>



Christopher Whipps
Part of the [Biology Commons](#)
SUNY-Syracuse, NY, cwhipps@esf.edu

Recommended Citation

McAllister, Chris T.; Hnida, John A.; Robison, Henry W.; Durden, Lance A.; and Whipps, Christopher (2021) "Eimeria lancasterensis (Apicomplexa: Eimeriidae), Three Nematodes (Heligmosomoidea: Boehmiellidae, Heligmonellidae), and a Flea (Siphonaptera: Ceratophyllidae) from the Eastern Fox Squirrel, *Sciurus niger* (Rodentia: Sciuridae) in Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 75 , Article 11.

DOI: <https://doi.org/10.54119/jaas.2021.7502>

Available at: <https://scholarworks.uark.edu/jaas/vol75/iss1/11>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in *Journal of the Arkansas Academy of Science* by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Eimeria lancasterensis (Apicomplexa: Eimeriidae), Three Nematodes (Heligmosomoidea: Boehmiellidae, Heligmonellidae), and a Flea (Siphonaptera: Ceratophyllidae) from the Eastern Fox Squirrel, *Sciurus niger* (Rodentia: Sciuridae) in Arkansas

Cover Page Footnote

The Arkansas Game and Fish Commission issued a Scientific Collecting Permit to CTM. We thank Drs. S.L. Gardner and G. Racz (HWML) for expert curatorial assistance, J.M. Kinsella (Missoula, MT) for nematode identifications, and L.M. Hardy (Ouachita Mountains Biological Station) for providing gratis housing and laboratory space for CTM. We also thank the Keck DNA Sequencing Facility at Yale University (New Haven, CT) for their assistance with DNA sequencing service.

***Eimeria lancasterensis* (Apicomplexa: Eimeriidae), Three Nematodes (Heligmosomoidea: Boehmiellidae, Heligmonellidae), and a Flea (Siphonaptera: Ceratophyllidae) from the Eastern Fox Squirrel, *Sciurus niger* (Rodentia: Sciuridae) in Arkansas**

C.T. McAllister^{1*}, J.A. Hnida², H.W. Robison³, L.A. Durden⁴, and C.M. Whipps⁵

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

²Department of Microbiology and Immunology, Midwestern University, Glendale, AZ 85308

³9717 Wild Mountain Drive, Sherwood, AR 72120

⁴Department of Biology, Georgia Southern University, Statesboro, GA 30458

⁵Environmental Biology, SUNY College of Environmental Science & Forestry, 1 Forestry Drive, Syracuse, NY 13210

*Correspondence: cmcallister@se.edu

Running Title: Parasites of Eastern Fox Squirrel

Abstract

In Arkansas, the eastern fox squirrel (*Sciurus niger*) is a common inhabitant of the state. Although information is available on ectoparasites of this host in Arkansas, little is known about the endoparasites of this squirrel. A single specimen from Montgomery County was examined and found to harbor the following: a coccidian (*Eimeria lancasterensis*), three nematodes, *Boehmiella wilsoni*, *Citellinema bifurcatum*, and *Sciurodendrium hassalli*, and a flea, *Orchopeas howardi*. We document these nematodes from an Arkansas *S. niger* for the first time, and add mensural and molecular information on *E. lancasterensis* from this host.

Introduction

The eastern fox squirrel, *Sciurus niger* (L., 1758) is the largest tree squirrel in the Western Hemisphere that occurs naturally in temperate forests over most of eastern North America (Hall 1981; Koprowski 1994). In Arkansas, *S. niger* is found statewide (Sealander and Heidt 1990). It inhabits a diversity of deciduous and mixed-forest habitats, but is more common in forest patches (Nixon and Hansen 1987). Fox squirrels feed heavily on tree seeds during much of the year (Koprowski 1994).

Although *S. niger* has been the subject of several studies of its coccidian parasites (Knipling and Becker 1935; Levine and Ivens 1965; Joseph 1972, 1973a, b, 1975; McAllister and Upton 1989; Spurgin and Hnida 2002; Motruik-Smith *et al.* 2009; Ozmen *et al.* 2009), there are no surveys reporting coccidia in any specimen from Arkansas.

Eastern fox squirrels have also been reported to be host of a suite of helminth parasites (Rausch and Tiner 1948; Flyger and Gates 1992). In Arkansas, Davidson (1976) examined some *S. niger* from the Ozarks in Stone County for parasites. There are no other reports of any helminth parasite from this host in the state. Here we report new records for parasites from a *S. niger* from the Ouachitas of Arkansas as well as include additional figures, mensural, and sequence data for a coccidian.

Materials and Methods

On 16 October 2020, an adult squirrel was hit and killed by an automobile on St. Hwy. 8, 3.2 km west of Black Springs, Montgomery County (34° 27' 16.29" N, -93° 46' 20.2872" W). It was opportunistically collected and immediately taken to the lab and processed for parasites. The pelage was brushed over a white enamel tray for ectoparasites. Any found were placed in a vial of 70% (v/v) ethanol and later cleared in 10% (w/v) potassium hydroxide, dehydrated through an ethanol series, further cleared in xylene, and slide-mounted in Canada balsam. A mid-ventral incision was made to expose the viscera and the gastrointestinal (GI) tract from the throat to anus was removed, rinsed in 0.9% (w/v) saline, and organs (including heart, liver, lungs, spleen, and kidneys) were placed in individual Petri dishes. Several 10 cm sections of the GI tract were cut, split lengthwise, and examined under a stereomicroscope for endoparasites. Feces from the rectum was collected and placed in 2.5% (w/v) potassium dichromate. A fecal flotation was accomplished with Sheather's sugar solution (sp. gr. 1.30). Nematodes were examined as temporary mounts

Parasites of Eastern Fox Squirrel

in glycerol.

For analysis of the DNA sequence of the *Eimeria* species, feces in 2.5% (w/v) potassium dichromate was sent to the Fish and Wildlife Disease Laboratory at SUNY-ESF. DNA was extracted using the Quick-DNA™ Fecal/Soil Microbe Miniprep Kit (Zymo Research Corp, Irvine, CA) with modifications described in Whipps *et al.* (2020). PCR was performed in 50 µL reaction volumes in Quick-Load® Taq 2X Master Mix (New England Biolabs, Ipswich, MA), 0.25 µM of each primer and 3 µL of template DNA. Overlapping fragments targeting the SSU ribosomal DNA were amplified with primers Eimeria1F (5'-GAT TCA TAG TAA CCG AAC GG) with 18R (Whipps *et al.*, 2003), and Eimeria2F (5'-GGG CAT TCG TAT TTA ACT GTC) with 18R. Amplifications were performed on a C1000™ Thermal Cycler (BioRad Laboratories, Hercules, CA) with initial denaturation at 95°C for 3 min, followed by 35 cycles of 94°C for 30 sec, 56°C for 45 sec, 68°C for 90 sec, and a final extension at 72°C for 7 min. Product amplification was evaluated by observation on a 1% (w/v) agarose gel and the remainder of the sample purified using the E.Z.N.A. Cycle Pure Kit (Omega Bio-Tek, Norcross, GA). DNA was quantified using a DNA spectrophotometer (NanoDrop Technologies Wilmington, Delaware). Sequencing used amplification primers with the ABI BigDye Terminator Cycle Sequencing Ready Reaction Kit v3.1, using the ABI3730xl Genetic Analyzer (Applied Biosystems, Foster City, CA). Sequences were assembled manually in BioEdit (Hall 1999) and identity analyzed by GenBank BLAST search.

A host photovoucher was deposited in the Eastern Oklahoma State College Collection, Idabel, OK. Voucher specimens of ectoparasites were deposited in the General Ectoparasite Collection in the Department of Biology at Georgia Southern University, Statesboro, GA. Endoparasites were deposited in the Harold W. Manter Laboratory (HWML) of Parasitology, University of Nebraska, Lincoln, NE, or samples were retained for molecular analyses.

APICOMPLEXA: EIMERIORINA: EIMERIIDAE

***Eimeria lancasterensis* Joseph, 1969** – Oocysts (Fig. 1A–C, HWML 216668) of this coccidian were being passed in feces. Oocysts ($n = 20$) were ellipsoidal, 23.5×14.3 ($18\text{--}29 \times 11\text{--}19$) µm, with a length/width ratio (L/W) of 1.6 (1.3–1.8). Bilayered wall was 1.4 (1.1–1.7) with a smooth, occasionally lightly pitted or sculptured, colorless to light yellow outer layer, ~2/3 total thickness; inner layer light yellow.




Figure 1. Sporulated oocysts of *Eimeria lancasterensis* from *Sciurus niger* from Montgomery County, Arkansas. A. Stieda body (SB) and sporocyst residuum (SR). B. Polar granule (PG) and posterior refractile body (PRB). C. Oocyst wall (OW) and sporocyst (SP). Scale bars = 10 µm.

Micropyle and oocyst residuum was absent but 1–3 sometimes bilobed polar granule(s) were present. Sporocysts ($n = 20$) were ellipsoidal, (L × W) 11.5×6.7 ($10\text{--}13 \times 6\text{--}8$) µm with an L/W ratio of 1.7 (1.4–2.2). Nipple-like Stieda body was present but subStieda and paraStieda bodies were absent. Sporocyst residuum was composed of various-sized granules forming a compact sphere, or a dense irregular mass located between and across the sporozoites, or a combination of both within the same sporocyst. Sporozoites (not measured) were elongate, anterior end tapered, posterior end rounded with a large, ellipsoidal posterior refractile body. The 1,472 nucleotide SSU DNA sequence for this specimen was submitted to GenBank (accession MZ831509). Our sequence was identical to a sequence from *E. lancasterensis* from eastern gray squirrels, *Sciurus carolinensis* Gmelin in Italy (GenBank accession KT360976) over 1,224 nucleotides.

This is one of the most prevalent coccidians infecting members of the rodent family Sciuridae. It has been reported previously from *S. niger* in Texas (McAllister and Upton 1989a), Nebraska (Spurgin and Hnida 2002), and Virginia and Wyoming (Motriuk-Smith *et al.* 2009). In addition, this coccidian has been reported from *S. carolinensis* from Italy (Hofmannová *et al.* 2016), from a red squirrel, *Sciurus vulgaris* (L.) in Turkey (Ozmen *et al.* 2009) and from *S. carolinensis* in Massachusetts (Joseph 1969, 1972), Florida (Forrester *et al.* 1977), Texas (McAllister and Kessler 2002), and Arkansas (McAllister and Kessler 2002). Although the latter authors reported *E. lancasterensis* in the state from *S. carolinensis*, no mensural data or photomicrographs were provided. Therefore, this is the first report of measurements, accompanying photomicrographs, and molecular data on *E. lancasterensis* from an Arkansas host.

NEMATODA: HELIGMOSOMOIDEA: BOEHMIELLIDAE

***Boehmiella wilsoni* Lucker, 1943.** – Two specimens (HWML 112234) were found in the stomach. *Boehmiella* spp. are principally characterized by having chitinized sheaths on the lateral and externo-dorsal rays of the bursa. They have short, complex unbranched spicules and females are didelphic. This nematode was described infecting *S. carolinensis* in Florida, Georgia, Minnesota, and West Virginia (Lucker 1943; Rausch and Tiner 1948). Coyner *et al.* (1996) reported it from *S. niger* from Florida, and Whitaker and Mumford (2009) from Indiana. Davidson (1976) reported *B. wilsoni* from *S. carolinensis* from Stone County, Arkansas. More recently, *B. wilsoni* was found in Deppe's squirrel, *Sciurus deppei* Peters in México (Falcon-Ordáz and García-Prieto 2004); in brown agouti, *Dasyprocta variegata* Tschudi in Bolivia (Mollericono *et al.* 2016); and in Ferreira's spiny tree-rat, *Mesomys hispidis* (Desmarest) in Brazil (Andrade-Silva *et al.* 2020). We document *B. wilsoni* in a *S. niger* from Arkansas for the first time.

TRICHOSTRONGYLOIDEA: HELIGMONELLIDAE

***Sciurodendrium hassalli* (Price, 1928).** – Approximately 30 specimens (HWML 112233) were found in the small intestine. Price (1928) originally described this nematode from *S. carolinensis* from Maryland. *Sciurodendrium* spp. are loosely coiled parasites and are characterized by having most of the cuticular ridges discontinuous and scalloped. Species are determined by the pattern of the bursal rays and females are monodelphic. The distribution of *S. hassalli* in sciurids is widespread. Chandler (1942) reported 100% prevalence in fox squirrels from eastern Texas, while Eckerlin (1993) found 50% prevalence in *S. niger* from Maryland and Virginia. It has also been reported from *S. niger* from Florida (Coyner *et al.* 1996), Ohio (Katz 1938) and Tennessee (Reiber and Byrd 1942). Davidson (1976) reported *S. hassalli* from *S. carolinensis* from Stone County. We document *S. hassalli* from an Arkansas eastern fox squirrel for the first time.

***Citellinema bifurcatum* Hall, 1916.** – Two specimens (HWML 112232) were recovered from the small intestine. The type host is the Wyoming ground squirrel, *Urocyon elegans* (Kennicott) (see Hall 1916). *Citellinema* spp. are tightly coiled parasites characterized by an asymmetrical bursa with a greatly reduced dorsal ray. The spicules are short (380–400 µm) and deeply bifurcated and females are didelphic. It is a common among sciurids where it occurs in

squirrels over a range from Colorado, Wyoming, and Saskatchewan, Canada to Maine (Reiber and Byrd 1942). This nematode has also been reported from *S. niger* from Florida (Coyner *et al.* 1996), Indiana (Whitaker and Mumford 2009), Tennessee (Reiber and Byrd 1942), and Ohio (Katz 1938). Davidson (1976) reported *C. bifurcatum* from *S. carolinensis* from Stone County. This nematode is reported from an Arkansas eastern fox squirrel for the first time.

ARTHROPODA: INSECTA: SIPHONAPTERA: CERATOPHYLLIDAE

***Orchopeas howardi* (Baker, 1895).** – a single female (L3851) was recovered. This flea is a common ectoparasite of sciurids, including *S. niger* (Whitaker *et al.* 1976; Lewis 2000). Schiefer and Lancaster (1970) and McAllister *et al.* (2013) reported *O. howardi* previously from *S. niger* from the Arkansas Ozarks. Other hosts from the state include *S. carolinensis*, southern flying squirrel, *Glaucomys volans* (L.), and raccoon, *Procyon lotor* (L.) (McAllister *et al.* 2017). This flea has been reported to transmit North American strains of the causative agent of sporadic epidemic typhus (*Rickettsia prowazekii*), which is maintained enzootically in flying squirrel populations (McDade 1987). Human cases of this disease have been serologically confirmed and recorded in Arkansas (McDade 1987). We report *O. howardi* from a host from the Ouachita uplands of the state for the first time.

In conclusion, we document, for the first time, three nematodes from a *S. niger* from Arkansas. Two of these, *S. hassalli* and *C. bifurcatum*, which have direct life cycles, are proposed to be core species of *S. niger* (Kinsella 1991) and we concur. Although only a single *S. niger* was examined herein it yielded these new records as well as extra mensural and molecular data on the coccidian, *E. lancasterensis*. Additional eastern fox squirrels in Arkansas should be examined for parasites from the southern and eastern parts of its range in the state.

Acknowledgments

The Arkansas Game and Fish Commission issued a Scientific Collecting Permit to CTM. We thank Drs. S.L. Gardner and G. Racz (HWML) for expert curatorial assistance, J.M. Kinsella (Missoula, MT) for nematode identifications, and L.M. Hardy (Ouachita Mountains Biological Station) for providing gratis housing and laboratory space for CTM. We also thank

Parasites of Eastern Fox Squirrel

the Keck DNA Sequencing Facility at Yale University (New Haven, CT) for their assistance with DNA sequencing service.

Literature Cited

- Andrade-Silva BE, RV Vilela, EJ Lopes-Torres, SF Costa-Neto, and A Maldonado.** 2020. *Boehmiella wilsoni* (Nematoda, Heligmosomoidea, Boehmiellidae fam. nov.), found in Amazonian rodents. *International Journal for Parasitology: Parasites and Wildlife* 13:119–129.
- Chandler AC.** 1942. Helminths of tree squirrels in southeast Texas. *Journal of Parasitology* 28:135–140.
- Coyner DF, JB Wooding, and D Forrester.** 1996. A comparison of parasitic helminths and arthropods from two subspecies of fox squirrels (*Sciurus niger*) in Florida. *Journal of Wildlife Diseases* 32:492–497.
- Davidson WR.** 1976. Endoparasites of selected populations of gray squirrels (*Sciurus carolinensis*) in the Southeastern United States. *Proceedings of the Helminthological Society of Washington* 43:211–217.
- Eckerlin RP.** 1993. Helminth parasites from two populations of the Delmarva fox squirrel (*Sciurus niger cinereus*) in Maryland and Virginia. In: Moncrief ND, Edwards JW, and Tappe PA, editors. *Proceedings of the Second Symposium on Southeastern Fox Squirrels, Sciurus niger*. Special Publication Number 1, Virginia Museum of Natural History (Martinsville, VA). p. 53–56.
- Falcón-Ordaz J and L García-Prieto.** 2004. Análisis morfológico de algunos Trichostrongilinos (Strongylida) depositados en La Colección Nacional de Helmintos del Instituto de Biología, UNAM, México. *Revista de Biología Tropical* 52:377–386.
- Flyger V and JE Gates.** 1982. Fox and gray squirrels. In: Chapman JA and GA Feldhamer, editors. *Wild mammals of North America*. Johns Hopkins University Press (Baltimore, MD). p. 209–229.
- Hall MC.** 1916. Nematode parasites of mammals of the orders Rodentia, Lagomorpha and Hyacoidea. *Proceedings of the United States National Museum* 50:1–258.
- Hall TA.** 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41:95–98.
- Hofmannová L, C Romeo, L Štohanzlová, D Jirsová, VM Mazzamuto, LA Wauters, N Ferrari, and D Modrý.** 2016. Diversity and host specificity of coccidia (Apicomplexa: Eimeriidae) in native and introduced squirrel species. *European Journal of Protistology* 56:1–14.
- Joseph T.** 1969. The coccidia of the grey squirrel *Sciurus carolinensis* with descriptions of two new species. *Dissertation Abstracts International (Science and Engineering)* 30:2961.
- Joseph T.** 1972. *Eimeria lancasterensis* Joseph, 1969 and *E. confusa* Joseph, 1969 from the grey squirrel *Sciurus carolinensis*. *Journal of Protozoology* 19:143–150.
- Joseph T.** 1973a. Occurrence of *Eimeria ontarioensis* Lee and Dorney, 1971, in the fox squirrel, *Sciurus niger rufiventer*, in Indiana. *Journal of Parasitology* 59:584–585.
- Joseph T.** 1973b. Eimerians occurring in or infective to both the fox squirrel *Sciurus niger rufiventer* and the gray squirrel *S. carolinensis*. *Journal of Protozoology* 20:509.
- Joseph T.** 1975. Experimental transmission of *Eimeria confusa* Joseph 1969 to the fox squirrel. *Journal of Wildlife Diseases* 11:402–403.
- Katz JS.** 1938. A survey of the parasites found in and on the fox squirrel (*Sciurus niger rufiventer* Geoffrey) and the southern gray squirrel (*Sciurus carolinensis carolinensis* Gmelin) in Ohio. [MS thesis]. The Ohio State University, Columbus, Ohio. 58 p. Available at: https://etd.ohiolink.edu/apexprod/rws_etd/send_file/send?accession=osu1394725814&disposition=inline.
- Kinsella JM.** 1991. Comparison of helminths of three species of mice, *Peromyscus floridanus*, *Peromyscus gossypinus*, and *Peromyscus polionotus*, from southern Florida. *Canadian Journal of Zoology* 69:3078–3083.
- Knipling EF and ER Becker.** 1935. A coccidium from the fox squirrel, *Sciurus niger rufiventer*, Geoffrey. *Journal of Parasitology* 21:417–418.
- Koprowski JL.** 1994. *Sciurus niger*. *Mammalian Species* 479:1–9.
- Langham RF, RL Rausch, and JF Williams.** 1990. Cysticerci of *Taenia mustelae* in the fox squirrel. *Journal of Wildlife Disease* 26:295–296.
- Levine ND and V Ivens.** 1965. The coccidian parasites (Protozoa: Apicomplexa) of rodents. *Illinois Biological Monographs* 33:1–365.
- Lewis RE.** 2000. A taxonomic review of the North American genus *Orchopeas* Jordan, 1933 (Siphonaptera: Ceratophyllidae: Ceratophyllinae). *Journal of Vector Ecology* 25:164–189.

- Lucker JT.** 1943. A new trichostrongylid nematode from the stomachs of American squirrels. *Journal of the Washington Academy of Science* 33:75–79.
- McAllister CT, MB Connior, and LA Durden.** 2013. Ectoparasites of sciurid rodents in Arkansas, including new state records for *Neohaematopinus* spp. (Phthiraptera: Anoplura: Polyplacidae). *Journal of the Arkansas Academy of Science* 67:197–199.
- McAllister CT, LA Durden, and HW Robison.** 2016. The ticks (Arachnida: Acari: Ixodida) of Arkansas. *Journal of the Arkansas Academy of Science* 70:141–154.
- McAllister CT, LA Durden, HW Robison, and MB Connior.** 2017. The fleas (Arthropoda: Insecta: Siphonaptera) of Arkansas. *Journal of the Arkansas Academy of Science* 71:69–76.
- McAllister CT and SJ Upton.** 1989. *Eimeria lancasterensis* (Apicomplexa: Eimeriidae) from the eastern fox squirrel, *Sciurus niger* (Rodentia: Sciuridae), in North-Central Texas. *Journal of Parasitology* 75:642–644.
- McDade JE.** 1987. Flying squirrels and their ectoparasites: disseminators of epidemic typhus. *Parasitology Today* 3:85–87.
- Mollericonn JL, A Nascimento, and R Nallar.** 2016. Primer reporte em Bolivia de *Boehmiella wilsoni* (Nematoda: Strongylida) em *Dasyprocta variegata* (Rodentia: Dasyproctidae). *Ecología en Bolivia* 51:169–174.
- Motriuk-Smith D, RS Seville, CE Oliver, DL Hofmann, and AW Smith.** 2009. Species of *Eimeria* (Apicomplexa: Eimeriidae) from tree squirrels (*Sciuru sniger*) (Rodentia: Sciuridae) and analysis of the ITS1, ITS2, and 5.8S rDNA. *Journal of Parasitology* 95:191–197.
- Nixon CM and LP Hansen.** 1987. Managing forests to maintain populations of gray and fox squirrels. *Illinois Department of Conservation Technical Bulletin* 5:1–35.
- Ozmen O, BA Yukari, and M Haligür.** 2009. First report of *Eimeria lancasterensis* in a red squirrel (*Sciurus vulgaris* L.) in Turkey. *Turkiye Parazitologii Dergisi* 33:245–247.
- Price EW.** 1928. Two new nematode worms from rodents. *Proceedings of the United States National Museum* 74:1–5.
- Rausch R and JD Tiner.** 1948. Studies on the parasitic helminths of the North Central states. I. Helminths of Sciuridae. *American Midland Naturalist* 39:728–747.
- Reiber RJ and EE Byrd.** 1942. Some nematodes from mammals of Reelfoot Lake in Tennessee. *Journal of the Tennessee Academy Science* 17:78–89.
- Sealander JA and GA Heidt.** 1990. Arkansas mammals: Their natural history, classification, and distribution. University of Arkansas Press (Fayetteville, AR). 308 p.
- Schiefer BA and JL Lancaster Jr.** 1970. Some Siphonaptera of Arkansas. *Journal of the Kansas Entomological Society* 43:177–181.
- Walters BL, JO Whitaker Jr, NS Gikas, and WJ Wrenn.** 2011. Host and distribution lists of chiggers (Trombiculidae and Leeuwenhoekiiidae), of North America wild vertebrates north of Mexico. Faculty Publications of the Harold W. Manter Laboratory of Parasitology (Lincoln, NE). Paper 697.183 p.
- Whipps CM, RD Adlard, MS Bryant, RGJ Lester, V Findlay, and ML Kent.** 2003. First report of three *Kudoa* species from Eastern Australia: *Kudoa thyrsites* from Mahi Mahi (*Coryphaena hippurus*), *Kudoa amamiensis* and *Kudoa minithyrsites* n. sp. from Sweeper (*Pempheris ypsilychnus*). *Journal of Eukaryotic Microbiology* 50:215–219.
- Whipps CM, AE Cheeseman, KA Lindsay, and JB Cohen.** 2020. Evaluation of cottontail pellets collected in suboptimal conditions for DNA analysis. *Wildlife Society Bulletin* 44:182–190.
- Whitaker JO Jr, BL Walters, LK Castor, CM Ritzi, and N Wilson.** 2007. Host and distribution lists of mites (Acari), parasitic and phoretic, in the hair or on the skin of North American wild mammals north of Mexico: Records since 1974. Faculty Publications of the Harold W. Manter Laboratory of Parasitology (Lincoln, NE). Paper 1. 17.
- Whitaker JO Jr and RE Mumford.** Mammals of Indiana. Revised and enlarged edition. Indiana University Press (Bloomington, IN). 661 p.
- Whitaker JO Jr, EJ Spika, and V Brack Jr.** 1976. Ectoparasites of squirrels of the genus *Sciurus* from Indiana. *Proceedings of the Indiana Academy of Science* 105:277–280.