



<http://hdl.handle.net/10826/ETD-2012-05-74>

dc.contributor.advisor	Haggard, Brian E.	=LDR 01440njm a2000145a 4500
dc.creator	Bailey, Iain M.	=042 \\\\$adc
dc.date.accessioned	2012-04-13T18:42:28Z	=720 \\\\$aBailey, Iain M.\$eauthor
dc.date.available	2012-04-13T18:42:28Z	=690 \\\\$aHaggard, Brian E.\$econtributor
dc.date.created	2012-05	=690 \\\\$aAgriculture
dc.date.issued	2012-04-13	=260 \\\\$c2012
dc.date.submitted	May 2012	=500 \\\\$a2012
dc.identifier.uri	http://hdl.handle.net/10826/ETD-2012-05-74	=520 \\\\$aThe addition of chemical amendments to land applied poultry litter is paramount to the maintenance of acceptable phosphorus (P) and nitrogen (N) levels in stormwater runoff from pasture. Chitin is the chief component of shrimp shells, and together with its more processed form chitosan, may show promising P, N, and metal ion chelation characteristics by virtue of its chemical structure and unique charge distribution. The objective of this study was to measure the effects of aluminum sulfate (alum, Al ₂ (SO ₄) ₃) on aqueous solubility of P and ammonia (NH ₃) release against those of chitin and three varieties of chitosan in poultry litter. The experiments revealed that while chitin has little to no effect on water extractable P (WEP), all three varieties of chitosan show decreases in WEP not significantly different from those of alum. While alum significantly decreased NH ₃ volatilization from litter samples, neither chitin nor chitosan significantly affected released NH ₃ .
dc.description.abstract	The addition of chemical amendments to land applied poultry litter is paramount to the maintenance of acceptable phosphorus (P) and nitrogen (N) levels in stormwater runoff from pasture. Chitin is the chief component of shrimp shells, and together with its more processed form chitosan, may show promising P, N, and metal ion chelation characteristics by virtue of its chemical structure and unique charge distribution. The objective of this study was to measure the effects of aluminum sulfate (alum, Al ₂ (SO ₄) ₃) on aqueous solubility of P and ammonia (NH ₃) release against those of chitin and three varieties of chitosan in poultry litter. The experiments revealed that while chitin has little to no effect on water extractable P (WEP), all three varieties of chitosan show decreases in WEP not significantly different from those of alum. While alum significantly decreased NH ₃ volatilization from litter samples, neither chitin nor chitosan significantly affected released NH ₃ .	=024 8\ http://hdl.handle.net/10826/ETD-2012-05-74
dc.format.mimetype	application/pdf	=653 \\\\$aBiological and Agricultural Engineering
dc.language.iso	eng	=245 00\$aChitosan reduces water solubility of phosphorus in poultry litter h[electronic resource] / cby Iain M. Bailey
dc.subject	Agriculture	LEADER 00000nam a2200003a 4500
dc.subject	Environmental engineering	005 20120619153015.0
dc.subject	Agriculture engineering	006 m d
dc.title	Chitosan reduces water solubility of phosphorus in poultry litter	007 cr un ---uuuuu
dc.date.updated	2012-04-13T18:42:34Z	008 061912s2012 xx o 000 0 eng d
dc.identifier.slug	10826/ETD-2012-05-74	049 AFUT
thesis.degree.name	Bachelor of Science in Biological Engineering	100 1 Bailey, Iain M
thesis.degree.level	Undergraduate	245 10 Chitosan reduces water solubility of phosphorus in poultry litter h[electronic resource] / cby Iain M. Bailey
thesis.degree.grantor	University of Arkansas, Fayetteville	260 c2012
thesis.degree.discipline	Biological Engineering	300 1 online resource
thesis.degree.department	Biological and Agricultural Engineering	500 "May 2012"
dc.type.material	text	502 Thesis (Honors)--University of Arkansas, Fayetteville, 2012
dc.type.genre	thesis	520 3 The addition of chemical amendments to land applied poultry litter is paramount to the maintenance of acceptable phosphorus (P) and nitrogen (N) levels in stormwater runoff from pasture. Chitin is the chief component of shrimp shells, and together with its more processed form chitosan, may show promising P, N, and metal ion chelation characteristics by virtue of its chemical structure and unique charge distribution. The objective of this study was to measure the effects of aluminum sulfate (alum, Al ₂ (SO ₄) ₃) on aqueous solubility of P and ammonia (NH ₃) release against those of chitin and three varieties of chitosan in poultry litter. The experiments revealed that while chitin has little to no effect on water extractable P (WEP), all three varieties of chitosan show decreases in WEP not significantly different from those of alum. While alum significantly decreased NH ₃ volatilization from litter samples, neither chitin nor chitosan significantly affected released NH ₃ .
dc.contributor.committeeMember	Liang, Yi	538 System Requirements: Internet access, World Wide Web browser, Adobe Acrobat Reader. Mode of access: World Wide Web
dc.contributor.committeeMember	Zaharoff, David A.	655 7 Electronic theses. 2AFU
		690 Dept.: Biological and Agricultural Engineering
		690 Advisor: Haggard, Brian E
		690 Agriculture
		690 Environmental engineering
		690 Agriculture engineering
		830 0 UA undergraduate honors paper
		856 40 uhttp://hdl.handle.net/10826/ETD-2012-05-74

