

5-2018

The Influence of Social Determinants of Health on High School Students' Health Literacy

Rebecca Welch

Follow this and additional works at: <http://scholarworks.uark.edu/nursuht>

 Part of the [Health and Physical Education Commons](#), [Health Services Research Commons](#), and the [Public Health Education and Promotion Commons](#)

Recommended Citation

Welch, Rebecca, "The Influence of Social Determinants of Health on High School Students' Health Literacy" (2018). *The Eleanor Mann School of Nursing Undergraduate Honors Theses*. 61.
<http://scholarworks.uark.edu/nursuht/61>

This Thesis is brought to you for free and open access by the The Eleanor Mann School of Nursing at ScholarWorks@UARK. It has been accepted for inclusion in The Eleanor Mann School of Nursing Undergraduate Honors Theses by an authorized administrator of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, ccmiddle@uark.edu.

The Influence of Social Determinants of Health on High School Students'

Health Literacy

An honors research project submitted in partial fulfillment of the requirements of the degree of Bachelor of Science of Nursing

By
Rebecca Welch

May 2018
The University of Arkansas

Dr. Kelly Vowell Johnson
Research Mentor

Dr. Marilou Shreve, DNP, APRN
Committee Member

Cathy Hale, APRN
Committee Member

This research was supported by a University of Arkansas Honors College Research Grant and an Honors College Conference/Workshop Travel Grant

Abstract

The broad topic of health literacy has been assessed in a variety of studies. The most extensive research has been completed on how a lack of health literacy contributes to specific disease progressions. However, there is a deficit of research on how general health literacy contributes to overall individual health in high school students. The purpose of this study was to evaluate the health literacy scores of Northwest Arkansas high school students and examine the impact that age, gender, and ethnicity have on their health literacy. Health literacy was assessed using the Health Literacy Questionnaire (HLQ). A cross-sectional research design was utilized at a high school in Northwest Arkansas. A total of 100 high school students completed the HLQ. The results indicated no significant correlation between students' age, gender, or ethnicity and their health literacy. The total student population scored lowest on scale 5 of the HLQ, meaning they find it difficult to understand most health information, especially when they receive conflicting messages. Determining the health literacy deficits and identifying social determinants that affect health literacy could help high schools alter their health curriculum to meet student needs.

Background and Significance

Health literacy, known as the ability to access, understand, appraise, and apply health-related knowledge, is a major factor for health promotion and disease prevention (Sorensen, Van den Brouke, Fullam, Doyle, & Brand, 2012). Conversely, low health literacy has been linked to increased hospitalizations, healthcare costs, and mortality. In order to improve the health literacy in Arkansas, it's imperative to first understand what

areas of health literacy are lacking. Therefore, the primary objective of this study was to evaluate the health literacy of high school students, while the secondary objective was to detect any correlating disparities. By assessing these aspects of health literacy, we can better understand where our current education is lacking in health promotion and disease prevention.

To conduct this research, a sample size of 100 high school health students completed the Health Literacy Questionnaire (HLQ). The questionnaire covers all aspects of health literacy and normally takes 30 minutes or less to finish. Nurses and teachers alike will be able to use the results of the study to focus education on student needs. With resulting improved health literacy, students will graduate with improved confidence in their furtherance of well-being.

Literature Review

The broad topic of health literacy has been assessed in a variety of studies. The most extensive research has been completed on how a lack of health literacy contributes to specific disease progressions. However, there is a deficit of research on how general health literacy contributes to overall individual health in high school students. The focus points of this literature review will discuss studies regarding consequences of low literacy rates, disparities in health literacy, and ways that assessing health literacy can improve it.

The literature is clear that health literacy is a determinant of actual health outcomes. Multiple authors have linked low health literacy to increased risk for illness (Dodson, Beauchamp, Batterham, & Osborne 2015; Chari, Warsh, Ketterer, Hossain, & Sharif, 2014). Dodson et al. (2015) expressed the importance of increasing health literacy rates by

pointing out that low health literacy is associated with increased hospital admissions and readmissions, poor medication adherence, increased adverse medication events, increased mortality, lower functional status, and increased healthcare costs. Many of these findings are also related to obesity, which is highly correlated to low health literacy (Chari et al. 2014; Lam & Yang 2012). Obesity is a preventable condition that is a main cause of chronic diseases such as hypertension and diabetes. Chari et al. (2014) described obesity as a large and growing problem which is strongly linked to health literacy. While demonstrating that adolescent health literacy is correlated to adolescent obesity, it is also proven that parental health literacy correlates to their children's obesity. However, by positively influencing the health literacy of adolescents, it will improve their children's weight and health outcomes if the current adolescents so choose to become parents in the future. This is because it is hard to "unlearn" health literacy; it's a skill that will be continually beneficial. Lam and Yang's (2012) also completed a study that deals with obesity and its relation to health literacy. This study population was children in China ages 12-16, and although their studied population differs from our sample, the findings are in agreement with the 2014 study by Chari. Both studies clearly found correlation between low health literacy and being overweight or obese. These findings are significant because the results revealed low health literacy as a causation factor for the most widespread and costly diseases. Now that it is known how improved health literacy can improve health, steps can be taken to make health literacy education a priority.

Identifying low health literacy as a major cause for multiple health issues has provoked recent studies to be conducted for detection of causes and factors that lead to low health literacy (Rikard, Thompson, McKinney, & Beauchamp, 2016; Wilkinson, Haslem,

& Prusak 2016). Both Rikard et al. (2016) and Wilkinson et al. (2016) explored the correlation of gender to health literacy rates; both found females to have overall higher health literacy. Wilkinson's et al. (2016) study focused on high school students, and although it specifically assessed the health-related fitness knowledge of the students instead of overall health literacy, the findings are still applicable. This study was conducted in Washington and has helped schools in the state better educate students on their health-related fitness knowledge.

As stated earlier, the study by Rikard et al. (2016) supported Wilkinson et al. (2016) concerning significant inequality in scores between genders. However, the research of Rikard et al. (2016) used a large national sample size to demonstrate the many other disparities in current health literacy. The disparities that this study explores are age, gender, and ethnicity; these three factors were also included in the national study that surveyed U.S. adults. The results of the National Assessment of Adult Literacy were that minorities, males, and older people have lower health literacy levels than younger people, females, and Caucasians (Rikard et al., 2016). Discovering these disparities which lead to low health literacy can help target focused education towards specific groups or cohorts.

The previously noted studies support, proper integration of health literacy education is a clear necessity. The struggle in teaching health literacy comes when determining the most effective approach. Heo et al. (2016) and MacDonald (2015) both conducted research about the implementation and progress in improving student's physical literacy and health knowledge. HealthCorps, an organization which provides health programs for New York high schools, has been found to increase health literacy in school age children (Heo et al., 2016). The program was found to not only increase student knowledge, but

to improve health promoting behaviors. One reason the New York high school programs are so practical are because they assess their students' needs before beginning education. By using an annual community needs assessment instrument, HealthCorps is able to focus their teachings on the identified lacking components of health (such as nutrition, physical fitness, or mental resiliency). This information further justifies the need for health literacy assessments in Northwest Arkansas schools in order to create effective teaching programs similar to HealthCorps.

MacDonald's study in 2015 differs by solely evaluating physical literacy of high school students. Physical literacy is an important part of overall health literacy, and getting students to plan their improvement goals in physical fitness is a huge accomplishment. As educators support students in making their goals, it is important for them to allow the students to choose their own methods in which to progress their fitness levels. However, if students are physically illiterate or health illiterate, it becomes much more of a challenge to create a realistic and beneficial plan. To further demonstrate how health literacy assessments can improve teaching, Sorensen et al. (2012) established different perceptions and definitions of health literacy. They also proposed an integrative model for health literacy that portrays the most important dimensions of the concept. On an individual level, the model shows that people need to have access to health information, as well as the capacity to understand, appraise, and apply it. Determining whether student needs lie in the accessing, understanding, appraising, or applying elements of health literacy can greatly aid in planning more effective teaching.

Research Design and Methods

This cross-sectional study utilized a single site convenience sample of 100 high school students from a Northwest Arkansas high school. This research was reviewed and approved by the University of Arkansas Institutional Review Board. Consent was obtained through the Office of Research Compliance Institutional Review Board. Passive consent was given by participants two weeks prior to completing the questionnaire. All students took the 44-question survey during their health class and completed it within 30 minutes. A health literacy profile with scores for 9 HLQ scales was generated for each participant.

Instruments

The HLQ consist of 9 rating scales, with a total of 44 questions. The scales are:

1	Feeling understood and supported by healthcare providers (HPS)
2	Having sufficient information to manage my health (HSI)
3	Actively managing my health (AMH)
4	Social support for health (SS)
5	Appraisal of health information (CA)
6	Ability to actively engage with healthcare providers (AE)
7	Navigating the healthcare system (NHS)
8	Ability to find good health information (FHI)
9	Understand health information well enough to know what to do (UHI)

The first five scales are measured in the first 23 questions, which use an agree/disagree format. The options correlating to given statements are: Strongly disagree, Disagree, Agree, Strongly Agree. Scales 4-9 are measured in the last 21 questions by the subjects ranking tasks on how easy or challenging they are to complete. Answer options for the subsequent questions are: Cannot do or always difficult, Usually difficult, Sometimes difficult, Usually easy, Always easy. Demographic information was also obtained on the last page of the HLQ. The HLQ has strong construct validity, reliability, and provides a means to evaluating differences (Osborne, Batterham, Elsworth, Hawkins, & Buchbinder, 2013). There were no identifiers beyond the demographic information (age, gender, and ethnicity) associated with the HLQ, to ensure confidentiality for participants.

Data Analysis

After collecting the data from the Northwest Arkansas High School students, the data was input into an Excel scoring sheet provided with the HLQ. Each participants age, gender, and ethnicity were then coded and added to the spreadsheet to distinguish variables among the disparities.

The supporting Excel scoring sheet was used to calculate the mean scores and standard deviations for each individual participant. Correlations and t-tests were then performed to further analyze the data.

The HLQ scale averages weren't directly compared due to the first 5 scales were one to four and the last four scales were one to five. Therefore, to rank the mean scores on each HLQ scale, the mean scores were divided by the highest possible score for that scale to come up with comparison scores.

Participant Demographics

A total of 100 High School students completed the HLQ for this study. All participants completed the consent process. The participants were all high school students attending a high school in Northwest Arkansas and enrolled in a health course. The HLQ was taken on December 6th, 2017. Participants were predominately Hispanic (53%), followed by Pacific Islanders (18%). Other ethnicities included Caucasian (14%), African American (4%), Native American (2%), and Other (9%). The student population included 41 females and 56 males ranging in age from 15 to 19 years old. A small number of participants did not mark gender (.03), age (.04) or ethnicity (0.5).

Results

The comparison scores were analyzed to determine the highest and lowest priority aspects of health literacy that needed improvement. The lowest scored scale was HLQ's scale five, which measured the ability to appraise health information. Students indicated they have the needed support to maintain good health by scoring highest on scale four. Age, gender, or ethnicity were not significantly correlated to health literacy in high school students in this study.

	Std.	.51131	.69494	.50170	.49013	.4548	.57588	.78079	.81311	.63734	.73407
	Deviati on	054945 0471		046272 4217	490041 9853		198274 9202	264551 9624	962343 3464	755142 4057	623890 7310
3	Mean	3.6818 181818 18182	3.5625	3.4375 000000 00000	3.1500 000000 00000	3.500	3.0000 000000 00000	4.2000 000000 00000	4.1250 000000 00000	3.8500 000000 00000	4.1500 000000 00000
	N	4	4	4	4	4	4	4	4	4	4
	Std.	.17108	.42696	.51538	.44347	.1155	.28284	.28284	.15957	.19148	.19148
	Deviati on	469379 7518		820320 2208	115652 1669		271247 4619	271247 4619	118462 6056	542155 1267	542155 1268
4	Mean	3.4431 818181 81818	3.2500	3.0000 000000 00000	3.5000 000000 00000	3.200	2.4000 000000 00000	3.5000 000000 00000	4.0000 000000 00000	4.0000 000000 00000	3.9000 000000 00000
	N	2	2	2	2	2	2	2	2	2	2
	Std.	.08035	.70711	.00000	.14142	.2828	.00000	.14142	.00000	.00000	.14142
	Deviati on	304331 6653		000000 0000	135623 7310		000000 0000	135623 7310	000000 0000	000000 0000	135623 7310
5	Mean	3.1518 582396 79537	2.5395	2.8289 473684 21052	3.0526 315789 47368	3.284	2.7052 631578 94736	3.3684 210526 31579	3.3859 649122 80702	3.4315 789473 68422	3.5368 421052 63158
	N	19	19	19	19	19	19	19	19	19	19
	Std.	.44699	.76948	.64578	.55714	.3962	.52225	.85443	.60362	.68722	.56195
	Deviati on	907280 7379		027512 6680	392819 5530		332742 9990	459640 4057	237699 5002	681972 8673	148694 9016
6	Mean	3.1590 909090 90909	2.5000	3.0000 000000 00000	3.1333 333333 33333	3.133	2.9333 333333 33333	3.2666 666666 66667	3.3888 888888 88889	3.4666 666666 66667	3.4000 000000 00000
	N	3	3	3	3	3	3	3	3	3	3
	Std.	.38836	.50000	.25000	.61101	.2309	.41633	.61101	.67357	.23094	.34641
	Deviati on	380660 5342		000000 0000	009266 0779		319989 3226	009266 0779	531405 4564	010767 5850	016151 3776
To tal	Mean	3.1770 168020 47401	2.6789	2.8745 614035 08773	2.8000 000000 00000	3.116	2.5378 947368 42106	3.5515 789473 68421	3.5228 070175 43861	3.6610 526315 78948	3.6189 473684 21052
	N	95	95	95	95	95	95	95	95	95	95
	Std.	.50594	.74970	.56268	.55120	.5133	.56270	.80356	.75465	1.0647	.66465
	Deviati on	082708 9674		859252 5889	757182 2125		224683 1329	428834 8803	870916 9991	439745 04200	392320 0623

	N	96	100	100	100	100	100	100	100	100	100	100
Scale 4 (SS)	Pearson Correlation	.005	.549**	.392**	.459**	.337**	1	.297**	.402**	.281**	.321**	.297**
	Sig. (2-tailed)	.963	.000	.000	.000	.001		.003	.000	.005	.001	.003
	N	96	100	100	100	100	100	100	100	100	100	100
Scale 5 (CA)	Pearson Correlation	-.050	.682**	.498**	.471**	.529**	.297**	1	.444**	.451**	.423**	.430**
	Sig. (2-tailed)	.631	.000	.000	.000	.000	.003		.000	.000	.000	.000
	N	96	100	100	100	100	100	100	100	100	100	100
Scale 6 (AE)	Pearson Correlation	-.116	.830**	.646**	.453**	.361**	.402**	.444**	1	.778**	.465**	.657**
	Sig. (2-tailed)	.259	.000	.000	.000	.000	.000	.000		.000	.000	.000
	N	96	100	100	100	100	100	100	100	100	100	100
Scale 7 (NHS)	Pearson Correlation	-.066	.827**	.636**	.477**	.424**	.281**	.451**	.778**	1	.444**	.665**
	Sig. (2-tailed)	.524	.000	.000	.000	.000	.005	.000	.000		.000	.000
	N	96	100	100	100	100	100	100	100	100	100	100
Scale 8 (FHI)	Pearson Correlation	-.042	.692**	.430**	.434**	.147	.321**	.423**	.465**	.444**	1	.438**
	Sig. (2-tailed)	.686	.000	.000	.000	.143	.001	.000	.000	.000		.000
	N	96	100	100	100	100	100	100	100	100	100	100
Scale 9 (UHI)	Pearson Correlation	-.069	.771**	.564**	.472**	.395**	.297**	.430**	.657**	.665**	.438**	1
	Sig. (2-tailed)	.503	.000	.000	.000	.000	.003	.000	.000	.000	.000	
	N	96	100	100	100	100	100	100	100	100	100	100

** . Correlation is significant at the 0.01 level (2-tailed).

Age was not an indicator of health literacy as shown by the large (greater than .05) p-values in the ‘age’ row (Table 3). By finding no significant values and no substantial effect size, it is apparent the sample size was sufficient in concluding that high schoolers’ age does not impact health literacy scores. These results may be due to the limited variance (4 years) in the ages surveyed.

Table 4: Gender analysis (Sex 1=females; Sex 2=males)

Group Statistics					
	Sex	N	Mean	Std. Deviation	Std. Error Mean
Total	1	41	3.2000592997473	.50523523874090	.07890448787285
			31	5	5
	2	56	3.1607237239504	.50118759179095	.06697400910657
			68	6	4
Scale 1 (HPS)	1	41	2.7622	.77248	.12064
	2	56	2.6339	.72608	.09703
Scale 2 (HSI)	1	41	2.8861788617886	.50406545043256	.07872179763212
			18	4	8
	2	56	2.8571428571428	.59897097906846	.08004086386766
			57	4	9
Scale 3 (AMH)	1	41	2.7902439024390	.52905806772450	.08262498869405
			24	8	3
	2	56	2.8285714285714	.57768008942171	.07719574906346
			30	9	5
Scale 4 (SS)	1	41	3.132	.5415	.0846
	2	56	3.082	.4999	.0668
Scale 5 (CA)	1	41	2.5365853658536	.53560997823088	.08364822520539
			58	4	2
	2	56	2.5446428571428	.58025688444692	.07754008778276
			57	5	0
Scale 6 (AE)	1	41	3.6292682926829	.80568104807021	.12582624015948
			27	0	0
	2	56	3.5000000000000	.79085684257932	.10568269096134
			01	9	5
Scale 7 (NHS)	1	41	3.5487804878048	.80815625417694	.12621280240861
			79	6	8

	2	56	3.5178571428571	.70972933482823	.09484157172393
			42	5	6
Scale 8 (FHI)	1	41	3.6195121951219	.59633024080417	.09313113703432
			51	1	6
	2	56	3.6821428571428	1.3010635010506	.17386192354886
			58	73	5
Scale 9 (UHI)	1	41	3.6731707317073	.66446384034193	.10377181758515
			17	4	0
	2	56	3.5607142857142	.66433249381440	.08877516366980
			85	2	8

Table 4 shows the score averages between the 41 females and 56 males who were surveyed. There was no significant difference in scores by gender in any of the 9 scales. The mean scores of each gender per scale were very close, meaning that gender is not a disparity for health literacy of high school students in this study.

Discussion

The current level of health literacy among high school students in this study demonstrates a need for additional support and education to better evaluate health materials. Since this Northwest Arkansas study did not find any disparities, educational measures should be aimed at the entire student population. The HLQ provides information from 9 scales which create a health literacy profile for each student. These results can be averaged to create a health literacy profile of the school to supply the most insight when interpreting strengths and weaknesses in health literacy of the total student population. The lowest scores were on the scale concerning students' abilities to evaluate the accuracy of health information.

High school students need improvement in their ability to appraise health information (O'Dell, 2012). Since students get so many of their facts from the internet, it

is alarming that studies have concluded high-schoolers are unable to differentiate accurate from inaccurate websites (Kortum, Edwards, & Richards-Kortum, 2018). This is especially troubling since high school students are one of the most likely populations to utilize technology to self-diagnose and self-treat. The consequences of implementing inaccurate health information can jeopardize patient safety and lifelong health outcomes. With the abundance of conflicting messages on health advertised in the media, it is no surprise that the ability to appraise health information is a challenge for students. This is also the reason that more education should be given in high schools on how to differentiate poor sources and false information from those that are reliable and high quality.

Educating children and adolescents on health literacy, and specifically health information appraisal, should be done in a way that engages the students and encourages critical thinking. It's important for students to realize their own beliefs about health before they can further examine their established knowledge and evaluate the materials either supporting or contrasting their initial theories. It's also valuable to present students with multidimensional or contrasting health messages during teaching sessions to prepare them for assessing the validness of messages when they leave the classroom (Fairbrother, Curtis, & Goyder, 2016).

Limitations

The study had several limitations which should be considered. The most significant limitation faced in this study was the restricted sample size. The small

population of health students sampled delivered inconclusive results for the correlation of ethnicity to health literacy.

Another limitation of the study is that it is based on self-reported data, and therefore may have inaccurate data. The use of rating scales could have impacted study results due to students having diverse styles of responding to this type of questionnaire; some people frequent extreme answers such as ‘strongly agree’ or ‘strongly disagree’, while others stick to more impartial choices (Hoskin, 2012).

Despite the questionnaire being proven highly valid and reliable, the participants may have rushed through the questionnaire, interpreted statements differently, or not fully comprehended the questions. All surveys were given in English despite many students not speaking English as their first language. It is unknown if this affected the questionnaire results.

Social desirability bias may have also affected the results of this study because the participants knew the purpose of the study was to determine their health literacy. However, the investigators encouraged participants to respond honestly and reinforced the confidentiality of the results.

Lastly, the site of the sampling may have delivered results that don’t accurately reflect the health literacy or demographics of other Northwest Arkansas High Schools. The school used in this study had a higher number of Hispanic and Marshallese students than many other Northwest Arkansas schools. Further research on Health Literacy in other areas of Arkansas and other states should be done in the future to compare with this study.

Conclusions

Age, gender, and ethnicity were not shown to be positively correlated with students' health literacy scores in this study. Overall, students scored the lowest in their ability to evaluate the reliability of health information. The results from this study indicate that education is needed for high school students in appraising health information. Further research is necessary to fully understand the health literacy dynamics in Northwest Arkansas high school students. Determining the health literacy needs of high school students and how social determinants of health can affect health literacy is important in supporting and educating high school students. Future studies using the HLQ should implement larger sample sizes with a wider age range to get more accurate results on possible disparities.

References

- Chari, R., Warsh, J., Ketterer, T., Hossain, J., & Sharif, I. (2014). Association between health literacy and child and adolescent obesity. *Patient Education and Counseling, 94*(1), 61-66.
- Dodson S, Beauchamp A, Batterham RW and Osborne RH. Information sheet 10: About the Health Literacy Questionnaire (HLQ). In Ophelia Toolkit: A step-by-step guide for identifying and responding to health literacy needs within local communities. 2014. Retrieved from www.ophelia.net.au
- Dodson, S., Beauchamp, A., Batterham, R., & Osborne, R. (2015). Health literacy, inequity, and health outcomes. *Belges De Santé Publique, 72*(1), 11.
- Fairbrother H, Curtis P, Goyder E. (2016). Making health information meaningful: Children's health literacy practices. *SSM - Population Health. 2*:476–84. <https://doi.org/10.1016/j.ssmph.2016.06.005>
- Heo, M., Irvin, E., Ostrovsky, N., Isasi, C., Blank, A. E., Lounsbury, D. W., . . . Wylie-Rosett, J. (2016). Behaviors and knowledge of HealthCorps New York city high school students: Nutrition, mental health, and physical activity. *Journal of School Health, 86*(2), 84-95.
- Hoskin, R. (2012, March 3). The dangers of self-report. Science Brainwaves. Retrieved from <http://www.sciencebrainwaves.com/the-dangers-of-self-report/>
- Kortum, P., Edwards, C. and Richards-Kortum, R. (2008). The impact of inaccurate Internet health information in a secondary school learning environment. *Journal of Medical Internet Research, 10*(2). doi: 10.2196/jmir.986

- Lam, L. T., & Yang, L. (2014). Is low health literacy associated with overweight and obesity in adolescents: An epidemiology study in a 12-16years old population, Nanning, China, 2012. *Archives of Public Health = Archives Belges De Santé Publique*, 72(1), 11.
- MacDonald, L. C. (2015). Moving high school students toward physical literacy. *Journal of Physical Education, Recreation & Dance*, 86(7), 23-27.
- O'Dell, R. (2012). Appraisal skills, health literacy and the patient-provider relationship: considerations as the health care consumer turns to the internet to inform their care. *Online Journal of Public Health Informatics*, 4(1), ojphi.v4i1.3684.
<http://doi.org/10.5210/ojphi.v4i1.3684>
- Osborne RH, Batterham R, Elsworth GR, Hawkins M, & Buchbinder, R. The grounded theory, psychometric development and initial validation of the health literacy Questionnaire (HLQ). *BMC Public Health*. 2013;13:658.
- Rikard, R. V., Thompson, M. S., McKinney, J., & Beauchamp, A. (2016). Examining health literacy disparities in the United States: a third look at the National Assessment of Adult Literacy (NAAL). *BMC Public Health*, 16(1), 975.
<http://doi.org/10.1186/s12889-016-3621-9>
- Sørensen, K., Van den Broucke, S., Fullam, J., Doyle, G., Pelikan, J., ... Brand, H. (2012). Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health*, 12(1). doi:10.1186/1471-2458-12-80
- Wilkinson, C., Haslem, L., & Prusak, K. (2016). Students' health-related fitness knowledge, perceived competence, and physical activity. *Research Quarterly for Exercise and Sport*, 87(S2), A57.