

2016

## Gastrointestinal Parasites of the Northern River Otter, *Lontra canadensis* (Carnivora: Mustelidae), from Arkansas

C. T. McAllister

*Eastern Oklahoma State College, cmcallister@se.edu*

A. Surf

*Henderson State University*

R. Tumlison

*Henderson State University*

C. R. Burse

*Pennsylvania State University-Shenango*

T. J. Fayton

*University of Southern Mississippi*

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Population Biology Commons](#), and the [Zoology Commons](#)

---

### Recommended Citation

McAllister, C. T.; Surf, A.; Tumlison, R.; Burse, C. R.; and Fayton, T. J. (2016) "Gastrointestinal Parasites of the Northern River Otter, *Lontra canadensis* (Carnivora: Mustelidae), from Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 70, Article 49. Available at: <http://scholarworks.uark.edu/jaas/vol70/iss1/49>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact [scholar@uark.edu](mailto:scholar@uark.edu), [cmiddle@uark.edu](mailto:cmiddle@uark.edu).

## Gastrointestinal Parasites of the Northern River Otter, *Lontra canadensis* (Carnivora: Mustelidae), from Arkansas

C.T. McAllister<sup>1\*</sup>, A. Surf<sup>2</sup>, R. Tumilson<sup>2</sup>, C.R. Bursey<sup>3</sup>, and T.J. Fayton<sup>4</sup>

<sup>1</sup>Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745, <sup>2</sup>Department of Biology, Henderson State University, Arkadelphia, AR 71999, <sup>3</sup>Department of Biology, Pennsylvania State University-Shenango, Sharon, PA 68421  
<sup>4</sup>Gulf Coast Research Laboratory, University of Southern Mississippi, Ocean Springs, MS 39564

\*Correspondence: cmcallister@se.edu

Running Title: Helminths of River Otter from Arkansas

The northern river otter, *Lontra canadensis* (Schreber) ranges from Alaska and most of Canada south to northern California and northern Utah in the west and from Newfoundland southward to Florida in the east (Reid 2006). In Arkansas, *L. canadensis* is found statewide in rivers, creeks, bayous, and lakes bordered by timber (Sealander and Heidt 1990). The river otter primarily feeds on fish, but will also consume aquatic invertebrates, amphibians, reptiles, birds, and other mammals (Tumilson et al. 1986, Larivière and Walton 1998, Melquist et al. 2003, Ligon and Reasor 2007).

The river otter has been a common subject of several surveys on its helminth parasites. These previous studies were conducted on specimens from 15 US states, including Alabama, Alaska, Arkansas, Florida, Georgia, Louisiana, Maryland, Massachusetts, Michigan, Montana, New York, North Carolina, Oklahoma, Oregon, and Tennessee, and Newfoundland and Ontario, Canada (summarized by Fleming et al. 1977, Stuht 1978, Shoop and Corkum 1981, Tumilson et al. 1984, Addison et al. 1988, Snyder et al. 1989, Forrester 1992, Hoberg et al. 1997, Kollars et al. 1997, Kimber and Kollias 2000, Feldhamer et al. 2003, Dronen 2009, Crait et al. 2015, McAllister and Fayton 2015). Although it appears the helminth fauna of *L. canadensis* is fairly well known, there are obvious gaps in our knowledge, particularly for populations in Arkansas. Indeed, only a single nematode parasite has been reported from river otters from Arkansas (Tumilson et al. 1986). Here, we report more detailed information on a survey of gastrointestinal helminth parasites of *L. canadensis* from the state.

During the 2013–2015 fur trapping seasons, 38 adult river otter carcasses (21 males, 17 females) were obtained from licensed fur buyers in Arkansas ( $n = 2$ ), Clark ( $n = 1$ ), Crawford ( $n = 5$ ), Franklin ( $n = 2$ ), Lonoke ( $n = 1$ ), Pike ( $n = 4$ ), Polk ( $n = 5$ ), Perry ( $n = 1$ ), Prairie ( $n = 9$ ), Pulaski ( $n = 1$ ), Scott ( $n = 1$ ), St. Francis

( $n = 1$ ), Sebastian ( $n = 2$ ), Union ( $n = 1$ ), White ( $n = 1$ ) and Yell ( $n = 1$ ) counties (Fig. 1). Most of the carcasses were obtained frozen, but many had remained at ambient temperatures for 24 hrs or more before being frozen. These conditions may have limited the diversity of parasites we were able to locate. After thawing, a mid-ventral incision was made to expose the gastrointestinal tract, which was split lengthwise from the esophagus to anus and its contents rinsed into a sieve then examined for helminths. Several 15 cm segments of tissue were cut and placed in Petri dishes and their contents rinsed in 0.9% w/v saline. Trematodes were removed from the stomach and upper small intestine, transferred to 70% v/v ethanol, stained in acetocarmine or Ehrlich's hematoxylin, cleared in methyl salicylate and mounted in Canada balsam. Nematodes and acanthocephalans originally fixed in 70% v/v ethanol were examined as temporary mounts in glycerol. Host voucher specimens (skulls only) were deposited in the Henderson State University Collection, Arkadelphia, as HSU 719, 763, 784, 809, 867, 870; parasite vouchers were deposited in the Harold W. Manter Laboratory of




Fig. 1. Sixteen counties in Arkansas where river otters were collected.

## Helminths of River Otter from Arkansas

Parasitology (HWML), University of Nebraska, Lincoln, NE.

A total of 37 of 38 (97%) of the *L. canadensis* harbored at least 1 helminth; 1 (3%) of the river otters from Polk County harbored a multiple infection of 2 helminths. Two specimens positive for the nematode, *Dracunculus insignis* (Leidy, 1858) will be reported elsewhere by RT. An annotated list of the parasites found follows:

### **Trematoda: Digenea: Echinostomatidae, *Bashkirovitrema canadense* Dronen, 2009**

Twenty-four of 38 (63%) of the river otters were found to be infected with *B. canadense* (HWML 94121) in the gastrointestinal tract; the mean intensity was  $8.4 \pm 18.2$ , range 1–61 worms. Dronen (2009) erected *B. canadense* to accommodate specimens of *Bashkirovitrema* Skrjabin, 1944 harvested from the intestine of mustelids, including *L. canadensis* and the American mink, *Neovision vision* (Schreber), from Florida, Georgia, Louisiana, New York, North Carolina, and Ontario, Canada (Dronen 2009) and found this species to be morphologically most similar to *Bashkirovitrema incrassatum* (Diesing, 1850), which parasitizes the Neotropical otter, *Lontra longicaudis* (Olfers, 1818), in Brazil, South America. Recently, McAllister et al. (2015) reported *B. canadensis* from Oklahoma; we report this species from Arkansas for the first time.

### **Strigeidida: Clinostomatidae, *Clinostomum* sp.**

Seven *Clinostomum* sp. (HWML 94120) were found in the stomach of a single (3%) *L. canadensis* collected on 16 February 2015 from the Little Missouri River, Clark County. This parasite (yellow grub) is typically found in the metacercarial stage in fishes and amphibians with fish-eating birds (herons, egrets) serving as definitive hosts (McAllister et al. 2010). In the current case, this parasite is considered an incidental finding of a host which has a piscivorous diet. However, this is the first time *Clinostomum* sp. has been reported from a river otter.

### **Nematoda: Dioctophymatoidea: Dioctophymatidae, *Dioctophyma renale* (Goeze, 1782)**

Twelve of 38 (32%) of *L. canadensis* from Crawford, Franklin, Pike, Polk, Prairie, Pulaski, Sebastian, St. Francis and Yell counties harbored larval and pre-adults of this nematode (HWML 94122) in the large intestine; the mean intensity was  $3.8 \pm 9.0$ , range 1–31 worms. This cosmopolitan nematode, often referred to as the giant kidney worm has been found

encapsulated in fishes; larvae can be transmitted along the food chain of paratenic hosts (Karmanova 1961). It is possible that American bullfrogs (*Lithobates catesbeianus*) become infected with *D. renale* by eating infected frogs of different species (Mace and Anderson 1975). Definitive hosts are carnivorous mammals, notably mink, wolves, coyotes, foxes, dogs, raccoons, and weasels. *Dioctophyma renale* has been previously reported from Arkansas (Hallberg 1953) and from *L. canadensis* from unknown localities (Kimber et al. 2000). However, we report *D. renale* in river otters from Arkansas for the first time.

### **Acanthocephala gen sp. (unknown genus)**

Three female acanthocephalans (HWML 94123) were found in the large intestine of 3 of 38 (8%) river otters collected in February 2014, 2 hosts from Prairie and 1 host from Scott counties. As with the *Clinostomum* sp. found herein, this is considered to be an incidental finding of river otter food habits. Other acanthocephalans, also considered incidental findings, have been previously reported in *L. canadensis* from Alabama, Alaska, Florida, Oregon, Tennessee, Wyoming and Newfoundland, Canada (Schmidt 1969, Smith and Threlfall 1973, Fleming et al. 1977, Forrester 1992, Hoberg et al. 1997, Kollars et al. 1997, Crait et al. 2015).

In summary, we document a low parasite species richness as well a new distributional record for a trematode parasite of *L. canadensis* in Arkansas. The parasite list for river otters is much more extensive (see Kimber and Kollias 2000 for summary). Therefore, in order to further increase our knowledge of its parasites in the state, we suggest future surveys to include examination of fresh material, their major visceral organs, and specimens from other counties of the state, particularly northern Arkansas.

### **Acknowledgments**

We thank the trappers who provided otter carcasses and Dr. S.L. Gardner (HWML) for expert curatorial assistance. The Arkansas Game and Fish Commission issued a Scientific Collecting Permit to RT.

### **Literature Cited**

Addison EM, MA Strickland, AB Stephenson and J Hoeve. 1988. Cranial lesions possibly associated with *Skrjabingylus* (Nematoda: Metastrongyloidea) infections in martens, fishers, and otters. Canadian Journal of Zoology 66:2155–2159.

- Crait JR, AD McIntosh, EC Greiner and M Ben-David.** 2015. The influence of changing prey availability on the prevalence of *Diphyllbothrium* in river otters from Yellowstone National Park. *Journal of Parasitology* 101:240–243.
- Dronen ND.** 2009. Re-evaluation of the species composition of *Bashkirovitrema* Skrjabin, 1944 (Digenea: Echinostomatidae), with the description of two new species of this genus and the proposal of *Kostadinovatrema novaeguiniense* n. g., n. sp. *Systematic Parasitology* 74:169–185.
- Feldhamer GA, BC Thompson and JA Chapman (eds).** 2003. Wild mammals of North America: biology, management, and conservation. 2<sup>nd</sup> edition. The John Hopkins University Press (Baltimore). 1225 p.
- Fleming WJ, CF Dixon and JW Lovett.** 1977. Helminth parasites of river otters (*Lutra canadensis*) from southeastern Alabama. *Proceedings of the Helminthological Society of Washington* 44:131–35.
- Forrester DJ.** 1992. Parasites and diseases of wild mammals in Florida. University of Florida Press (Gainesville). 459 p.
- Hallberg CW.** 1953. *Diocetophyma renale* (Goeze, 1782) a study of the migration routes to the kidneys of mammals and resultant pathology. *Transactions of the American Microscopical Society* 72:351–363.
- Hoberg EP, CJ Henry, OR Hedstrom and RA Grove.** 1997. Intestinal helminths of river otters (*Lutra canadensis*) from the Pacific Northwest. *Journal of Parasitology* 83:105–110.
- Karmanova EM.** 1961. The first report of *Diocetophyma renale* in fish. *Akademi Nauka SSSR* 11:118–121.
- Kimber KR and GV Kollias II.** 2000. Infectious and parasitic diseases and contaminant-related problems of North American river otters (*Lontra canadensis*): a review. *Journal of Zoo and Wildlife Medicine* 31:452–472.
- Kollars TM, RE Lizotte and WE Wilhelm.** 1997. Gastrointestinal helminths in the river otter (*Lutra canadensis*) in Tennessee. *Journal of Parasitology* 83:158–160.
- Larivière S and LR Walton.** 1998. *Lontra canadensis*. *Mammalian Species* 587:1–8.
- Ligon DB and J Reasor.** 2007. Predation on alligator snapping turtles (*Macrochelys temminckii*) by northern river otters (*Lontra canadensis*). *Southwestern Naturalist* 52:608–610.
- Mace TF and RC Anderson.** 1975. Development of the giant kidney worm, *Diocetophyma renale* (Goeze, 1782) (Nematoda: Diocetophymatoidea). *Canadian Journal of Zoology* 53:1552–1568.
- McAllister CT and TJ Fayton.** 2015. *Bashkirovitrema canadense* (Trematoda: Digenea: Echinostomatidae) from northern river otter, *Lontra canadensis* (Carnivora: Mustelidae), from southeastern Oklahoma. *Proceedings of the Oklahoma Academy of Science* 95:67–70.
- McAllister CT, CR Bursey, JA Crawford, AR Kuhns, C Shaffer and SE Trauth.** 2010. Metacercariae of *Clinostomum* (Trematoda: Digenea) from three species of *Ambystoma* (Caudata: Ambystomatidae) from Arkansas and Illinois, U.S.A. *Comparative Parasitology* 77:25–30.
- Melquist WE, PJ Polechla Jr and D Toweill.** 2003. River otter: *Lontra canadensis*. In Feldhamer GA, Thompson BC and Chapman JA, editors. Wild mammals of North America: biology, management, and conservation. Johns Hopkins University Press (Baltimore). p 708–734.
- Reid FA.** 2006. Mammals of North America. Houghton Mifflin Company (Boston). 579 p.
- Sealander JA and GA Heidt.** 1990. Arkansas mammals: their natural history, classification, and distribution. University of Arkansas Press (Fayetteville). 308 p.
- Schmidt GD.** 1979. *Paracanthocephalus rauschi* sp. n. (Acanthocephala: Paracanthocephalidae) from grayling, *Thymallus arcticus* (Pallas) in Alaska. *Canadian Journal of Zoology* 47:383–385.
- Shoop WL and KC Corkum.** 1981. Some trematodes of mammals in Louisiana, USA. *Tulane University Studies in Zoology and Botany* 22:109–121.
- Smith FR and W Threlfall.** 1973. Helminths of some mammals from Newfoundland. *American Midland Naturalist* 90:215–218.
- Snyder DE, AN Hamir, VF Nettles and CE Rupprecht.** 1989. *Dirofilaria immitis* in a river otter (*Lutra canadensis*) from Louisiana. *Journal of Wildlife Diseases* 25:629.
- Stuht JN.** 1978. Paragonomiasis in a river otter. State of Michigan Department of Natural Resources Wildlife Division Report Number 2824, Case Report (WPL No. 74-741). Michigan State University, Lansing, Michigan.
- Tumilson R, R Smith, J Hudspeth, PJ Polechia and AW King.** 1984. Dracunculiasis in some Arkansas carnivores. *Journal of Parasitology* 70:440.
- Tumilson R, AW King and VR McDaniel.** 1986. The river otter in Arkansas. IV. Winter food habits in eastern Arkansas. *Proceedings of the Arkansas Academy of Science* 40:76–77.